



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2024 – 02:49 PM EDT

PDB ID : 6BMN  
Title : Structure of human DHHC20 palmitoyltransferase, space group P63  
Authors : Rana, M.S.; Lee, C.-J.; Banerjee, A.  
Deposited on : 2017-11-15  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	<div> <div>25%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
2	B	295	<div> <div>14%</div> <div>92%</div> <div>7%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called human DHHC20 palmitoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total	C	N	O	S	0	0	0
			2332	1551	371	393	17			

- Molecule 2 is a protein called human DHHC20 palmitoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	294	Total	C	N	O	S	0	0	0
			2405	1598	381	407	19			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	THR	-	expression tag	UNP Q5W0Z9

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is 3'-PHOSPHATE-ADENOSINE-5'-DIPHOSPHATE (three-letter code: PAP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	O	0	0
			2	2		
6	B	23	Total	O	0	0
			23	23		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: human DHHC20 palmitoyltransferase



- Molecule 2: human DHHC20 palmitoyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.54Å 99.54Å 158.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	86.20 – 2.25 86.20 – 2.25	Depositor EDS
% Data completeness (in resolution range)	94.8 (86.20-2.25) 79.9 (86.20-2.25)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.20 (at 2.25Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, $R_{free}$	0.257 , 0.278 0.263 , 0.281	Depositor DCC
$R_{free}$ test set	38057 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.8	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 62.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.063 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4812	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CME, ZN, PAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/2409	0.38	0/3274
2	B	0.25	0/2476	0.39	0/3364
All	All	0.25	0/4885	0.38	0/6638

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2332	0	2242	12	0
2	B	2405	0	2321	13	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	10	0	0	0	0
4	B	5	0	0	1	0
5	B	31	0	11	6	0
6	A	2	0	0	0	0
6	B	23	0	0	0	0
All	All	4812	0	4574	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:403:PAP:O4'	5:B:403:PAP:C1'	1.66	1.24
2:B:126:ARG:NH1	4:B:404:PO4:O4	2.30	0.64
5:B:403:PAP:H5'2	5:B:403:PAP:C8	2.29	0.62
2:B:156:CME:HB2	2:B:159:VAL:HG22	1.81	0.61
1:A:88:ASN:HA	1:A:91:LYS:HE2	1.84	0.59
1:A:189:VAL:O	1:A:193:PHE:N	2.35	0.58
2:B:143:SER:OG	5:B:403:PAP:N6	2.30	0.56
1:A:39:VAL:O	1:A:43:PHE:HB3	2.08	0.54
5:B:403:PAP:H5'2	5:B:403:PAP:H8	1.90	0.53
2:B:154:HIS:NE2	2:B:156:CME:SD	2.78	0.52
2:B:69:TYR:CZ	2:B:73:ILE:HD11	2.45	0.51
1:A:5:THR:OG1	1:A:6:LEU:N	2.45	0.49
1:A:239:ARG:NE	1:A:256:GLY:O	2.46	0.49
1:A:69:TYR:CZ	1:A:73:ILE:HD11	2.48	0.49
1:A:169:LYS:NZ	1:A:282:ILE:O	2.38	0.47
5:B:403:PAP:H8	5:B:403:PAP:O3B	2.14	0.47
1:A:126:ARG:NH2	1:A:139:ALA:O	2.46	0.46
1:A:203:ASP:O	1:A:207:LYS:HG2	2.15	0.46
2:B:37:LEU:O	2:B:42:ILE:HG13	2.16	0.46
1:A:149:ILE:HB	1:A:152:MET:HB2	2.00	0.43
2:B:43:PHE:HE1	2:B:48:ASN:HB2	1.83	0.43
1:A:37:LEU:O	1:A:42:ILE:HG13	2.19	0.42
2:B:265:LYS:O	2:B:269:GLN:HG2	2.19	0.42
2:B:275:LYS:HG2	2:B:278:TRP:CH2	2.53	0.42
2:B:156:CME:HE2	2:B:158:TRP:HD1	1.85	0.42
1:A:43:PHE:CE1	2:B:276:LYS:HB2	2.56	0.41
2:B:276:LYS:HE3	2:B:277:TYR:CE2	2.56	0.41
5:B:403:PAP:C8	5:B:403:PAP:O3B	2.68	0.41
2:B:293:PRO:HA	2:B:299:MET:HG2	2.01	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/295 (97%)	281 (99%)	4 (1%)	0	100	100
2	B	291/295 (99%)	285 (98%)	6 (2%)	0	100	100
All	All	576/590 (98%)	566 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/268 (92%)	245 (99%)	3 (1%)	67	76
2	B	258/267 (97%)	256 (99%)	2 (1%)	79	86
All	All	506/535 (95%)	501 (99%)	5 (1%)	73	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	THR
1	A	162	CYS
1	A	215	PHE
2	B	162	CYS
2	B	297	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CME	B	156	2	8,9,10	1.77	1 (12%)	6,9,11	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CME	B	156	2	-	4/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	156	CME	O-C	4.23	1.36	1.20

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	156	CME	CE-SD-SG-CB
2	B	156	CME	N-CA-CB-SG
2	B	156	CME	SD-CE-CZ-OH
2	B	156	CME	CZ-CE-SD-SG

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	156	CME	3	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PO4	B	404	-	4,4,4	0.98	0	6,6,6	0.48	0
4	PO4	A	403	-	4,4,4	0.95	0	6,6,6	0.46	0
4	PO4	A	404	-	4,4,4	0.97	0	6,6,6	0.46	0
5	PAP	B	403	-	28,33,33	4.62	12 (42%)	37,52,52	1.47	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PAP	B	403	-	-	10/17/37/37	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	403	PAP	O4'-C1'	19.52	1.66	1.40
5	B	403	PAP	P-O3'	7.02	1.71	1.59
5	B	403	PAP	O4'-C4'	-5.78	1.32	1.45
5	B	403	PAP	PA-O3A	5.42	1.65	1.59
5	B	403	PAP	PB-O1B	5.17	1.66	1.50
5	B	403	PAP	O3'-C3'	-3.54	1.32	1.44

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	403	PAP	C6-N6	3.19	1.45	1.34
5	B	403	PAP	PA-O5'	3.17	1.71	1.59
5	B	403	PAP	C3'-C4'	2.70	1.59	1.52
5	B	403	PAP	C1'-N9	-2.52	1.43	1.49
5	B	403	PAP	C2-N3	2.32	1.35	1.32
5	B	403	PAP	C2-N1	2.08	1.37	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	403	PAP	N3-C2-N1	-6.26	120.17	128.67
5	B	403	PAP	C3'-C2'-C1'	3.55	107.69	99.89
5	B	403	PAP	P-O3'-C3'	-2.21	117.53	123.43
5	B	403	PAP	C4'-O4'-C1'	-2.01	108.08	109.92

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	403	PAP	C5'-O5'-PA-O2A
5	B	403	PAP	C4'-C5'-O5'-PA
5	B	403	PAP	C3'-C4'-C5'-O5'
5	B	403	PAP	O4'-C4'-C5'-O5'
5	B	403	PAP	PA-O3A-PB-O2B
5	B	403	PAP	PA-O3A-PB-O3B
5	B	403	PAP	C5'-O5'-PA-O1A
5	B	403	PAP	C5'-O5'-PA-O3A
5	B	403	PAP	C3'-O3'-P-O2
5	B	403	PAP	PB-O3A-PA-O2A


There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	404	PO4	1	0
5	B	403	PAP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	289/295 (97%)	1.46	75 (25%) 2 2	68, 96, 132, 170	0
2	B	293/295 (99%)	0.98	40 (13%) 8 8	55, 71, 108, 127	0
All	All	582/590 (98%)	1.22	115 (19%) 3 3	55, 86, 122, 170	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	298	GLY	5.9
2	B	297	VAL	5.5
2	B	48	ASN	5.0
1	A	197	TRP	5.0
1	A	158	TRP	4.3
2	B	88	ASN	4.2
2	B	6	LEU	4.0
2	B	273	ASP	4.0
1	A	228	PHE	4.0
1	A	211	LEU	3.9
1	A	233	TRP	3.9
2	B	197	TRP	3.7
1	A	6	LEU	3.7
1	A	13	VAL	3.6
1	A	111	ALA	3.6
1	A	215	PHE	3.5
1	A	159	VAL	3.5
1	A	203	ASP	3.5
1	A	98	PHE	3.4
2	B	127	TYR	3.4
1	A	96	LYS	3.4
2	B	113	ALA	3.3
1	A	192	TYR	3.3
1	A	48	ASN	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	205	ARG	3.2
1	A	251	SER	3.2
1	A	252	TYR	3.1
1	A	112	ARG	3.1
1	A	210	VAL	3.1
1	A	47	GLU	3.1
1	A	204	THR	3.1
1	A	194	ILE	3.1
2	B	167	ASN	3.1
1	A	9	CYS	3.0
1	A	79	SER	3.0
1	A	202	THR	3.0
2	B	227	LEU	2.9
1	A	208	PHE	2.9
2	B	98	PHE	2.9
1	A	5	THR	2.9
2	B	230	TYR	2.9
1	A	167	ASN	2.9
2	B	47	GLU	2.8
2	B	263	CYS	2.8
2	B	134	ILE	2.8
1	A	144	ALA	2.8
1	A	214	PHE	2.7
2	B	78	ALA	2.7
1	A	254	PRO	2.7
2	B	77	PRO	2.7
1	A	193	PHE	2.7
2	B	44	GLY	2.7
2	B	9	CYS	2.7
1	A	212	PHE	2.7
1	A	296	LEU	2.6
1	A	250	PHE	2.6
2	B	86	LEU	2.6
1	A	10	CYS	2.6
2	B	299	MET	2.5
1	A	24	PHE	2.5
1	A	189	VAL	2.5
2	B	224	VAL	2.5
1	A	279	LEU	2.5
1	A	116	ILE	2.5
1	A	7	TRP	2.5
2	B	245	PHE	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	121	ALA	2.4
1	A	280	LEU	2.4
2	B	220	PHE	2.4
1	A	119	THR	2.4
2	B	232	CYS	2.4
1	A	196	PHE	2.4
1	A	124	THR	2.4
1	A	273	ASP	2.4
1	A	275	LYS	2.4
1	A	227	LEU	2.4
2	B	84	PHE	2.4
1	A	81	SER	2.3
1	A	234	LEU	2.3
1	A	123	LYS	2.3
1	A	84	PHE	2.3
1	A	206	ALA	2.3
2	B	13	VAL	2.3
2	B	283	PHE	2.3
1	A	100	GLN	2.3
1	A	263	CYS	2.2
2	B	194	ILE	2.2
2	B	91	LYS	2.2
1	A	90	GLU	2.2
1	A	232	CYS	2.2
2	B	290	CYS	2.2
1	A	230	TYR	2.2
1	A	8	ARG	2.2
1	A	237	LYS	2.2
1	A	285	SER	2.1
2	B	296	LEU	2.1
1	A	160	ASN	2.1
2	B	111	ALA	2.1
2	B	40	PHE	2.1
1	A	49	GLY	2.1
1	A	291	SER	2.1
1	A	50	LYS	2.1
1	A	213	LEU	2.1
2	B	7	TRP	2.1
2	B	85	TYR	2.1
1	A	220	PHE	2.1
2	B	12	ARG	2.1
2	B	196	PHE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	262	GLY	2.1
1	A	287	GLY	2.1
2	B	49	GLY	2.1
1	A	74	PHE	2.0
1	A	264	SER	2.0
1	A	231	HIS	2.0
1	A	113	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CME	B	156	10/11	0.95	0.10	56,69,83,89	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	PAP	B	403	31/31	0.74	0.19	72,112,132,165	0
4	PO4	A	404	5/5	0.81	0.08	105,111,116,124	0
4	PO4	A	403	5/5	0.84	0.10	88,96,103,104	0
4	PO4	B	404	5/5	0.87	0.10	79,81,94,99	0
3	ZN	A	402	1/1	0.99	0.03	93,93,93,93	0
3	ZN	B	402	1/1	0.99	0.04	70,70,70,70	0
3	ZN	B	401	1/1	1.00	0.01	69,69,69,69	0
3	ZN	A	401	1/1	1.00	0.02	93,93,93,93	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers ⓘ

There are no such residues in this entry.