



wwPDB EM Validation Summary Report ⓘ

Apr 24, 2025 – 12:37 PM EDT

PDB ID : 9BMO / pdb_00009bmo
EMDB ID : EMD-44705
Title : State-6 of motor domain from full-length human dynein-1 in 5mM AMPPNP
Authors : Chai, P.; Zhang, K.
Deposited on : 2024-05-02
Resolution : 3.60 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev117
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	4646	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 21867 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytoplasmic dynein 1 heavy chain 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	2711	21755	13852	3757	4035	111	0	0

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
2	A	1	27	10	5	10	2	0
2	A	1	27	10	5	10	2	0
2	A	1	27	10	5	10	2	0

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).

ALA
GLU
GLU
ILE
SER
ASP
ALA
ILE
ARG
GLU
LYS
MET
LYS
GLN
LYS
ASN
TYR
MET
SER
SER
ASN
PRO
SER
TYR
TYR
GLU
GLU
VAL
SER
ASN
ILE
ARG
ALA
SER
SER
LEU
ALA
CYS
GLY
PRO
MET
VAL
VAL
TRP
ILE
ALA
GLN
LEU
ASN
TYR
ALA
ASP
MET
LEU
LYS
ARG
ASN
GLU

LEU
GLN
LYS
LEU
GLU
ASP
ASP
ALA
LYS
ASP
GLN
GLN
LYS
ALA
ASN
ASN
GLU
VAL
GLU
GLN
MET
SER
ILE
ARG
ASP
LEU
GLU
ALA
SER
ILE
ALA
ARG
TYR
LYS
GLU
GLY
TYR
VAL
LEU
LYS
LEU
ASP
MET
ALA
VAL
GLU
ALA
LYS
V3472
N3473
R3474
S3475

T3476
A3477
L3478
L3479
R3480
S3481
L3482
S3483
A3484
E3485
R3486
E3487
R3488
L3489
E3490
K3491
T3492
S3493
E3494
T3495
F3496
K3497
N3498
Q3499
R3500
S3501
T3502
L3508
L3509
S3510
A3511
A3512
F3513
L3514
A3515
Y3516
A3517
G3518
Y3519
F3520
K3524
K3525
Q3526
F3529
L3536
Q3542
F3543
R3544
T3545
D3546
L3547
A3548
R3549
L3553

S3566
D3570
D3571
L3572
E3575
M3579
S3583
R3582
L3588
I3589
I3590
D3591
G3594
Q3595
T3597
M3601
N3602
D3606
R3607
K3608
R3611
T3612
D3617
A3618
R3619
R3620
E3624
R3628
F3629
G3630
V3635
Q3636
D3637
V3638
E3639
D3642
P3643
L3645
N3646
L3649
N3650
V3653

R3654
R3655
T3656
G3657
R3658
R3659
V3660
L3661
L3662
T3663
L3664
G3665
D3666
Q3667
D3668
L3669
D3670
L3671
F3675
L3679
S3680
D3683
P3684
T3685
V3686
D3691
L3692
C3693
S3694
V3699
N3700
T3702
V3703
T3704
R3705
S3706
C3712
E3715
V3716
L3717
K3718
V3724
D3725
E3726
K3727
R3728
S3729
D3730
L3731
L3732
K3733

L3734
Q3735
G3736
E3737
F3738
Q3739
L3740
R3741
L3742
R3743
Q3744
L3745
E3746
K3747
S3748
L3749
L3750
Q3751
A3752
L3753
N3754
E3755
V3756
K3757
G3758
R3759
L3760
L3761
D3762
D3763
D3764
T3765
L3766
L3767
T3768
T3769
L3770
E3771
N3772
L3773
K3774
R3775
E3776
A3777
A3778
K3779
R3780
T3781
R3782
K3783
V3784
E3785
E3786
D3788
L3789
V3790
N3791
Q3792
E3793

V3794
E3795
T3796
Q3799
S3805
C3808
S3817
L3818
K3819
L3824
Y3825
L3829
Y3836
V3839
L3840
D3851
Q3854
R3855
L3856
S3857
L3858
I3859
L3863
G3874
M3875
L3876
D3879
H3880
I3881
M3885
L3886
L3887
A3888
L3892
T3895
V3896
G3897
E3898
D3902
A3903
E3904
L3909

N3912
E3913
L3914
V3915
L3916
S3917
G3919
S3920
L3924
L3927
T3928
V3929
E3930
Q3931
A3932
E3933
A3934
V3935
V3936
R3937
L3938
K3945
B3946
L3947
N4063
L3948
V3951
E3955
L3959
P3966
E3967
P3971
L3973
K3974
S3975
E3976
T3981
L3982
L3983
G3984
Q3985
H3988
L3989
L3990
L3991
L3992
R3997

P3998
D3999
R4000
M4004
M4007
S4010
L4013
G4014
E4015
M4021
L4027
T4028
H4029
I4030
I4031
V4035
D4050
L4058
A4059
A4060
E4061
Q4062
N4063
T4064
Q4065
I4066
I4069
A4070
I4071
G4072
S4073
A4074
E4075
G4076
F4077
H4078
Q4079
A4080
D4081
W4083
V4084
M4085
L4096
V4099
A4102
W4105

L4106
L4109
K4112
L4113
L4124
L4130
N4131
P4132
K4133
L4138
R4143
L4144
F4145
V4146
K4154
T4160
C4170
K4171
R4176
L4177
R4178
L4179
F4186
H4187
A4188
L4189
L4190
Q4191
L4194
W4201
S4202
D4211
L4212
D4217
D4220
L4223
L4238
M4247
I4251

R4255
V4256
M4258
E4259
D4261
Q4262
R4263
S4277
E4281
F4282
K4283
V4287
W4288
D4289
G4290
H4291
K4292
D4293
L4294
Q4295
M4296
P4297
R4301
Q4307
V4308
V4309
E4310
L4311
L4312
S4319
W4320
L4321
M4325
R4329
D4338
M4339
K4342
M4343
L4344
K4345
M4346
Q4347
MET
LEU
GLU
ASP
GLU
ASP

ASP
LEU
ALA
TYR
ALA
GLU
THR
GLU
LYS
THR
THR
THR
ASP
SER
THR
ASP
GLY
ARG
P4374
A4375
W4376
M4377
R4378
T4379
L4380
H4381
T4382
T4383
L4388
R4400
L4405
L4423
L4431
C4438
E4439
K4441
K4442
K4443
Q4444
T4445
M4446
L4447
L4448
R4449
T4450
L4451
I4452
N4453
V4456
R4457

R4462
S4463
H4466
M4473
I4476
Q4477
K4487
Q4488
L4489
Q4490
N4491
I4492
K4505
F4515
V4516
P4517
E4518
I4521
T4522
Q4526
V4527
V4528
L4536
E4537
E4538
Q4549
G4550
A4551
T4552
L4553
D4554
L4575
S4578
I4581
W4592
V4593
K4594
M4597
T4598
E4599
V4604
V4609

Y4610	D4617	E4625	T4628	V4642
L4611	L4618	F4635	F4638	T4645
N4612	I4619	R4638	R4638	GLU
F4613				
T4614				

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	82309	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	45000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.305	Depositor
Minimum map value	-0.690	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.038	Depositor
Recommended contour level	0.2	Depositor
Map size (Å)	333.312, 333.312, 333.312	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.302, 1.302, 1.302	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/22220	0.52	3/30121 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4132	PRO	CA-N-CD	-12.08	94.59	111.50
1	A	4132	PRO	N-CD-CG	-6.11	94.04	103.20
1	A	2494	LEU	CA-CB-CG	5.62	128.24	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	21755	0	21789	606	0
2	A	81	0	36	8	0
3	A	31	0	12	7	0
All	All	21867	0	21837	606	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 606 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2452:LEU:HD23	1:A:2729:ARG:HD2	1.57	0.84
1:A:3570:ASP:OD1	1:A:3571:ASP:N	2.20	0.74
1:A:4388:LEU:HD21	1:A:4431:LEU:HB3	1.70	0.74
1:A:2935:LEU:HD13	1:A:2943:LYS:HB3	1.70	0.73
1:A:3736:GLY:O	1:A:3740:LEU:N	2.22	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	2699/4646 (58%)	2641 (98%)	58 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	2407/4125 (58%)	2343 (97%)	64 (3%)	40 65

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4289	ASP
1	A	4447	TYR
1	A	2664	ASP
1	A	2615	MET
1	A	4451	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2857	HIS
1	A	3602	ASN
1	A	4526	GLN
1	A	3650	ASN
1	A	2730	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	A	4702	-	28,33,33	0.67	0	34,52,52	0.60	1 (2%)
2	ADP	A	4703	-	24,29,29	0.92	1 (4%)	29,45,45	1.17	2 (6%)
2	ADP	A	4704	-	24,29,29	0.86	0	29,45,45	1.21	2 (6%)
2	ADP	A	4701	-	24,29,29	0.84	0	29,45,45	1.29	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	4702	-	-	0/18/38/38	0/3/3/3
2	ADP	A	4703	-	-	5/12/32/32	0/3/3/3
2	ADP	A	4704	-	-	5/12/32/32	0/3/3/3
2	ADP	A	4701	-	-	1/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	4703	ADP	PA-O3A	2.12	1.61	1.59

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	4701	ADP	N3-C2-N1	-3.78	123.54	128.67
2	A	4703	ADP	N3-C2-N1	-3.74	123.60	128.67
2	A	4704	ADP	N3-C2-N1	-3.69	123.66	128.67
2	A	4703	ADP	C4-C5-N7	-2.62	106.57	109.34
2	A	4704	ADP	C4-C5-N7	-2.55	106.64	109.34

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	4703	ADP	C5'-O5'-PA-O2A
2	A	4704	ADP	C5'-O5'-PA-O3A
2	A	4704	ADP	O4'-C4'-C5'-O5'
2	A	4704	ADP	C3'-C4'-C5'-O5'
2	A	4703	ADP	O4'-C4'-C5'-O5'



There are no ring outliers.


3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	4702	ATP	7	0
2	A	4703	ADP	2	0
2	A	4701	ADP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-44705. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 146




Y Index: 120




Z Index: 135


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map




6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_44705_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 154 nm³; this corresponds to an approximate mass of 139 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	3.63	4.07	3.67
Unmasked-calculated*	6.10	8.33	6.51

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.10 differs from the reported value 3.6 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-44705 and PDB model 9BMO. Per-residue inclusion information can be found in section [3](#) on page [5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).

9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7280	 0.3970
A	 0.7280	 0.3970

