



wwPDB EM Validation Summary Report ⓘ

May 19, 2026 – 12:16 PM EDT

PDB ID : 9DVU / pdb_00009dvu
EMDB ID : EMD-47223
Title : CryoEM structure of RpaA bound to PkaiBC DNA and Syn7942 RNAP-SigA holoenzyme
Authors : Fang, M.; Gu, Y.; Matyszewski, M.; Leanca, M.; LiWang, A.; Yuzenkova, Y.; Corbett, K.D.; Golden, S.E.
Deposited on : 2024-10-08
Resolution : 3.70 Å (reported)
Based on initial model : 8SYI

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11569 (3.20 - 4.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	105	
2	2	105	
3	A	309	
3	B	309	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	C	1100	<div><div></div><div>67%26%5%</div></div>
5	D	624	<div><div></div><div>73%25%</div></div>
6	E	76	<div><div></div><div>42%61%11%28%</div></div>
7	F	1318	<div><div></div><div>12%65%25%8%</div></div>
8	G	399	<div><div></div><div>58%18%23%</div></div>
9	R	249	<div><div></div><div>73%20%5%</div></div>
9	S	249	<div><div></div><div>67%22%7%</div></div>

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 35249 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	1	61	Total	C	N	O	P	0	0
			1253	600	219	373	61		

- Molecule 2 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	2	52	Total	C	N	O	P	0	0
			1067	509	199	307	52		

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	A	278	Total	C	N	O	S	0	0
			2147	1346	375	421	5		
3	B	215	Total	C	N	O	S	0	0
			1644	1032	290	318	4		

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	C	1042	Total	C	N	O	S	0	0
			8222	5166	1454	1575	27		

- Molecule 5 is a protein called DNA-directed RNA polymerase subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	D	617	Total	C	N	O	S	0	0
			4926	3107	892	906	21		

- Molecule 6 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	E	55	Total	C	N	O	S	0	0
			451	281	81	87	2		

- Molecule 7 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	F	1209	Total	C	N	O	S	0	0
			9265	5771	1643	1830	21		

- Molecule 8 is a protein called RNA polymerase sigma factor SigA1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	G	308	Total	C	N	O	S	0	0
			2491	1564	456	466	5		

- Molecule 9 is a protein called DNA-binding dual master transcriptional regulator RpaA.

Mol	Chain	Residues	Atoms						AltConf	Trace
9	R	236	Total	C	N	O	P	S	0	0
			1904	1206	330	358	1	9		
9	S	232	Total	C	N	O	P	S	0	0
			1876	1191	325	350	1	9		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	121	GLN	ARG	conflict	UNP Q31S42
S	121	GLN	ARG	conflict	UNP Q31S42

- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
10	D	1	Total	Mg	0
			1	1	

- Molecule 11 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
11	D	1	Total	Zn	0
			1	1	

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
11	F	1	1	1	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-template DNA



- Molecule 2: Template DNA





Chain B: 




• Molecule 4: DNA-directed RNA polymerase subunit beta

Chain C: 







• Molecule 5: DNA-directed RNA polymerase subunit gamma




• Molecule 6: DNA-directed RNA polymerase subunit omega




• Molecule 7: DNA-directed RNA polymerase subunit beta'






• Molecule 9: DNA-binding dual master transcriptional regulator RpaA

Chain R: 73% 20% 5%



• Molecule 9: DNA-binding dual master transcriptional regulator RpaA

Chain S: 67% 22% 7%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	221552	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.017	Depositor
Minimum map value	-0.111	Depositor
Average map value	0.006	Depositor
Map value standard deviation	0.019	Depositor
Recommended contour level	0.15	Depositor
Map size (\AA)	478.72, 478.72, 478.72	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.1968, 1.1968, 1.1968	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, PHD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1	0.36	0/1403	0.53	0/2165
2	2	0.35	0/1198	0.50	0/1846
3	A	0.26	0/2178	0.41	0/2961
3	B	0.26	0/1670	0.38	0/2274
4	C	0.31	0/8376	0.45	0/11340
5	D	0.28	0/5019	0.38	0/6793
6	E	0.17	0/454	0.45	0/608
7	F	0.23	0/9383	0.40	0/12707
8	G	0.23	0/2521	0.44	0/3389
9	R	0.27	0/1927	0.57	0/2612
9	S	0.27	0/1899	0.53	0/2574
All	All	0.27	0/36028	0.44	0/49269

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	B	0	1
4	C	0	5
6	E	0	1
7	F	0	1
8	G	0	2
9	R	0	3
9	S	0	2
All	All	0	15

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	B	183	ARG	Sidechain
4	C	247	ARG	Sidechain
4	C	253	ARG	Sidechain
4	C	339	ARG	Sidechain
4	C	624	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	1253	0	694	50	0
2	2	1067	0	586	35	0
3	A	2147	0	2134	33	0
3	B	1644	0	1632	35	0
4	C	8222	0	8196	192	0
5	D	4926	0	4951	112	0
6	E	451	0	458	6	0
7	F	9265	0	9364	238	0
8	G	2491	0	2549	67	0
9	R	1904	0	1913	50	0
9	S	1876	0	1891	70	0
10	D	1	0	0	0	0
11	D	1	0	0	0	0
11	F	1	0	0	0	0
All	All	35249	0	34368	776	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 776 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:304:MET:HE2	8:G:190:GLN:HB3	1.36	1.07
3:B:147:TYR:H	3:B:170:MET:HE3	1.33	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:109:MET:HE3	4:C:113:GLY:HA2	1.51	0.91
1:1:29:DT:H3'	9:S:204:ARG:HH12	1.36	0.89
4:C:259:VAL:HG21	4:C:313:ARG:HD2	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	270/309 (87%)	265 (98%)	5 (2%)	0	100	100
3	B	209/309 (68%)	202 (97%)	7 (3%)	0	100	100
4	C	1036/1100 (94%)	988 (95%)	48 (5%)	0	100	100
5	D	613/624 (98%)	602 (98%)	10 (2%)	1 (0%)	43	72
6	E	51/76 (67%)	50 (98%)	1 (2%)	0	100	100
7	F	1199/1318 (91%)	1143 (95%)	56 (5%)	0	100	100
8	G	304/399 (76%)	294 (97%)	10 (3%)	0	100	100
9	R	231/249 (93%)	224 (97%)	5 (2%)	2 (1%)	14	45
9	S	227/249 (91%)	218 (96%)	5 (2%)	4 (2%)	6	34
All	All	4140/4633 (89%)	3986 (96%)	147 (4%)	7 (0%)	44	72

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
9	R	85	LEU
9	S	85	LEU
5	D	79	ARG
9	S	139	GLY
9	S	140	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	234/262 (89%)	221 (94%)	13 (6%)	19	46
3	B	178/262 (68%)	173 (97%)	5 (3%)	38	58
4	C	887/938 (95%)	847 (96%)	40 (4%)	24	49
5	D	532/539 (99%)	522 (98%)	10 (2%)	50	65
6	E	50/68 (74%)	49 (98%)	1 (2%)	48	64
7	F	1001/1093 (92%)	941 (94%)	60 (6%)	17	44
8	G	263/344 (76%)	256 (97%)	7 (3%)	39	59
9	R	207/217 (95%)	201 (97%)	6 (3%)	37	57
9	S	204/217 (94%)	199 (98%)	5 (2%)	42	61
All	All	3556/3940 (90%)	3409 (96%)	147 (4%)	28	52

5 of 147 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	F	756	VAL
9	S	54	LEU
7	F	857	VAL
8	G	183	MET
4	C	929	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
5	D	546	GLN
7	F	389	ASN
9	S	174	HIS
6	E	28	HIS
7	F	277	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	PHD	R	53	9	9,11,12	4.09	1 (11%)	9,15,17	1.85	3 (33%)
9	PHD	S	53	9	9,11,12	4.03	1 (11%)	9,15,17	1.76	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PHD	R	53	9	-	4/8/11/13	-
9	PHD	S	53	9	-	4/8/11/13	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	R	53	PHD	P-OD1	12.10	1.81	1.59
9	S	53	PHD	P-OD1	11.94	1.81	1.59

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	S	53	PHD	OD1-P-OP1	-3.71	97.61	109.47
9	R	53	PHD	OD1-P-OP1	-3.68	97.70	109.47
9	S	53	PHD	OP3-P-OP2	2.34	116.59	107.80
9	R	53	PHD	OP3-P-OP2	2.22	116.14	107.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	R	53	PHD	CA-CB-CG	2.02	117.11	112.78

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	S	53	PHD	CG-OD1-P-OP1
9	R	53	PHD	C-CA-CB-CG
9	R	53	PHD	CA-CB-CG-OD1
9	S	53	PHD	CA-CB-CG-OD1
9	S	53	PHD	CA-CB-CG-OD2

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	R	53	PHD	4	0
9	S	53	PHD	5	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-47223. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 199



Y Index: 201




Z Index: 185


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.15. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 335 nm^3 ; this corresponds to an approximate mass of 302 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.270 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-47223 and PDB model 9DVU. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.15 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.15).

9.4 Atom inclusion ⓘ



At the recommended contour level, 94% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.15) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.8310	<div></div> 0.4350
1	<div></div> 0.8720	<div></div> 0.3630
2	<div></div> 0.8140	<div></div> 0.3200
A	<div></div> 0.8780	<div></div> 0.4770
B	<div></div> 0.9210	<div></div> 0.5210
C	<div></div> 0.8900	<div></div> 0.5000
D	<div></div> 0.9170	<div></div> 0.5200
E	<div></div> 0.3550	<div></div> 0.4210
F	<div></div> 0.7380	<div></div> 0.3790
G	<div></div> 0.8290	<div></div> 0.4090
R	<div></div> 0.8100	<div></div> 0.3450
S	<div></div> 0.8020	<div></div> 0.3160

1.0

0.0

<0.0