



wwPDB EM Validation Summary Report ⓘ

Aug 9, 2026 – 05:57 PM JST

PDB ID : 9LRW / pdb_00009lrw
EMDB ID : EMD-63345
Title : Cryo-EM structure of Fission yeast centromeric nucleosome Class 2
Authors : Xiong, Y.; Zang, J.
Deposited on : 2025-02-01
Resolution : 3.04 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13952 (2.54 - 3.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	
1	E	120	
2	B	103	
2	F	103	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	132	 71% 26%
3	G	132	 55% 14% 31%
4	D	126	 63% 10% 26%
4	H	126	 57% 14% 29%
5	I	147	 5% 56% 39% 5%
6	J	147	 5% 48% 47% 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11213 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3-like centromeric protein cnp1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	81	Total	C	N	O	S	0	0
			618	390	121	103	4		
1	E	97	Total	C	N	O	S	0	0
			761	476	148	133	4		

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	B	78	Total	C	N	O	0	0
			623	395	120	108		
2	F	78	Total	C	N	O	0	0
			626	396	120	110		

- Molecule 3 is a protein called Histone H2A-alpha.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	98	Total	C	N	O	0	0
			742	466	146	130		
3	G	91	Total	C	N	O	0	0
			685	432	135	118		

- Molecule 4 is a protein called Histone H2B-alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	93	Total	C	N	O	S	0	0
			723	455	127	140	1		
4	H	90	Total	C	N	O	S	0	0
			698	441	121	135	1		

- Molecule 5 is a DNA chain called DNA (140-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	140	Total	C	N	O	P	0	0
			2848	1354	518	837	139		


- Molecule 6 is a DNA chain called DNA (140-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	140	Total	C	N	O	P	0	0
			2889	1367	541	841	140		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3-like centromeric protein cnp1




- Molecule 1: Histone H3-like centromeric protein cnp1



- Molecule 2: Histone H4



- Molecule 2: Histone H4



- Molecule 3: Histone H2A-alpha

Chain C:  71% 26%

MET SER GLY LYS SER GLY LYS LYS ALA VAL ALA LYS S15 V32 L65 R83 P111 H112 TLE ASN ALA HIS LEU LEU PRO LYS THR SER GLY ARG THR GLY LYS PRO SER GLN GLU LEU

- Molecule 3: Histone H2A-alpha

Chain G:  55% 14% 31%

MET SER GLY LYS SER GLY LYS LYS ALA VAL ALA LYS S18 Q17 V32 Y41 R44 V45 G46 A47 P50 Y59 L60 A61 A62 L65 E66 L67 Q86 R90 E94 L95 I104 G107 GLY VAL PRO ASN TLE ASN ALA HIS LEU

PRO LYS THR SER ARG THR GLY LYS PRO SER GLN LEU

- Molecule 4: Histone H2B-alpha

Chain D:  63% 10% 26%

MET SER ALA GLU LYS PRO ALA LYS LYS ALA GLY LYS LYS PRO ARG ASP THR MET LYS SER ALA ASP LYS LYS ARG GLY K31 S39 Y40 I41 Y42 K43 V44 L45 K46 Q47 D68 E69 F70 E71 R72 R92 R99 L100 G114 V118 S123 ALA

GLN

- Molecule 4: Histone H2B-alpha

Chain H:  57% 14% 29%

MET SER ALA GLU LYS LYS PRO ALA LYS LYS LYS PRO ARG ASP THR MET LYS SER ALA ASP LYS LYS ARG GLY ASN ARG K34 E35 T36 Y37 S38 L45 Q57 A58 M59 R60 I61 L62 N63 I69 F70 S78 T88 L106 A107 K108 H109

A110 V111 T112 E113 G114 S123 SER ALA GLN

- Molecule 5: DNA (140-MER)


Chain I:  5% 56% 39% 5%

DA T-72 C-71 G-70 A-69 G-68 A-67 C-62 G-61 G-60 T-59 G-58 C-57 G-56 G-55 A-54 C-46 A-45 G-41 G-40 T-36 A-35 C-29 T-28 C-27 T-26 C-21 C-20 G-19 C-18 T-17 G-8 G-7 C-2 G-1 C0 T1 G2 C5 C6 C7 C8 G9 T15 A16 A17 A23 A24 G25

G26 G27 T31 A32 C33 T34 C35 C36 C37 T38 A39 C45 A46 G47 G51 G54 A57 T62 A63 T64 A65 C66 A67 DT DC DC DC DA DA DT

- Molecule 6: DNA (140-MER)

Chain J:  5% 48% 47% 5%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	832775	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	55	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.664	Depositor
Minimum map value	-0.283	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.019	Depositor
Recommended contour level	0.1	Depositor
Map size ($\text{\AA}$)	273.92, 273.92, 273.92	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.07, 1.07, 1.07	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/626	0.26	0/845
1	E	0.15	0/770	0.25	0/1037
2	B	0.17	0/630	0.29	0/846
2	F	0.16	0/633	0.25	0/850
3	C	0.14	0/752	0.25	0/1016
3	G	0.14	0/693	0.31	0/935
4	D	0.15	0/733	0.27	0/989
4	H	0.17	0/708	0.30	0/956
5	I	0.18	0/3191	0.36	0/4918
6	J	0.17	0/3244	0.34	0/5010
All	All	0.17	0/11980	0.32	0/17402

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	618	0	613	12	0
1	E	761	0	765	14	0
2	B	623	0	664	19	0
2	F	626	0	666	11	0
3	C	742	0	778	4	0
3	G	685	0	719	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	723	0	734	10	0
4	H	698	0	713	14	0
5	I	2848	0	1572	41	0
6	J	2889	0	1572	57	0
All	All	11213	0	8796	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:70:ARG:HB2	2:F:81:THR:HG22	1.69	0.74
2:F:30:ILE:O	2:F:56:ARG:NH1	2.21	0.73
2:F:32:LYS:HE2	2:F:36:ARG:HH21	1.55	0.70
4:H:38:SER:HA	4:H:59:MET:HE3	1.74	0.70
5:I:68:DG:H1	6:J:68:DC:H42	1.38	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/120 (66%)	77 (98%)	2 (2%)	0	100	100
1	E	95/120 (79%)	94 (99%)	1 (1%)	0	100	100
2	B	76/103 (74%)	75 (99%)	1 (1%)	0	100	100
2	F	76/103 (74%)	75 (99%)	1 (1%)	0	100	100
3	C	96/132 (73%)	94 (98%)	2 (2%)	0	100	100
3	G	89/132 (67%)	88 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	D	91/126 (72%)	89 (98%)	2 (2%)	0	100	100
4	H	88/126 (70%)	85 (97%)	3 (3%)	0	100	100
All	All	690/962 (72%)	677 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/104 (57%)	59 (100%)	0	100	100
1	E	76/104 (73%)	76 (100%)	0	100	100
2	B	65/81 (80%)	65 (100%)	0	100	100
2	F	66/81 (82%)	66 (100%)	0	100	100
3	C	73/97 (75%)	73 (100%)	0	100	100
3	G	65/97 (67%)	65 (100%)	0	100	100
4	D	79/105 (75%)	79 (100%)	0	100	100
4	H	77/105 (73%)	77 (100%)	0	100	100
All	All	560/774 (72%)	560 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
4	D	57	GLN
1	E	44	GLN
1	E	80	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-63345. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 114




Y Index: 117




Z Index: 140


6.3.2 Raw map



X Index: 114



Y Index: 117




Z Index: 140


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 128 nm^3 ; this corresponds to an approximate mass of 115 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.329 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.329 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.04	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.42	3.87	3.48

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.42 differs from the reported value 3.04 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-63345 and PDB model 9LRW. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).

9.4 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 91% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9130	0.4360
A	0.9550	0.5190
B	0.9500	0.5050
C	0.9250	0.5090
D	0.9320	0.5030
E	0.9330	0.5160
F	0.9520	0.5230
G	0.8510	0.4600
H	0.8720	0.4590
I	0.9070	0.3740
J	0.9070	0.3770

1.0

0.0

<0.0