



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 03:12 PM EDT

PDB ID : 4ZC0
Title : Structure of a dodecameric bacterial helicase
Authors : Bazin, A.; Cherrier, M.V.; Gutsche, I.; Timmins, J.; Terradot, L.
Deposited on : 2015-04-15
Resolution : 6.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 6.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1002 (9.50-3.90)
Clashscore	141614	1066 (9.50-3.90)
Ramachandran outliers	138981	1000 (9.50-3.90)
Sidechain outliers	138945	1000 (9.50-3.86)
RSRZ outliers	127900	1004 (9.50-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	 3% 60% 25% 15%
1	B	521	 9% 59% 23% 17%
1	C	521	 2% 20% 8% 72%
1	D	521	 19% 8% 72%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	TBR	B	501	-	-	X	-

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9140 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replicative DNA helicase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	0	0
			3491	2199	611	667	14			
1	B	434	Total	C	N	O	S	0	0	0
			3397	2141	595	647	14			
1	C	148	Total	C	N	O	S	0	0	0
			1122	714	193	212	3			
1	D	146	Total	C	N	O	S	0	0	0
			1112	708	191	210	3			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-32	MET	-	initiating methionine	UNP O25916
A	-31	HIS	-	expression tag	UNP O25916
A	-30	HIS	-	expression tag	UNP O25916
A	-29	HIS	-	expression tag	UNP O25916
A	-28	HIS	-	expression tag	UNP O25916
A	-27	HIS	-	expression tag	UNP O25916
A	-26	HIS	-	expression tag	UNP O25916
A	-25	GLY	-	expression tag	UNP O25916
A	-24	LYS	-	expression tag	UNP O25916
A	-23	PRO	-	expression tag	UNP O25916
A	-22	ILE	-	expression tag	UNP O25916
A	-21	PRO	-	expression tag	UNP O25916
A	-20	ASN	-	expression tag	UNP O25916
A	-19	PRO	-	expression tag	UNP O25916
A	-18	LEU	-	expression tag	UNP O25916
A	-17	LEU	-	expression tag	UNP O25916
A	-16	GLY	-	expression tag	UNP O25916
A	-15	LEU	-	expression tag	UNP O25916
A	-14	ASP	-	expression tag	UNP O25916
A	-13	SER	-	expression tag	UNP O25916
A	-12	THR	-	expression tag	UNP O25916

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	GLU	-	expression tag	UNP O25916
A	-10	ASN	-	expression tag	UNP O25916
A	-9	LEU	-	expression tag	UNP O25916
A	-8	TYR	-	expression tag	UNP O25916
A	-7	PHE	-	expression tag	UNP O25916
A	-6	GLN	-	expression tag	UNP O25916
A	-5	GLY	-	expression tag	UNP O25916
A	-4	ILE	-	expression tag	UNP O25916
A	-3	ASP	-	expression tag	UNP O25916
A	-2	PRO	-	expression tag	UNP O25916
A	-1	PHE	-	expression tag	UNP O25916
A	0	THR	-	expression tag	UNP O25916
B	-32	MET	-	initiating methionine	UNP O25916
B	-31	HIS	-	expression tag	UNP O25916
B	-30	HIS	-	expression tag	UNP O25916
B	-29	HIS	-	expression tag	UNP O25916
B	-28	HIS	-	expression tag	UNP O25916
B	-27	HIS	-	expression tag	UNP O25916
B	-26	HIS	-	expression tag	UNP O25916
B	-25	GLY	-	expression tag	UNP O25916
B	-24	LYS	-	expression tag	UNP O25916
B	-23	PRO	-	expression tag	UNP O25916
B	-22	ILE	-	expression tag	UNP O25916
B	-21	PRO	-	expression tag	UNP O25916
B	-20	ASN	-	expression tag	UNP O25916
B	-19	PRO	-	expression tag	UNP O25916
B	-18	LEU	-	expression tag	UNP O25916
B	-17	LEU	-	expression tag	UNP O25916
B	-16	GLY	-	expression tag	UNP O25916
B	-15	LEU	-	expression tag	UNP O25916
B	-14	ASP	-	expression tag	UNP O25916
B	-13	SER	-	expression tag	UNP O25916
B	-12	THR	-	expression tag	UNP O25916
B	-11	GLU	-	expression tag	UNP O25916
B	-10	ASN	-	expression tag	UNP O25916
B	-9	LEU	-	expression tag	UNP O25916
B	-8	TYR	-	expression tag	UNP O25916
B	-7	PHE	-	expression tag	UNP O25916
B	-6	GLN	-	expression tag	UNP O25916
B	-5	GLY	-	expression tag	UNP O25916
B	-4	ILE	-	expression tag	UNP O25916
B	-3	ASP	-	expression tag	UNP O25916

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	PRO	-	expression tag	UNP O25916
B	-1	PHE	-	expression tag	UNP O25916
B	0	THR	-	expression tag	UNP O25916
C	-32	MET	-	initiating methionine	UNP O25916
C	-31	HIS	-	expression tag	UNP O25916
C	-30	HIS	-	expression tag	UNP O25916
C	-29	HIS	-	expression tag	UNP O25916
C	-28	HIS	-	expression tag	UNP O25916
C	-27	HIS	-	expression tag	UNP O25916
C	-26	HIS	-	expression tag	UNP O25916
C	-25	GLY	-	expression tag	UNP O25916
C	-24	LYS	-	expression tag	UNP O25916
C	-23	PRO	-	expression tag	UNP O25916
C	-22	ILE	-	expression tag	UNP O25916
C	-21	PRO	-	expression tag	UNP O25916
C	-20	ASN	-	expression tag	UNP O25916
C	-19	PRO	-	expression tag	UNP O25916
C	-18	LEU	-	expression tag	UNP O25916
C	-17	LEU	-	expression tag	UNP O25916
C	-16	GLY	-	expression tag	UNP O25916
C	-15	LEU	-	expression tag	UNP O25916
C	-14	ASP	-	expression tag	UNP O25916
C	-13	SER	-	expression tag	UNP O25916
C	-12	THR	-	expression tag	UNP O25916
C	-11	GLU	-	expression tag	UNP O25916
C	-10	ASN	-	expression tag	UNP O25916
C	-9	LEU	-	expression tag	UNP O25916
C	-8	TYR	-	expression tag	UNP O25916
C	-7	PHE	-	expression tag	UNP O25916
C	-6	GLN	-	expression tag	UNP O25916
C	-5	GLY	-	expression tag	UNP O25916
C	-4	ILE	-	expression tag	UNP O25916
C	-3	ASP	-	expression tag	UNP O25916
C	-2	PRO	-	expression tag	UNP O25916
C	-1	PHE	-	expression tag	UNP O25916
C	0	THR	-	expression tag	UNP O25916
D	-32	MET	-	initiating methionine	UNP O25916
D	-31	HIS	-	expression tag	UNP O25916
D	-30	HIS	-	expression tag	UNP O25916
D	-29	HIS	-	expression tag	UNP O25916
D	-28	HIS	-	expression tag	UNP O25916
D	-27	HIS	-	expression tag	UNP O25916

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-26	HIS	-	expression tag	UNP O25916
D	-25	GLY	-	expression tag	UNP O25916
D	-24	LYS	-	expression tag	UNP O25916
D	-23	PRO	-	expression tag	UNP O25916
D	-22	ILE	-	expression tag	UNP O25916
D	-21	PRO	-	expression tag	UNP O25916
D	-20	ASN	-	expression tag	UNP O25916
D	-19	PRO	-	expression tag	UNP O25916
D	-18	LEU	-	expression tag	UNP O25916
D	-17	LEU	-	expression tag	UNP O25916
D	-16	GLY	-	expression tag	UNP O25916
D	-15	LEU	-	expression tag	UNP O25916
D	-14	ASP	-	expression tag	UNP O25916
D	-13	SER	-	expression tag	UNP O25916
D	-12	THR	-	expression tag	UNP O25916
D	-11	GLU	-	expression tag	UNP O25916
D	-10	ASN	-	expression tag	UNP O25916
D	-9	LEU	-	expression tag	UNP O25916
D	-8	TYR	-	expression tag	UNP O25916
D	-7	PHE	-	expression tag	UNP O25916
D	-6	GLN	-	expression tag	UNP O25916
D	-5	GLY	-	expression tag	UNP O25916
D	-4	ILE	-	expression tag	UNP O25916
D	-3	ASP	-	expression tag	UNP O25916
D	-2	PRO	-	expression tag	UNP O25916
D	-1	PHE	-	expression tag	UNP O25916
D	0	THR	-	expression tag	UNP O25916


- Molecule 2 is HEXATANTALUM DODECABROMIDE (three-letter code: TBR) (formula: $\text{Br}_{12}\text{Ta}_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	Br	Ta	0	0
			18	12	6		



● Molecule 1: Replicative DNA helicase



● Molecule 1: Replicative DNA helicase



ASN	ALA	LEU	LYS	LEU
ALA	GLU	SER	LEU	LYS
PRO	ASP	THR	LEU	LYS
PRO	ASN	LEU	PHE	LEU
THR	LYS	ALA	ARG	ASP
ARG	ILE	ARG	GLU	ASP
TYR	ASP	GLU	LEU	LYS
GLU	LEU	LEU	ILE	GLU
ASP	LEU	GLU	PRO	VAL
PRO	LYS	PRO	GLU	ILE
ILE	GLU	ILE	GLU	ILE
ASP	GLY	ILE	ILE	ILE
HIS	LYS	ALA	ALA	PHE
SER	ILE	LEU	GLN	SER
HIS	GLU	VAL	ILE	GLU
GLU	GLU	LEU	GLN	ARG
LEU	ALA	LEU	LEU	LEU
GLU	GLN	ASN	ASN	GLN
GLY	GLN	ASN	LEU	GLN
GLN	GLN	ASN	LEU	GLN
GLY	GLN	ASN	LEU	GLN
VAL	VAL	ASN	LEU	LYS
LYS	LYS	GLU	GLU	LEU
VAL	VAL	ASN	ASN	LEU
ASP	VAL	ARG	ARG	SER
TYR	ASN	ASP	ASP	GLN
ASP	GLU	ASP	ASP	GLN
ILE	GLU	ASP	HIS	LYS
ILE	ILE	LYS	LYS	LYS
VAL	ARG	ARG	GLU	ASP
THR	ARG	PRO	LEU	LEU
THR	ILE	ILE	GLY	THR
THR	HIS	LEU	ILE	THR
THR	LYS	SER	ILE	SER
THR	GLN	ASP	ASP	ASP
THR	GLN	ASP	ASP	ASP
THR	ASN	VAL	THR	GLN
THR	ARG	LYS	LYS	TRP
THR	ASN	LEU	GLU	ASN
THR	GLY	PHE	ARG	ASN
THR	ALA	LEU	HIS	LYS
THR	ALA	TYR	GLU	ALA
THR	THR	ARG	GLN	LYS
THR	GLY	GLY	ILE	CYS
THR	THR	TYR	ALA	PHE
THR	VAL	ILE	GLU	ASP
THR	VAL	TYR	ILE	HIS
THR	THR	TYR	SER	LYS
THR	THR	GLN	LEU	ARG
THR	ARG	MET	GLU	GLN
THR	PHE	ARG	THR	THR

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	283.47Å 283.47Å 283.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.24 – 6.70 47.24 – 6.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.24-6.70) 99.9 (47.24-6.70)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.43 (at 6.67Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.258 , 0.299 0.266 , 0.305	Depositor DCC
R_{free} test set	355 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	579.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 363.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.037 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	9140	wwPDB-VP
Average B, all atoms (Å ²)	315.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TBR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.25	0/3536	0.46	0/4757
1	B	0.25	0/3439	0.45	0/4624
1	C	0.28	0/1140	0.50	1/1542 (0.1%)
1	D	0.28	0/1130	0.47	0/1528
All	All	0.26	0/9245	0.46	1/12451 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	-2	PRO	N-CA-CB	5.69	110.13	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3491	0	3456	93	2
1	B	3397	0	3357	80	2
1	C	1122	0	1062	29	0
1	D	1112	0	1062	35	0
2	B	18	0	0	6	0
All	All	9140	0	8937	226	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:LEU:HG	2:B:501:TBR:BR9	2.15	1.01
1:B:301:LYS:NZ	1:B:305:LYS:O	2.07	0.86
1:A:301:LYS:NZ	1:A:305:LYS:O	2.08	0.84
1:C:99:GLU:O	1:C:103:ASN:ND2	2.13	0.81
1:B:39:ASP:O	1:B:108:ARG:NH2	2.14	0.80

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:ALA:O	1:B:107:LYS:NZ[9_555]	1.94	0.26
1:A:107:LYS:NZ	1:B:124:SER:O[9_555]	2.02	0.18

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	433/521 (83%)	423 (98%)	10 (2%)	0	100	100
1	B	422/521 (81%)	409 (97%)	13 (3%)	0	100	100
1	C	144/521 (28%)	138 (96%)	5 (4%)	1 (1%)	22	63
1	D	142/521 (27%)	136 (96%)	5 (4%)	1 (1%)	22	63
All	All	1141/2084 (55%)	1106 (97%)	33 (3%)	2 (0%)	47	81

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	124	SER
1	C	73	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/461 (80%)	365 (99%)	4 (1%)	73	84
1	B	356/461 (77%)	348 (98%)	8 (2%)	52	71
1	C	110/461 (24%)	109 (99%)	1 (1%)	78	87
1	D	110/461 (24%)	109 (99%)	1 (1%)	78	87
All	All	945/1844 (51%)	931 (98%)	14 (2%)	65	80

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	119	GLU
1	B	333	GLU
1	D	111	PHE
1	B	458	ASN
1	C	119	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	472	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TBR	B	501	-	0,36,36	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	TBR	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/521 (85%)	0.18	17 (3%) 40 36	58, 291, 466, 550	0
1	B	434/521 (83%)	0.55	45 (10%) 6 9	75, 341, 550, 550	0
1	C	148/521 (28%)	0.29	8 (5%) 25 26	118, 270, 425, 539	0
1	D	146/521 (28%)	-0.03	2 (1%) 75 66	114, 260, 386, 539	0
All	All	1173/2084 (56%)	0.30	72 (6%) 21 20	58, 301, 502, 550	0

The worst 5 of 72 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	388	LEU	6.7
1	B	467	MET	6.4
1	B	203	ALA	5.6
1	B	466	ASP	4.7
1	B	318	MET	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TBR	B	501	18/18	0.90	0.18	550,550,550,550	18

6.5 Other polymers [i](#)

There are no such residues in this entry.