



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 27, 2026 – 06:10 PM JST

PDB ID : 26AO / pdb_000026ao
Title : E. coli peptidyl-prolyl cis-trans isomerase, mutant F4/F5Phe
Authors : Frkic, R.L.; Jackson, C.J.
Deposited on : 2026-04-24
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1380 (1.16-1.12)
Clashscore	190562	1393 (1.16-1.12)
Ramachandran outliers	187476	1369 (1.16-1.12)
Sidechain outliers	187428	1369 (1.16-1.12)
RSRZ outliers	180081	1379 (1.16-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	
1	B	170	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5492 atoms, of which 2567 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidyl-prolyl cis-trans isomerase B.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	170	Total	C	F	H	N	O	S	0	4	0
			2650	850	5	1291	241	254	9			
1	B	170	Total	C	F	H	N	O	S	0	2	0
			2622	842	5	1276	238	253	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	HIS	-	expression tag	UNP P23869
A	166	HIS	-	expression tag	UNP P23869
A	167	HIS	-	expression tag	UNP P23869
A	168	HIS	-	expression tag	UNP P23869
A	169	HIS	-	expression tag	UNP P23869
A	170	HIS	-	expression tag	UNP P23869
B	165	HIS	-	expression tag	UNP P23869
B	166	HIS	-	expression tag	UNP P23869
B	167	HIS	-	expression tag	UNP P23869
B	168	HIS	-	expression tag	UNP P23869
B	169	HIS	-	expression tag	UNP P23869
B	170	HIS	-	expression tag	UNP P23869


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	154	Total	O	0	0
			154	154		
2	B	66	Total	O	0	0
			66	66		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptidyl-prolyl cis-trans isomerase B



- Molecule 1: Peptidyl-prolyl cis-trans isomerase B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	34.91Å 39.20Å 61.62Å 78.04° 79.29° 67.01°	Depositor
Resolution (Å)	31.92 – 1.15 31.92 – 1.15	Depositor EDS
% Data completeness (in resolution range)	94.5 (31.92-1.15) 94.5 (31.92-1.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.15Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.197 , 0.226 0.197 , 0.227	Depositor DCC
R_{free} test set	4891 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	12.8	Xtriage
Anisotropy	0.245	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5492	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PF5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/1392	0.68	0/1877
1	B	0.35	0/1367	0.59	0/1845
All	All	0.42	0/2759	0.64	0/3722

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1359	1291	1273	2	0
1	B	1346	1276	1267	12	0
2	A	154	0	0	0	0
2	B	66	0	0	1	0
All	All	2925	2567	2540	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:132:VAL:HA	1:B:135:LYS:HE2	1.75	0.68
1:B:146:MET:HE2	1:B:146:MET:HA	1.83	0.61
1:B:21:PRO:HD2	1:B:71:GLU:OE2	2.05	0.57
1:B:131:ASP:O	1:B:135:LYS:HG3	2.09	0.53
1:B:132:VAL:HG22	1:B:135:LYS:HE2	1.91	0.52
1:B:114:SER:OG	1:B:116:GLN:NE2	2.46	0.48
1:B:77:LYS:HD2	1:B:109:ASN:ND2	2.28	0.48
1:B:132:VAL:HA	1:B:135:LYS:HG3	1.96	0.46
1:A:1[B]:MET:CE	1:A:169:HIS:CD2	2.98	0.46
1:A:33:GLU:O	1:A:33:GLU:HG3	2.18	0.44
1:B:59:MET:HE2	1:B:147:HIS:CG	2.55	0.42
1:B:72:ALA:HB1	1:B:110:PHE:HB2	2.02	0.41
1:B:168:HIS:CD2	1:B:168:HIS:C	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/170 (100%)	163 (96%)	7 (4%)	0	100	100
1	B	169/170 (99%)	160 (95%)	9 (5%)	0	100	100
All	All	339/340 (100%)	323 (95%)	16 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/143 (103%)	147 (100%)	0	100	100
1	B	145/143 (101%)	145 (100%)	0	100	100
All	All	292/286 (102%)	292 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	101	ASN
1	A	105	ASN
1	A	169	HIS
1	B	8	HIS
1	B	101	ASN
1	B	105	ASN
1	B	116	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PF5	B	4	1	15,16,17	1.34	1 (6%)	19,23,25	1.43	3 (15%)
1	PF5	A	4	1	15,16,17	1.08	1 (6%)	19,23,25	1.27	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PF5	B	4	1	-	2/5/6/8	0/1/1/1
1	PF5	A	4	1	-	2/5/6/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	4	PF5	CE1-CD1	3.06	1.43	1.37
1	A	4	PF5	CB-CG	-2.19	1.48	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4	PF5	CB-CG-CD2	-3.32	117.49	122.03
1	A	4	PF5	CB-CA-C	-3.31	105.27	111.47
1	B	4	PF5	CB-CG-CD1	2.60	125.59	122.03
1	A	4	PF5	CB-CG-CD1	-2.33	118.85	122.03
1	B	4	PF5	CB-CA-C	-2.16	107.43	111.47
1	A	4	PF5	CB-CG-CD2	2.05	124.84	122.03

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	4	PF5	CA-CB-CG-CD1
1	A	4	PF5	CA-CB-CG-CD2
1	B	4	PF5	CA-CB-CG-CD1
1	B	4	PF5	CA-CB-CG-CD2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	4	PF5	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	169/170 (99%)	0.85	12 (7%)	22 30	9, 17, 27, 42	2 (1%)
1	B	169/170 (99%)	2.05	84 (49%)	0 1	15, 29, 48, 71	1 (0%)
All	All	338/340 (99%)	1.45	96 (28%)	1 3	9, 21, 43, 71	3 (0%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	110	PHE	5.7
1	B	146	MET	5.3
1	B	145	GLY	5.1
1	B	169	HIS	4.9
1	B	91	PRO	4.6
1	B	90	ALA	4.5
1	B	160[A]	VAL	4.4
1	B	115	LEU	4.3
1	B	2	VAL	4.1
1	B	23	THR	4.0
1	B	111	SER	3.9
1	B	165	HIS	3.9
1	B	16	PHE	3.9
1	B	139	VAL	3.9
1	B	28	LEU	3.9
1	B	22	GLU	3.8
1	B	170	HIS	3.8
1	B	143	ARG	3.6
1	B	44	VAL	3.5
1	B	132	VAL	3.5
1	B	67	PRO	3.5
1	A	146	MET	3.4
1	B	24	VAL	3.3
1	B	106	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	156	ILE	3.2
1	A	33	GLU	3.2
1	B	168	HIS	3.2
1	B	76	LEU	3.1
1	B	114	SER	3.1
1	B	159[A]	SER	3.1
1	B	89	GLN	3.1
1	B	72	ALA	3.0
1	A	145	GLY	3.0
1	B	116	GLN	2.9
1	B	163	SER	2.9
1	A	148	GLN	2.9
1	B	15	THR	2.9
1	B	45	ILE	2.8
1	B	66	GLU	2.8
1	B	144	SER	2.7
1	B	152	LYS	2.7
1	B	147	HIS	2.7
1	B	27	PHE	2.7
1	B	6	THR	2.6
1	A	103	VAL	2.6
1	A	116	GLN	2.6
1	A	76	LEU	2.6
1	B	162	VAL	2.6
1	B	88	THR	2.6
1	B	31	CYS	2.5
1	B	60	LYS	2.5
1	B	17	ASP	2.5
1	B	107	PHE	2.5
1	B	123	PHE	2.5
1	B	40	ILE	2.4
1	B	21	PRO	2.4
1	B	55	PHE	2.4
1	B	94	ALA	2.4
1	B	153	GLU	2.4
1	B	20	ALA	2.3
1	B	138	GLY	2.3
1	B	118	TRP	2.3
1	B	81	GLY	2.3
1	B	142	GLY	2.3
1	B	86	ALA	2.3
1	B	25	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	115	LEU	2.3
1	B	3	THR	2.3
1	B	18	ASP	2.3
1	B	74	ASN	2.2
1	A	138	GLY	2.2
1	B	34	GLY	2.2
1	B	46	ASN	2.2
1	B	1	MET	2.2
1	B	48	PHE	2.2
1	B	141	THR	2.2
1	B	13	ILE	2.2
1	B	135	LYS	2.2
1	B	164	GLU	2.1
1	B	148	GLN	2.1
1	A	2[A]	VAL	2.1
1	B	7	ASN	2.1
1	B	37	ASN	2.1
1	B	57	PRO	2.1
1	B	124	ALA	2.1
1	B	140	ALA	2.1
1	B	68	ILE	2.1
1	B	131	ASP	2.1
1	B	121	CYS	2.1
1	B	58	GLY	2.1
1	A	106	ASP	2.0
1	B	9	GLY	2.0
1	B	109	ASN	2.0
1	A	163	SER	2.0
1	B	84	ALA	2.0
1	B	41	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PF5	B	4	16/17	0.88	0.11	20,23,28,29	0
1	PF5	A	4	16/17	0.96	0.06	11,13,14,15	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.