Full wwPDB EM Validation Report ⓘ

Oct 26, 2026 – 04:30 PM EDT

PDB ID	:	37XV / pdb_000037xv
EMDB ID	:	EMD-78609
Title	:	Polynucleotide kinase phosphatase bound to a nucleosome containing a nick with a 3'-phosphate and 5'-OH at SHL-5 (catalytic, composite)
Authors	:	Boesch, D.J.; Evans, J.J.; Weaver, T.M.
Deposited on	:	2026-08-12
Resolution	:	3.20 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	62% 9% 29%
1	E	135	65% 7% 28%
2	B	102	76% 20%
2	F	102	75% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	129	
3	G	129	
4	D	125	
4	H	125	
5	I	147	
6	J	24	
7	K	123	
8	L	528	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 14915 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	96	Total	C	N	O	S	0	0
			789	499	151	137	2		
1	E	97	Total	C	N	O	S	0	0
			800	505	155	138	2		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	CYS	engineered mutation	UNP Q71DI3
E	110	ALA	CYS	engineered mutation	UNP Q71DI3

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	82	Total	C	N	O	S	0	0
			657	416	128	112	1		
2	F	80	Total	C	N	O	S	0	0
			641	405	125	110	1		

- Molecule 3 is a protein called Histone H2A type 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	107	Total	C	N	O	0	0
			823	520	161	142		
3	G	107	Total	C	N	O	0	0
			823	520	161	142		

- Molecule 4 is a protein called Histone H2B type 1-C/E/F/G/I.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	94	Total	C	N	O	S	0	0
			736	461	134	139	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	93	Total	C	N	O	S	0	0
			725	455	130	138	2		

- Molecule 5 is a DNA chain called 601 I strand (non-damaged strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	147	Total	C	N	O	P	0	0
			2996	1423	545	881	147		

- Molecule 6 is a DNA chain called 601 J strand (damaged strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	24	Total	C	N	O	P	0	0
			497	236	88	148	25		

- Molecule 7 is a DNA chain called 601 K strand (damaged strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	K	123	Total	C	N	O	P	0	0
			2535	1199	478	736	122		

- Molecule 8 is a protein called Bifunctional polynucleotide phosphatase/kinase.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	L	376	Total	C	N	O	S	0	0
			2893	1850	513	517	13		

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
L	-6	MET	-	initiating methionine	UNP Q96T60
L	-5	HIS	-	expression tag	UNP Q96T60
L	-4	HIS	-	expression tag	UNP Q96T60
L	-3	HIS	-	expression tag	UNP Q96T60
L	-2	HIS	-	expression tag	UNP Q96T60
L	-1	HIS	-	expression tag	UNP Q96T60
L	0	HIS	-	expression tag	UNP Q96T60
L	1	GLY	-	expression tag	UNP Q96T60
L	171	ALA	ASP	engineered mutation	UNP Q96T60

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone H3.2

Chain A:




• Molecule 1: Histone H3.2

Chain E:




• Molecule 2: Histone H4

Chain B:



• Molecule 2: Histone H4

Chain F:




• Molecule 3: Histone H2A type 1


Chain C:






• Molecule 3: Histone H2A type 1






• Molecule 4: Histone H2B type 1-C/E/F/G/I







• Molecule 4: Histone H2B type 1-C/E/F/G/I






• Molecule 5: 601 I strand (non-damaged strand)






• Molecule 6: 601 J strand (damaged strand)





• Molecule 7: 601 K strand (damaged strand)






• Molecule 8: Bifunctional polynucleotide phosphatase/kinase

Chain L:


27%
67%
5%
29%


MET HIS HIS HIS HIS HIS GLY GLY GLU VAL GLU ALA PRO GLY ARG LEU TRP LEU GLU SER PRO PRO GLY GLY ALA PRO PRO ILE PHE LEU PRO SER ASP GLY GLN ALA LEU VAL LEU GLY ARG GLY PRO LEU THR GLN VAL THR ASP ARG LYS CYS SER ARG THR GLN VAL GLU LEU
VAL ALA ASP PRO GLU THR ARG THR VAL ALA VAL LYS GLN LEU GLY VAL ASN PRO SER THR THR GLY THR GLN GLU LEU LYS PRO GLY LEU GLU GLY SER LEU GLY VAL GLY ASP THR LEU TYR LEU VAL ASN GLY LEU HIS PRO LEU THR LEU ARG TRP GLU GLU THR ARG THR PRO GLU
SER GLN PRO ASP THR PRO PRO GLY THR PRO LEU VAL SER GLN ASP GLU LYS ARG ASP ALA GLU LEU PRO LYS LYS ARG MET ARG LYS SER ASN PRO G146 W147 E148 N149 L154 A158 A159 G160 V167 L172 D173 G174 T175 L176 I177 T178 T179 R180 S181 G182 P186 T187 G188
P189 S190 D191 I194 L195 Y196 P197 R201 K202 L203 A208 E209 K212 L213 F216 R224 G225 K226 L227 P228 A229 E230 E231 F232 E237 A238 V239 V240 E241 K242 L243 G244 V245 P246 F247 Q248 V249 L250 V251 T263 E271 Q272 A273 N274 D275 G276 T277 P278 I281 G282
D283 F286 V287 G288 D289 A290 A291 A295 N296 P299 G300 R301 K302 F306 R311 L312 F313 L317 G318 L319 K330 W331 P332 A333 A334 P339 A340 F341 V346 S347 R348 S349 G350 P351 L352 C353 L354 P355 E356 S357 R358 A359 L360 L361 S362 A363 S364 V369 A370
G375 A376 S388 A389 H393 D397 T408 K414 Q415 G416 R433 Y434 V435 A438 R439 A440 A441 G442 V443 F448 N460 R464 D468 S469 S470 H471 I472 P473 V474 A488 P489 T490 L491 A492 E493 G494 F495 S496 A497 I498 L499 E500 I501 P502 F503 R504 E508
P509 R510 L511 S519 E520 G521

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	12103	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	27.702	Depositor
Minimum map value	-12.416	Depositor
Average map value	-0.030	Depositor
Map value standard deviation	0.958	Depositor
Recommended contour level	5	Depositor
Map size (Å)	260.80002, 260.80002, 260.80002	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.65200007, 0.65200007, 0.65200007	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: THP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.15	0/801	0.31	0/1075
1	E	0.14	0/812	0.27	0/1089
2	B	0.15	0/664	0.30	0/889
2	F	0.14	0/648	0.28	0/868
3	C	0.12	0/833	0.25	0/1124
3	G	0.13	0/833	0.26	0/1124
4	D	0.12	0/747	0.28	0/1004
4	H	0.12	0/736	0.26	0/990
5	I	0.73	1/3357 (0.0%)	0.93	3/5174 (0.1%)
6	J	0.75	0/530	0.93	0/816
7	K	0.70	0/2847	0.84	0/4397
8	L	0.10	0/2969	0.25	0/4030
All	All	0.48	1/15777 (0.0%)	0.64	3/22580 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	128	DG	C3'-O3'	6.15	1.51	1.44

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	I	128	DG	P-O3'-C3'	6.05	126.96	119.70
5	I	126	DG	O4'-C4'-C3'	-5.40	102.76	106.00
5	I	95	DC	O4'-C4'-C3'	-5.12	102.93	106.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	789	0	826	8	0
1	E	800	0	839	9	0
2	B	657	0	706	3	0
2	F	641	0	684	3	0
3	C	823	0	884	5	0
3	G	823	0	884	6	0
4	D	736	0	756	4	0
4	H	725	0	743	3	0
5	I	2996	0	1650	18	0
6	J	497	0	272	5	0
7	K	2535	0	1380	21	0
8	L	2893	0	2862	13	0
All	All	14915	0	12486	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:PRO:HB3	2:B:53:GLU:HG3	1.80	0.64
5:I:17:DC:H2"	5:I:18:DC:C5	2.40	0.57
8:L:281:ILE:HG23	8:L:319:LEU:HD22	1.86	0.56
6:J:1:DA:H2"	6:J:2:DT:H5'	1.89	0.54
8:L:510:ARG:HG3	8:L:511:LEU:HG	1.89	0.54
7:K:82:DG:H2'	7:K:83:DT:C6	2.43	0.53
7:K:104:DC:H2'	7:K:105:DT:H71	1.90	0.53
7:K:35:DT:H1'	7:K:36:DA:H5'	1.92	0.52
5:I:130:DC:H2"	5:I:131:DA:N7	2.24	0.52
1:A:50:GLU:HA	1:A:53:ARG:HB2	1.90	0.52
5:I:144:DC:H2"	5:I:145:DG:C8	2.45	0.52
7:K:82:DG:H3'	7:K:83:DT:H71	1.91	0.51
5:I:47:DC:H2"	5:I:48:DT:H72	1.93	0.50
6:J:21:DA:H2'	6:J:22:DC:C5	2.47	0.50
5:I:96:DC:H2'	5:I:97:DA:C8	2.47	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:L:317:LEU:HB3	8:L:319:LEU:HD23	1.95	0.49
3:G:63:LEU:HD13	4:H:45:LEU:HB2	1.94	0.49
5:I:28:DC:H2"	5:I:29:DA:C8	2.47	0.49
1:E:61:LEU:HD12	2:F:37:LEU:HD23	1.95	0.49
7:K:89:DT:H2"	7:K:90:DA:C8	2.47	0.49
8:L:263:THR:HA	8:L:313:PHE:HE1	1.78	0.48
5:I:146:DA:H2"	5:I:147:DT:H5'	1.94	0.48
6:J:7:DT:H2'	6:J:8:DG:C8	2.48	0.48
1:A:79:LYS:HD3	1:A:82:LEU:HD13	1.95	0.48
3:C:102:ILE:HG23	4:D:61:ILE:HD13	1.95	0.48
6:J:3:DC:H2"	6:J:4:DG:C8	2.48	0.48
3:G:54:VAL:HG21	4:H:98:VAL:HG21	1.96	0.48
7:K:111:DC:H2"	7:K:112:DG:C8	2.48	0.48
1:A:108:ASN:HB2	2:B:43:VAL:HG22	1.96	0.47
5:I:82:DC:H2"	5:I:83:DG:C8	2.50	0.47
6:J:9:DT:H2"	6:J:10:DA:H5"	1.96	0.47
7:K:54:DC:H2'	7:K:55:DG:C8	2.50	0.47
8:L:355:PRO:HD2	8:L:358:ARG:HD2	1.97	0.46
7:K:138:DG:H1'	7:K:139:DA:H5'	1.97	0.46
1:E:46:VAL:HG21	5:I:83:DG:H3'	1.97	0.46
7:K:123:DC:H2"	7:K:124:DG:C8	2.50	0.46
7:K:137:DG:H2"	7:K:138:DG:C8	2.51	0.46
7:K:55:DG:H2"	7:K:56:DG:C8	2.51	0.45
1:A:62:ILE:HD11	2:B:37:LEU:HD11	1.98	0.45
3:G:95:LYS:HD2	4:H:103:PRO:HB3	1.98	0.45
1:A:113:HIS:CG	1:E:126:LEU:HD22	2.52	0.45
4:D:46:LYS:HA	4:D:46:LYS:HD3	1.72	0.45
5:I:121:DG:H2"	5:I:122:DG:C8	2.51	0.45
7:K:145:DG:H2"	7:K:146:DA:H5"	1.98	0.44
8:L:312:LEU:HD11	8:L:339:PRO:HG2	1.99	0.44
8:L:291:ALA:HB1	8:L:311:ARG:HH21	1.83	0.44
5:I:124:DA:H2"	5:I:125:DC:O2	2.17	0.44
8:L:203:LEU:HD22	8:L:213:LEU:HD13	1.99	0.44
3:C:63:LEU:HD13	4:D:45:LEU:HB2	1.99	0.44
3:C:44:GLY:HA2	7:K:112:DG:H5"	2.00	0.43
7:K:28:DT:H2"	7:K:29:DG:N7	2.33	0.43
5:I:57:DT:H2"	5:I:58:DT:C6	2.53	0.43
8:L:213:LEU:HB3	8:L:247:PHE:CE1	2.53	0.43
7:K:30:DG:H2"	7:K:31:DA:H8	1.83	0.43
5:I:60:DA:H2"	5:I:61:DA:C8	2.54	0.43
3:C:88:ARG:HA	3:C:88:ARG:HD3	1.82	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:K:41:DG:H2'	7:K:42:DT:H71	2.01	0.42
5:I:108:DT:H2"	5:I:109:DC:C6	2.55	0.42
5:I:137:DA:H2'	5:I:138:DT:H71	2.01	0.42
8:L:194:ILE:HG12	8:L:242:LYS:HE3	2.01	0.42
8:L:448:PHE:CE1	8:L:499:LEU:HD12	2.54	0.42
8:L:460:ASN:O	8:L:464:ARG:HG2	2.20	0.42
1:E:43:PRO:HA	5:I:83:DG:H5'	2.02	0.42
3:C:107:VAL:HG11	1:E:101:VAL:HG11	2.03	0.41
1:E:92:LEU:HD23	1:E:92:LEU:HA	1.94	0.41
7:K:128:DT:H2"	7:K:129:DC:C5	2.55	0.41
2:F:90:LEU:HB3	2:F:95:ARG:HB2	2.02	0.41
8:L:167:VAL:HG22	8:L:212:LYS:HB3	2.02	0.41
7:K:70:DG:H2"	7:K:71:DA:C8	2.56	0.41
3:G:16:THR:HA	7:K:31:DA:H5"	2.02	0.41
1:A:126:LEU:HD22	1:E:113:HIS:CG	2.56	0.41
4:D:33:ARG:HG3	7:K:123:DC:H4'	2.03	0.40
5:I:111:DC:H6	5:I:111:DC:H2'	1.69	0.40
5:I:66:DC:H2"	5:I:67:DG:C8	2.56	0.40
3:G:63:LEU:HD23	3:G:63:LEU:HA	1.92	0.40
3:G:65:LEU:HD12	3:G:93:LEU:HD12	2.02	0.40
1:A:49:ARG:C	1:A:51:ILE:H	2.29	0.40
1:E:85:GLN:HG3	7:K:50:DT:OP1	2.21	0.40
1:E:108:ASN:HB2	2:F:43:VAL:HG22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/135 (70%)	91 (97%)	3 (3%)	0	100	100
1	E	95/135 (70%)	93 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	80/102 (78%)	77 (96%)	3 (4%)	0	100	100
2	F	78/102 (76%)	75 (96%)	3 (4%)	0	100	100
3	C	105/129 (81%)	105 (100%)	0	0	100	100
3	G	105/129 (81%)	104 (99%)	1 (1%)	0	100	100
4	D	92/125 (74%)	90 (98%)	2 (2%)	0	100	100
4	H	91/125 (73%)	88 (97%)	3 (3%)	0	100	100
8	L	374/528 (71%)	368 (98%)	6 (2%)	0	100	100
All	All	1114/1510 (74%)	1091 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/109 (76%)	82 (99%)	1 (1%)	63	79
1	E	84/109 (77%)	84 (100%)	0	100	100
2	B	68/78 (87%)	67 (98%)	1 (2%)	57	76
2	F	66/78 (85%)	66 (100%)	0	100	100
3	C	83/98 (85%)	83 (100%)	0	100	100
3	G	83/98 (85%)	83 (100%)	0	100	100
4	D	81/105 (77%)	81 (100%)	0	100	100
4	H	80/105 (76%)	79 (99%)	1 (1%)	61	78
8	L	297/435 (68%)	297 (100%)	0	100	100
All	All	925/1215 (76%)	922 (100%)	3 (0%)	84	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	92	ARG
4	H	63	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
1	A	108	ASN
2	B	75	HIS
1	E	108	ASN
3	G	84	GLN
4	H	49	HIS
4	H	95	GLN
8	L	296	ASN
8	L	415	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	THP	J	24	6	22,25,26	0.81	0	32,37,40	0.95	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	THP	J	24	6	-	7/12/26/27	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	J	24	THP	O2P-P1-O1P	2.54	120.74	110.83
6	J	24	THP	O4-C4-C5	-2.24	122.35	124.92

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	J	24	THP	O4'-C1'-N1-C2
6	J	24	THP	O4'-C1'-N1-C6
6	J	24	THP	O4'-C4'-C5'-O5'
6	J	24	THP	C3'-C4'-C5'-O5'
6	J	24	THP	C2'-C1'-N1-C6
6	J	24	THP	C4'-C5'-O5'-P2
6	J	24	THP	C2'-C1'-N1-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-78609. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices


6.2.1 Primary map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 185



Y Index: 216




Z Index: 179


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 5.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 129 nm^3; this corresponds to an approximate mass of 116 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.312 Å^{-1}


8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-78609 and PDB model 37XV. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 5.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (5).

9.4 Atom inclusion (i)



At the recommended contour level, 86% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8150	0.3980
A	0.8740	0.5220
B	0.8920	0.5290
C	0.9000	0.5220
D	0.8880	0.5090
E	0.9080	0.5310
F	0.9200	0.5360
G	0.8400	0.5030
H	0.8660	0.4940
I	0.8840	0.4040
J	0.4770	0.1590
K	0.9670	0.4430
L	0.5200	0.1430

