wwPDB EM Validation Summary Report ⓘ

Oct 26, 2026 – 04:13 PM EDT

PDB ID	:	37XX / pdb_000037xx
EMDB ID	:	EMD-78611
Title	:	Polynucleotide kinase phosphatase bound to a nucleosome containing a nick with a 3'-phosphate and 5'-OH at SHL-5 (FHA, composite)
Authors	:	Boesch, D.J.; Evans, J.J.; Weaver, T.M.
Deposited on	:	2026-08-12
Resolution	:	3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	65% 6% 29%
1	E	135	71% 28%
2	B	102	75% 6% 20%
2	F	102	76% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	129	
3	G	129	
4	D	125	
4	H	125	
5	I	147	
6	J	24	
7	K	123	
8	L	528	

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 12776 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	96	Total	C	N	O	S	0	0
			789	499	151	137	2		
1	E	97	Total	C	N	O	S	1	0
			808	510	158	138	2		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	CYS	engineered mutation	UNP Q71DI3
E	110	ALA	CYS	engineered mutation	UNP Q71DI3

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	82	Total	C	N	O	S	0	0
			657	416	128	112	1		
2	F	80	Total	C	N	O	S	0	0
			641	405	125	110	1		

- Molecule 3 is a protein called Histone H2A type 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	107	Total	C	N	O	0	0
			823	520	161	142		
3	G	107	Total	C	N	O	0	0
			823	520	161	142		

- Molecule 4 is a protein called Histone H2B type 1-C/E/F/G/I.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	94	Total	C	N	O	S	1	0
			743	466	136	139	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	93	Total	C	N	O	S	1	0
			732	460	132	138	2		

- Molecule 5 is a DNA chain called 601 I strand (non-damaged strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	147	Total	C	N	O	P	0	0
			2996	1423	545	881	147		

- Molecule 6 is a DNA chain called 601 J strand (damaged strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	24	Total	C	N	O	P	0	0
			497	236	88	148	25		

- Molecule 7 is a DNA chain called 601 K strand (damaged strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	K	123	Total	C	N	O	P	0	0
			2535	1199	478	736	122		

- Molecule 8 is a protein called Bifunctional polynucleotide phosphatase/kinase.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	L	99	Total	C	N	O	S	0	0
			732	464	131	136	1		

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	-6	MET	-	initiating methionine	UNP Q96T60
L	-5	HIS	-	expression tag	UNP Q96T60
L	-4	HIS	-	expression tag	UNP Q96T60
L	-3	HIS	-	expression tag	UNP Q96T60
L	-2	HIS	-	expression tag	UNP Q96T60
L	-1	HIS	-	expression tag	UNP Q96T60
L	0	HIS	-	expression tag	UNP Q96T60
L	1	GLY	-	expression tag	UNP Q96T60
L	171	ALA	ASP	engineered mutation	UNP Q96T60


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Histone H3.2

Chain A:






• Molecule 1: Histone H3.2

Chain E:






• Molecule 2: Histone H4

Chain B:






• Molecule 2: Histone H4

Chain F:





• Molecule 3: Histone H2A type 1


Chain C:






• Molecule 3: Histone H2A type 1

Chain G:




• Molecule 4: Histone H2B type 1-C/E/F/G/I

Chain D:




• Molecule 4: Histone H2B type 1-C/E/F/G/I

Chain H:




• Molecule 5: 601 I strand (non-damaged strand)

Chain I:




• Molecule 6: 601 J strand (damaged strand)

Chain J:



• Molecule 7: 601 K strand (damaged strand)


Chain K:



• Molecule 8: Bifunctional polynucleotide phosphatase/kinase

Chain L:




MET HIS HIS HIS HIS HIS HIS GLY GLY GLU VAL GLU ALA PRO G8 P16 G17 G18 S26 D27 G28 Q29 A30 L31 V32 L33 P37 L38 T39 Q40 K45 C46 L53 V54 A55 D56 P57 E58 T59 R60 T61 S72 G75 T76 K80 E84 L87 G88 V89 G90
L93 W106 GLU GLU THR ARG THR PRO GLU SER GLN PRO ASP THR PRO PRO GLY THR PRO LEU VAL SER GLN ASP GLU LYS ARG ASP ALA GLU LEU PRO LYS LYS ARG MET ARG LYS SER ASN PRO GLY TRP GLU ASN LEU GLU LYS LEU LEU VAL PHE THR ALA ALA GLY VAL LYS
PRO GLN GLY LYS VAL ALA GLY PHE ALA LEU ASP GLY THR LEU ILE THR THR ARG SER GLY LYS VAL PHE PRO THR GLY PRO SER ASP TRP ARG ILE LEU TYR PRO GLU ILE PRO ARG LYS LEU ARG GLU LEU GLU LEU VAL ILE PHE THR ASN GLN MET SER ILE
GLY ARG GLY LYS LEU PRO ALA GLU GLU PHE LYS ALA LYS VAL GLU ALA VAL VAL GLU LYS LEU GLY VAL PRO PHE GLN VAL LEU VAL ALA THR HIS ALA GLY LEU TYR ARG LYS PRO VAL THR GLY MET TRP ASP HIS LEU GLN GLU GLN ALA ASN ASP GLY THR PRO ILE SER ILE GLY
ASP SER ILE PHE VAL GLY ASP ALA ALA GLY ARG PRO ALA ASN TRP ALA PRO GLY ARG LYS LYS LYS ASP PHE SER CYS ALA ASP ARG LEU PHE ALA LEU ASN LEU GLY LEU PRO PHE ALA THR PRO GLU GLU PHE PHE LEU LYS TRP PRO ALA ALA GLY PHE GLU LEU PRO ALA ASP PHE
PRO ARG THR VAL SER ARG SER GLY PRO LEU CYS LEU PRO GLU SER ARG ALA LEU LEU SER ALA SER PRO GLU VAL VAL VAL ALA VAL GLY PHE PRO GLY ALA GLY LYS SER THR PHE LEU LYS LYS HIS LEU VAL SER ALA GLY TYR VAL HIS VAL ASN ARG ASP THR LEU GLY SER TRP
GLN ARG CYS VAL THR THR CYS GLU THR ALA LEU LYS GLN GLY LYS ARG VAL ALA ILE ASP ASN THR ASN PRO ASP ALA ALA ALA SER ARG ALA ARG TYR VAL GLN CYS ALA ARG ALA ALA GLY VAL PRO CYS ARG CYS PHE LEU PHE THR ALA THR LEU GLU GLN ALA ARG HIS ASN ASN ARG
PHE ARG GLU MET THR ASP SER SER HIS ILE PRO VAL SER ASP MET VAL MET TYR GLY TYR ARG LYS GLN PHE GLU ALA PRO THR LEU ALA GLU GLY PHE SER ALA ILE LEU GLU ILE PRO PHE ARG LEU TRP VAL GLU PRO ARG LEU GLY ARG LEU TYR CYS GLN PHE SER GLU GLY

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	33821	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	32.615	Depositor
Minimum map value	-14.346	Depositor
Average map value	0.002	Depositor
Map value standard deviation	1.004	Depositor
Recommended contour level	5.8	Depositor
Map size (Å)	260.80002, 260.80002, 260.80002	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.65200007, 0.65200007, 0.65200007	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: THP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.27	0/801	0.37	0/1075
1	E	0.26	0/823	0.30	0/1103
2	B	0.29	0/664	0.37	0/889
2	F	0.28	0/648	0.34	0/868
3	C	0.25	0/833	0.28	0/1124
3	G	0.24	0/833	0.30	0/1124
4	D	0.25	0/758	0.30	0/1019
4	H	0.25	0/747	0.31	0/1005
5	I	0.83	0/3357	0.92	0/5174
6	J	0.83	0/530	1.02	1/816 (0.1%)
7	K	0.80	0/2847	0.86	0/4397
8	L	0.13	0/748	0.37	0/1024
All	All	0.60	0/13589	0.70	1/19618 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	J	19	DA	O4'-C4'-C3'	-5.24	102.85	106.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	789	0	826	5	0
1	E	808	0	852	1	0
2	B	657	0	706	4	0
2	F	641	0	684	1	0
3	C	823	0	884	5	0
3	G	823	0	884	4	0
4	D	743	0	763	6	0
4	H	732	0	750	5	0
5	I	2996	0	1650	16	0
6	J	497	0	272	4	0
7	K	2535	0	1380	14	0
8	L	732	0	750	1	0
All	All	12776	0	10401	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:PRO:HB3	2:B:53:GLU:HG3	1.82	0.62
5:I:28:DC:H2"	5:I:29:DA:C8	2.35	0.61
5:I:119:DC:H2"	5:I:120:DA:C8	2.35	0.61
1:A:50:GLU:HA	1:A:53:ARG:HB2	1.83	0.60
1:A:62:ILE:HD11	2:B:37:LEU:HD11	1.87	0.56

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/135 (70%)	94 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	96/135 (71%)	96 (100%)	0	0	100	100
2	B	80/102 (78%)	80 (100%)	0	0	100	100
2	F	78/102 (76%)	78 (100%)	0	0	100	100
3	C	105/129 (81%)	105 (100%)	0	0	100	100
3	G	105/129 (81%)	105 (100%)	0	0	100	100
4	D	93/125 (74%)	93 (100%)	0	0	100	100
4	H	92/125 (74%)	92 (100%)	0	0	100	100
8	L	97/528 (18%)	97 (100%)	0	0	100	100
All	All	840/1510 (56%)	840 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/109 (76%)	82 (99%)	1 (1%)	63	82
1	E	85/109 (78%)	85 (100%)	0	100	100
2	B	68/78 (87%)	67 (98%)	1 (2%)	57	80
2	F	66/78 (85%)	66 (100%)	0	100	100
3	C	83/98 (85%)	83 (100%)	0	100	100
3	G	83/98 (85%)	83 (100%)	0	100	100
4	D	82/105 (78%)	82 (100%)	0	100	100
4	H	81/105 (77%)	81 (100%)	0	100	100
8	L	80/435 (18%)	80 (100%)	0	100	100
All	All	711/1215 (58%)	709 (100%)	2 (0%)	84	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	THR
2	B	92	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
4	H	49	HIS
4	H	67	ASN
3	C	112	GLN
1	E	108	ASN
3	G	73	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	THP	J	24	6	22,25,26	0.52	0	32,37,40	1.01	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	THP	J	24	6	-	3/12/26/27	0/2/2/2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	J	24	THP	O4-C4-C5	-2.56	121.99	124.92
6	J	24	THP	O2P-P1-O1P	2.28	119.71	110.83
6	J	24	THP	O3P-P1-O3'	-2.16	97.45	105.85

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	J	24	THP	O4'-C4'-C5'-O5'
6	J	24	THP	C3'-C4'-C5'-O5'
6	J	24	THP	C3'-O3'-P1-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-78611. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices


6.2.1 Primary map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 182



Y Index: 215




Z Index: 179


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 5.8. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 99 nm^3; this corresponds to an approximate mass of 90 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.333 Å^{-1}


8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-78611 and PDB model 37XX. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 5.8 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (5.8).

9.4 Atom inclusion (i)



At the recommended contour level, 92% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (5.8) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8770	0.4930
A	0.8820	0.5500
B	0.8860	0.5520
C	0.8830	0.5550
D	0.8910	0.5430
E	0.9020	0.5610
F	0.9020	0.5620
G	0.8620	0.5460
H	0.8770	0.5350
I	0.9240	0.4930
J	0.6120	0.2460
K	0.9570	0.5090
L	0.5230	0.1390

