



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 2, 2024 – 10:51 PM EDT

PDB ID : 3ABZ  
Title : Crystal structure of Se-Met labeled Beta-glucosidase from Kluyveromyces marxianus  
Authors : Yoshida, E.; Hidaka, M.; Fushinobu, S.; Katayama, T.; Kumagai, H.  
Deposited on : 2009-12-25  
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	 88% 10% .
1	B	845	 3% 81% 13% . .
1	C	845	 85% 11% . .
1	D	845	 85% 12% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	5002	-	-	X	-
2	GOL	B	5006	-	-	X	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 29154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase I.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	844	Total	C	N	O	S	Se	0	0	0
			6606	4197	1110	1284	5	10			
1	B	810	Total	C	N	O	S	Se	0	0	0
			6351	4040	1065	1232	4	10			
1	C	836	Total	C	N	O	S	Se	0	0	0
			6550	4161	1102	1272	5	10			
1	D	833	Total	C	N	O	S	Se	0	0	0
			6521	4146	1096	1264	5	10			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	875	Total	O	0	0
			875	875		
3	B	828	Total	O	0	0
			828	828		
3	C	701	Total	O	0	0
			701	701		
3	D	674	Total	O	0	0
			674	674		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Beta-glucosidase I



#### • Molecule 1: Beta-glucosidase I



#### • Molecule 1: Beta-glucosidase I



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	245.68Å 148.64Å 119.78Å 90.00° 112.95° 90.00°	Depositor
Resolution (Å)	50.00 – 2.15 50.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.15) 99.8 (50.00-2.15)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.03 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.160 , 0.218 0.156 , 0.213	Depositor DCC
$R_{free}$ test set	10724 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.4	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	29154	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
 GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.07	1/6733 (0.0%)	0.88	8/9088 (0.1%)
1	B	1.13	7/6466 (0.1%)	0.90	11/8720 (0.1%)
1	C	1.01	3/6675 (0.0%)	0.85	6/9008 (0.1%)
1	D	0.99	4/6645 (0.1%)	0.84	5/8967 (0.1%)
All	All	1.05	15/26519 (0.1%)	0.87	30/35783 (0.1%)

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	795	GLU	CG-CD	5.84	1.60	1.51
1	A	643	TYR	CD1-CE1	-5.67	1.30	1.39
1	B	80	GLU	CB-CG	-5.64	1.41	1.52
1	C	199	GLU	CB-CG	5.53	1.62	1.52
1	C	771	VAL	CB-CG1	5.47	1.64	1.52

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	113	ARG	NE-CZ-NH1	-8.69	115.95	120.30
1	A	769	ARG	NE-CZ-NH1	-7.39	116.60	120.30
1	C	769	ARG	NE-CZ-NH1	-7.34	116.63	120.30
1	C	251	ARG	NE-CZ-NH1	-7.02	116.79	120.30
1	B	263	ARG	NE-CZ-NH2	6.89	123.75	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6606	0	6514	70	0
1	B	6351	0	6279	89	0
1	C	6550	0	6460	74	0
1	D	6521	0	6439	65	0
2	A	24	0	32	10	0
2	B	12	0	16	5	0
2	C	6	0	8	0	0
2	D	6	0	8	0	0
3	A	875	0	0	13	0
3	B	828	0	0	18	0
3	C	701	0	0	15	0
3	D	674	0	0	10	0
All	All	29154	0	25756	296	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 296 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:5001:GOL:H32	3:A:1983:HOH:O	1.54	1.05
1:C:734:ASP:O	1:C:735:ASP:HB3	1.57	1.04
1:A:737:ILE:HD11	1:A:796:LEU:HD23	1.41	1.01
1:C:338:ASN:HB3	1:C:575:ASP:HB2	1.53	0.91
1:D:134:LYS:HE2	3:D:2595:HOH:O	1.70	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	842/845 (100%)	808 (96%)	33 (4%)	1 (0%)	48	51
1	B	798/845 (94%)	764 (96%)	33 (4%)	1 (0%)	48	51
1	C	832/845 (98%)	796 (96%)	36 (4%)	0	100	100
1	D	827/845 (98%)	789 (95%)	38 (5%)	0	100	100
All	All	3299/3380 (98%)	3157 (96%)	140 (4%)	2 (0%)	48	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	524	GLY
1	B	405	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	716/706 (101%)	693 (97%)	23 (3%)	34	34
1	B	690/706 (98%)	670 (97%)	20 (3%)	37	38
1	C	710/706 (101%)	674 (95%)	36 (5%)	20	16
1	D	707/706 (100%)	678 (96%)	29 (4%)	26	25
All	All	2823/2824 (100%)	2715 (96%)	108 (4%)	28	27

5 of 108 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	487	PHE
1	C	736	LYS
1	D	708	ARG
1	C	508	PHE
1	C	633	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	388	HIS
1	D	443	HIS
1	B	388	HIS
1	B	338	ASN
1	D	525	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	5003	-	5,5,5	0.35	0	5,5,5	0.65	0
2	GOL	B	5005	-	5,5,5	0.56	0	5,5,5	1.07	0
2	GOL	C	5007	-	5,5,5	0.59	0	5,5,5	0.77	0
2	GOL	A	5001	-	5,5,5	0.47	0	5,5,5	0.75	0
2	GOL	B	5006	-	5,5,5	0.31	0	5,5,5	0.39	0
2	GOL	D	5008	-	5,5,5	0.51	0	5,5,5	1.02	0
2	GOL	A	5004	-	5,5,5	0.44	0	5,5,5	0.86	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	5002	-	5,5,5	0.82	0	5,5,5	1.78	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	5003	-	-	0/4/4/4	-
2	GOL	B	5005	-	-	2/4/4/4	-
2	GOL	C	5007	-	-	2/4/4/4	-
2	GOL	A	5001	-	-	2/4/4/4	-
2	GOL	B	5006	-	-	4/4/4/4	-
2	GOL	D	5008	-	-	2/4/4/4	-
2	GOL	A	5004	-	-	4/4/4/4	-
2	GOL	A	5002	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5002	GOL	C3-C2-C1	-3.24	99.91	111.80

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	5001	GOL	O2-C2-C3-O3
2	A	5002	GOL	C1-C2-C3-O3
2	A	5004	GOL	O1-C1-C2-C3
2	B	5006	GOL	O1-C1-C2-C3
2	B	5006	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5003	GOL	1	0
2	B	5005	GOL	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5001	GOL	1	0
2	B	5006	GOL	4	0
2	A	5002	GOL	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	834/845 (98%)	-0.67	4 (0%) 87 89	12, 25, 51, 69	0
1	B	800/845 (94%)	-0.50	28 (3%) 47 53	12, 22, 67, 98	0
1	C	826/845 (97%)	-0.26	8 (0%) 79 82	14, 32, 65, 94	0
1	D	823/845 (97%)	-0.24	7 (0%) 81 83	14, 34, 58, 73	0
All	All	3283/3380 (97%)	-0.42	47 (1%) 73 77	12, 28, 61, 98	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	445	PHE	4.9
1	B	408	ALA	4.3
1	B	538	THR	4.1
1	B	406	ALA	3.8
1	B	357	ALA	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	C	5007	6/6	0.84	0.12	33,38,40,46	0
2	GOL	D	5008	6/6	0.87	0.11	35,37,38,39	0
2	GOL	B	5006	6/6	0.88	0.09	50,51,54,58	0
2	GOL	A	5002	6/6	0.89	0.14	27,31,39,41	0
2	GOL	A	5001	6/6	0.91	0.10	29,37,38,41	0
2	GOL	A	5004	6/6	0.92	0.08	37,39,40,44	0
2	GOL	A	5003	6/6	0.93	0.10	30,37,44,48	0
2	GOL	B	5005	6/6	0.94	0.08	34,40,41,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.