



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 6, 2024 – 05:55 PM JST

PDB ID : 5AWC  
Title : Crystal structure of human TLR8 in complex with MB-564  
Authors : Tanji, H.; Ohto, U.; Shimizu, T.  
Deposited on : 2015-07-03  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.












Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	811	<div><div>79%12%8%</div></div>
1	B	811	<div><div>78%12%9%</div></div>
1	C	811	<div><div>77%12%10%</div></div>
1	D	811	<div><div>66%20%11%</div></div>
2	E	3	<div><div>33%67%</div></div>
2	G	3	<div><div>100%</div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	3	 33%67%
2	J	3	 100%
2	K	3	 100%
2	M	3	 100%
2	N	3	 67%33%
2	P	3	 100%
3	F	2	 100%
3	I	2	 100%
3	L	2	 100%
3	O	2	 50%50%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 24478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	749	Total	C	N	O	S	0	0	0
			6035	3862	1023	1131	19			
1	B	742	Total	C	N	O	S	0	0	0
			5979	3828	1015	1117	19			
1	C	727	Total	C	N	O	S	0	0	0
			5871	3760	998	1094	19			
1	D	724	Total	C	N	O	S	0	0	0
			5836	3735	990	1092	19			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	ARG	-	expression tag	UNP Q9NR97
A	24	SER	-	expression tag	UNP Q9NR97
A	25	PRO	-	expression tag	UNP Q9NR97
A	26	TRP	-	expression tag	UNP Q9NR97
A	828	GLU	-	expression tag	UNP Q9NR97
A	829	PHE	-	expression tag	UNP Q9NR97
A	830	LEU	-	expression tag	UNP Q9NR97
A	831	VAL	-	expression tag	UNP Q9NR97
A	832	PRO	-	expression tag	UNP Q9NR97
A	833	ARG	-	expression tag	UNP Q9NR97
B	23	ARG	-	expression tag	UNP Q9NR97
B	24	SER	-	expression tag	UNP Q9NR97
B	25	PRO	-	expression tag	UNP Q9NR97
B	26	TRP	-	expression tag	UNP Q9NR97
B	828	GLU	-	expression tag	UNP Q9NR97
B	829	PHE	-	expression tag	UNP Q9NR97
B	830	LEU	-	expression tag	UNP Q9NR97
B	831	VAL	-	expression tag	UNP Q9NR97
B	832	PRO	-	expression tag	UNP Q9NR97
B	833	ARG	-	expression tag	UNP Q9NR97
C	23	ARG	-	expression tag	UNP Q9NR97

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	24	SER	-	expression tag	UNP Q9NR97
C	25	PRO	-	expression tag	UNP Q9NR97
C	26	TRP	-	expression tag	UNP Q9NR97
C	828	GLU	-	expression tag	UNP Q9NR97
C	829	PHE	-	expression tag	UNP Q9NR97
C	830	LEU	-	expression tag	UNP Q9NR97
C	831	VAL	-	expression tag	UNP Q9NR97
C	832	PRO	-	expression tag	UNP Q9NR97
C	833	ARG	-	expression tag	UNP Q9NR97
D	23	ARG	-	expression tag	UNP Q9NR97
D	24	SER	-	expression tag	UNP Q9NR97
D	25	PRO	-	expression tag	UNP Q9NR97
D	26	TRP	-	expression tag	UNP Q9NR97
D	828	GLU	-	expression tag	UNP Q9NR97
D	829	PHE	-	expression tag	UNP Q9NR97
D	830	LEU	-	expression tag	UNP Q9NR97
D	831	VAL	-	expression tag	UNP Q9NR97
D	832	PRO	-	expression tag	UNP Q9NR97
D	833	ARG	-	expression tag	UNP Q9NR97

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	G	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	H	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	J	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	K	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	M	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	N	3	Total	C	N	O	0	0	0
			39	22	2	15			

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	P	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	L	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	O	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is 5-(4-azanylbutyl)-3-pentyl-quinolin-2-amine (three-letter code: M4D) (formula: C<sub>18</sub>H<sub>27</sub>N<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			21	18	3		
5	B	1	Total	C	N	0	0
			21	18	3		
5	C	1	Total	C	N	0	0
			21	18	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	N	0	0
			21	18	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	38	Total	O	0	0
			38	38		
6	B	42	Total	O	0	0
			42	42		
6	C	21	Total	O	0	0
			21	21		
6	D	22	Total	O	0	0
			22	22		




### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Toll-like receptor 8





#### • Molecule 1: Toll-like receptor 8







• Molecule 1: Toll-like receptor 8



• Molecule 1: Toll-like receptor 8





- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 33% 67%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 33% 67%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 100%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 100%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 100%

MAG1  
MAG2  
BMA3

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  67% 33%

MAG1  
MAG2  
BMA3

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P:  100%

MAG1  
MAG2  
BMA3

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  100%

MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:  50% 50%

MAG1  
MAG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.40Å 140.18Å 169.60Å 90.00° 90.47° 90.00°	Depositor
Resolution (Å)	47.48 – 2.50 47.48 – 2.50	Depositor EDS
% Data completeness (in resolution range)	94.7 (47.48-2.50) 94.9 (47.48-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.209 , 0.269 0.212 , 0.267	Depositor DCC
$R_{free}$ test set	6664 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.7	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 28.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.034 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	24478	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, M4D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.62	1/6159 (0.0%)	0.77	5/8351 (0.1%)
1	B	0.57	0/6102	0.74	0/8272
1	C	0.53	0/5992	0.71	2/8122 (0.0%)
1	D	0.56	0/5956	0.73	4/8076 (0.0%)
All	All	0.57	1/24209 (0.0%)	0.74	11/32821 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	D	0	4
All	All	0	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269	PRO	N-CD	5.08	1.54	1.47

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	217	PRO	C-N-CD	6.40	141.84	128.40
1	A	270	CYS	CA-CB-SG	6.22	125.20	114.00
1	D	216	VAL	C-N-CD	6.14	141.29	128.40
1	C	338	ARG	NE-CZ-NH1	-5.99	117.30	120.30
1	A	225	ARG	NE-CZ-NH1	5.89	123.25	120.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	779	ASP	Peptide
1	C	242	PHE	Peptide
1	D	161	ILE	Peptide
1	D	201	THR	Peptide
1	D	803	CYS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6035	0	6028	62	1
1	B	5979	0	5969	57	1
1	C	5871	0	5857	37	0
1	D	5836	0	5817	148	0
2	E	39	0	34	2	0
2	G	39	0	34	0	0
2	H	39	0	34	0	0
2	J	39	0	34	0	0
2	K	39	0	34	0	0
2	M	39	0	34	0	0
2	N	39	0	34	1	0
2	P	39	0	34	0	0
3	F	28	0	25	0	0
3	I	28	0	25	0	0
3	L	28	0	25	0	0
3	O	28	0	25	0	0
4	A	42	0	39	0	0
4	B	28	0	26	0	0
4	C	28	0	26	0	0
4	D	28	0	26	0	0
5	A	21	0	0	1	0
5	B	21	0	0	1	0
5	C	21	0	0	1	0
5	D	21	0	0	0	0
6	A	38	0	0	2	0
6	B	42	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	21	0	0	0	0
6	D	22	0	0	0	0
All	All	24478	0	24160	302	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 302 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:197:PHE:O	1:D:200:LEU:CD2	1.64	1.43
1:D:197:PHE:O	1:D:200:LEU:HD22	1.15	1.15
1:D:181:CYS:SG	1:D:187:CYS:CB	2.36	1.14
1:D:162:THR:HG1	1:D:164:GLU:HG2	1.01	1.09
1:D:177:LEU:O	1:D:208:LEU:HA	1.52	1.09

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:709:ASP:OD1	1:B:338:ARG:NH2[2_546]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	741/811 (91%)	691 (93%)	47 (6%)	3 (0%)	30 49
1	B	732/811 (90%)	675 (92%)	55 (8%)	2 (0%)	37 56
1	C	715/811 (88%)	655 (92%)	58 (8%)	2 (0%)	37 56
1	D	712/811 (88%)	612 (86%)	93 (13%)	7 (1%)	13 25

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2900/3244 (89%)	2633 (91%)	253 (9%)	14 (0%)	25	44

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	161	ILE
1	D	190	THR
1	D	284	GLN
1	B	735	VAL
1	D	246	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	696/755 (92%)	659 (95%)	37 (5%)	19	38
1	B	689/755 (91%)	658 (96%)	31 (4%)	23	46
1	C	676/755 (90%)	642 (95%)	34 (5%)	20	41
1	D	673/755 (89%)	622 (92%)	51 (8%)	11	22
All	All	2734/3020 (90%)	2581 (94%)	153 (6%)	17	36

5 of 153 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	162	THR
1	D	735	VAL
1	D	194	ASP
1	D	296	SER
1	D	779	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	416	ASN
1	D	139	GLN
1	C	686	GLN
1	D	202	ASN
1	B	388	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

32 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	E	1	1,2	14,14,15	0.94	1 (7%)	17,19,21	1.17	1 (5%)
2	NAG	E	2	2	14,14,15	0.89	0	17,19,21	1.41	3 (17%)
2	BMA	E	3	2	11,11,12	0.87	0	15,15,17	1.70	4 (26%)
3	NAG	F	1	1,3	14,14,15	1.01	1 (7%)	17,19,21	1.69	3 (17%)
3	NAG	F	2	3	14,14,15	0.71	0	17,19,21	1.68	2 (11%)
2	NAG	G	1	1,2	14,14,15	0.68	0	17,19,21	1.75	4 (23%)
2	NAG	G	2	2	14,14,15	0.89	1 (7%)	17,19,21	1.31	3 (17%)
2	BMA	G	3	2	11,11,12	0.63	0	15,15,17	1.37	1 (6%)
2	NAG	H	1	1,2	14,14,15	0.75	0	17,19,21	1.52	3 (17%)
2	NAG	H	2	2	14,14,15	0.66	0	17,19,21	1.14	0
2	BMA	H	3	2	11,11,12	0.88	0	15,15,17	1.58	3 (20%)
3	NAG	I	1	1,3	14,14,15	0.95	1 (7%)	17,19,21	2.01	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	I	2	3	14,14,15	0.42	0	17,19,21	1.32	3 (17%)
2	NAG	J	1	1,2	14,14,15	0.65	0	17,19,21	1.35	3 (17%)
2	NAG	J	2	2	14,14,15	0.77	0	17,19,21	1.28	1 (5%)
2	BMA	J	3	2	11,11,12	0.64	0	15,15,17	1.59	3 (20%)
2	NAG	K	1	1,2	14,14,15	0.98	1 (7%)	17,19,21	1.68	3 (17%)
2	NAG	K	2	2	14,14,15	0.70	0	17,19,21	1.45	3 (17%)
2	BMA	K	3	2	11,11,12	0.95	1 (9%)	15,15,17	1.64	4 (26%)
3	NAG	L	1	1,3	14,14,15	0.91	0	17,19,21	1.44	2 (11%)
3	NAG	L	2	3	14,14,15	0.69	0	17,19,21	1.80	6 (35%)
2	NAG	M	1	1,2	14,14,15	0.73	1 (7%)	17,19,21	1.50	5 (29%)
2	NAG	M	2	2	14,14,15	1.14	2 (14%)	17,19,21	1.60	4 (23%)
2	BMA	M	3	2	11,11,12	0.61	0	15,15,17	1.37	3 (20%)
2	NAG	N	1	1,2	14,14,15	0.59	0	17,19,21	1.43	2 (11%)
2	NAG	N	2	2	14,14,15	0.91	1 (7%)	17,19,21	1.38	2 (11%)
2	BMA	N	3	2	11,11,12	0.73	0	15,15,17	1.46	3 (20%)
3	NAG	O	1	1,3	14,14,15	0.81	1 (7%)	17,19,21	1.17	1 (5%)
3	NAG	O	2	3	14,14,15	0.77	0	17,19,21	0.82	0
2	NAG	P	1	1,2	14,14,15	0.62	0	17,19,21	1.55	2 (11%)
2	NAG	P	2	2	14,14,15	0.95	1 (7%)	17,19,21	1.20	3 (17%)
2	BMA	P	3	2	11,11,12	0.91	0	15,15,17	1.94	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	BMA	E	3	2	-	0/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	BMA	G	3	2	-	2/2/19/22	0/1/1/1
2	NAG	H	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
2	BMA	H	3	2	-	2/2/19/22	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1
2	NAG	J	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	J	2	2	-	0/6/23/26	0/1/1/1
2	BMA	J	3	2	-	1/2/19/22	0/1/1/1
2	NAG	K	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	K	2	2	-	0/6/23/26	0/1/1/1
2	BMA	K	3	2	-	0/2/19/22	0/1/1/1
3	NAG	L	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	0/6/23/26	0/1/1/1
2	NAG	M	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	M	2	2	-	0/6/23/26	0/1/1/1
2	BMA	M	3	2	-	0/2/19/22	0/1/1/1
2	NAG	N	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	N	2	2	-	2/6/23/26	0/1/1/1
2	BMA	N	3	2	-	0/2/19/22	0/1/1/1
3	NAG	O	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	O	2	3	-	0/6/23/26	0/1/1/1
2	NAG	P	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	P	2	2	-	0/6/23/26	0/1/1/1
2	BMA	P	3	2	-	0/2/19/22	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	K	1	NAG	O5-C1	-3.31	1.38	1.43
2	M	2	NAG	O5-C1	-3.05	1.38	1.43
2	E	1	NAG	O5-C1	-2.38	1.39	1.43
3	O	1	NAG	O5-C1	-2.35	1.40	1.43
2	N	2	NAG	O5-C1	-2.25	1.40	1.43

The worst 5 of 86 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	1	NAG	C1-C2-N2	-5.43	101.21	110.49
3	F	2	NAG	C1-O5-C5	5.13	119.15	112.19
2	K	1	NAG	O5-C1-C2	-4.75	103.78	111.29
2	P	3	BMA	C1-O5-C5	4.37	118.11	112.19
3	F	1	NAG	C1-C2-N2	-4.22	103.27	110.49

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	N	2	NAG	C4-C5-C6-O6
2	N	2	NAG	O5-C5-C6-O6
2	N	1	NAG	C4-C5-C6-O6
2	G	3	BMA	C4-C5-C6-O6
2	G	3	BMA	O5-C5-C6-O6



There are no ring outliers.



3 monomers are involved in 3 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2	NAG	1	0
2	N	1	NAG	1	0
2	E	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


























## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	M4D	B	901	-	22,22,22	1.41	2 (9%)	25,28,28	1.08	2 (8%)
4	NAG	B	907	1	14,14,15	0.77	0	17,19,21	1.35	3 (17%)
4	NAG	A	1010	1	14,14,15	0.56	0	17,19,21	1.39	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	M4D	A	1012	-	22,22,22	1.58	3 (13%)	25,28,28	0.98	2 (8%)
4	NAG	D	907	1	14,14,15	0.83	0	17,19,21	1.85	3 (17%)
4	NAG	D	911	1	14,14,15	0.53	0	17,19,21	0.93	0
4	NAG	C	1006	1	14,14,15	0.58	0	17,19,21	1.70	3 (17%)
4	NAG	A	1006	1	14,14,15	0.62	0	17,19,21	1.35	3 (17%)
4	NAG	B	911	1	14,14,15	0.62	0	17,19,21	1.23	2 (11%)
4	NAG	A	1011	1	14,14,15	0.81	1 (7%)	17,19,21	0.89	1 (5%)
4	NAG	C	1010	1	14,14,15	0.66	0	17,19,21	1.47	2 (11%)
5	M4D	D	901	-	22,22,22	1.13	4 (18%)	25,28,28	1.00	1 (4%)
5	M4D	C	1011	-	22,22,22	1.46	3 (13%)	25,28,28	0.96	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	M4D	B	901	-	-	0/10/10/10	0/2/2/2
4	NAG	B	907	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1010	1	-	2/6/23/26	0/1/1/1
5	M4D	A	1012	-	-	2/10/10/10	0/2/2/2
4	NAG	D	907	1	-	2/6/23/26	0/1/1/1
4	NAG	D	911	1	-	2/6/23/26	0/1/1/1
4	NAG	C	1006	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1006	1	-	0/6/23/26	0/1/1/1
4	NAG	B	911	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1011	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1010	1	-	0/6/23/26	0/1/1/1
5	M4D	D	901	-	-	0/10/10/10	0/2/2/2
5	M4D	C	1011	-	-	2/10/10/10	0/2/2/2

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1012	M4D	C7-C6	4.49	1.50	1.42
5	C	1011	M4D	C7-C6	4.19	1.50	1.42
5	B	901	M4D	C7-C6	4.16	1.50	1.42
5	A	1012	M4D	C1-C	3.96	1.49	1.42
5	C	1011	M4D	C1-C	3.46	1.48	1.42

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	907	NAG	O5-C1-C2	-5.46	102.66	111.29
4	B	911	NAG	C1-O5-C5	3.78	117.31	112.19
4	C	1006	NAG	C8-C7-N2	3.76	122.47	116.10
4	A	1010	NAG	O5-C5-C6	3.60	112.85	107.20
4	C	1010	NAG	C1-O5-C5	3.46	116.88	112.19

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	911	NAG	O5-C5-C6-O6
4	A	1010	NAG	O5-C5-C6-O6
4	D	907	NAG	O5-C5-C6-O6
4	A	1010	NAG	C4-C5-C6-O6
4	D	911	NAG	C4-C5-C6-O6

There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	901	M4D	1	0
5	A	1012	M4D	1	0
5	C	1011	M4D	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	749/811 (92%)	-0.34	8 (1%) 77 74	31, 46, 82, 114	0
1	B	742/811 (91%)	-0.25	12 (1%) 70 67	30, 50, 88, 118	0
1	C	727/811 (89%)	0.01	17 (2%) 61 58	34, 58, 106, 133	0
1	D	724/811 (89%)	0.41	55 (7%) 21 20	34, 72, 121, 150	0
All	All	2942/3244 (90%)	-0.05	92 (3%) 51 48	30, 53, 108, 150	0

The worst 5 of 92 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	100	VAL	5.5
1	D	100	VAL	5.0
1	D	815	VAL	4.8
1	D	732	LEU	4.5
1	D	58	PRO	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BMA	N	3	11/12	0.70	0.14	72,79,85,90	0
2	BMA	H	3	11/12	0.81	0.13	44,55,60,61	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BMA	E	3	11/12	0.81	0.10	44,48,54,54	0
2	BMA	K	3	11/12	0.83	0.11	44,54,62,63	0
2	BMA	G	3	11/12	0.83	0.10	52,56,63,70	0
2	BMA	P	3	11/12	0.83	0.13	53,60,77,78	0
2	BMA	M	3	11/12	0.85	0.09	59,66,77,81	0
3	NAG	L	2	14/15	0.86	0.09	47,55,62,65	0
2	BMA	J	3	11/12	0.89	0.08	50,55,60,60	0
3	NAG	I	2	14/15	0.91	0.08	53,60,65,66	0
3	NAG	O	2	14/15	0.91	0.09	58,62,67,70	0
3	NAG	F	2	14/15	0.92	0.08	43,51,57,60	0
2	NAG	N	2	14/15	0.92	0.09	58,66,75,82	0
2	NAG	P	2	14/15	0.93	0.07	37,47,51,55	0
2	NAG	N	1	14/15	0.94	0.08	55,61,69,69	0
2	NAG	K	2	14/15	0.95	0.06	39,45,49,52	0
2	NAG	G	2	14/15	0.95	0.08	39,44,51,53	0
3	NAG	O	1	14/15	0.95	0.07	38,42,44,51	0
2	NAG	K	1	14/15	0.95	0.08	38,41,44,45	0
3	NAG	F	1	14/15	0.96	0.06	32,34,42,42	0
2	NAG	G	1	14/15	0.96	0.05	34,37,38,41	0
2	NAG	H	1	14/15	0.96	0.07	35,40,44,46	0
2	NAG	M	1	14/15	0.96	0.06	45,46,50,50	0
2	NAG	M	2	14/15	0.96	0.07	44,49,53,54	0
2	NAG	H	2	14/15	0.96	0.06	36,45,54,58	0
2	NAG	E	1	14/15	0.97	0.06	30,33,39,50	0
3	NAG	I	1	14/15	0.97	0.06	34,37,41,47	0
2	NAG	P	1	14/15	0.97	0.05	37,44,50,55	0
3	NAG	L	1	14/15	0.97	0.06	38,41,45,48	0
2	NAG	J	1	14/15	0.97	0.05	32,38,40,41	0
2	NAG	J	2	14/15	0.97	0.05	37,39,47,51	0
2	NAG	E	2	14/15	0.97	0.06	30,34,42,44	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain E:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain G:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain J:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain K:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain M:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain N:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain P:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain F:**




$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain I:**


$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain L:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	C	1006	14/15	0.80	0.14	74,83,107,117	0
4	NAG	D	907	14/15	0.88	0.11	56,69,82,87	0
4	NAG	C	1010	14/15	0.89	0.09	78,82,87,88	0
4	NAG	B	907	14/15	0.89	0.12	55,62,70,72	0
4	NAG	B	911	14/15	0.91	0.07	56,59,67,70	0
4	NAG	A	1011	14/15	0.91	0.08	59,64,69,69	0
4	NAG	D	911	14/15	0.91	0.10	58,67,74,74	0
5	M4D	C	1011	21/21	0.91	0.13	48,52,61,63	0
4	NAG	A	1006	14/15	0.92	0.09	56,64,71,73	0
4	NAG	A	1010	14/15	0.92	0.09	60,70,81,85	0
5	M4D	B	901	21/21	0.95	0.09	39,42,46,47	0
5	M4D	A	1012	21/21	0.96	0.07	32,37,47,49	0
5	M4D	D	901	21/21	0.96	0.08	35,37,41,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around M4D C 1011:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around M4D B 901:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around M4D A 1012:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.