



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 1, 2025 – 10:49 pm BST

PDB ID : 2BIG / pdb\_00002big  
Title : Radiation damage of the Schiff base in phosphoserine aminotransferase (structure I)  
Authors : Dubnovitsky, A.P.; Ravelli, R.B.G.; Popov, A.N.; Papageorgiou, A.C.  
Deposited on : 2005-01-21  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1387 (1.30-1.30)
Clashscore	180529	1497 (1.30-1.30)
Ramachandran outliers	177936	1455 (1.30-1.30)
Sidechain outliers	177891	1455 (1.30-1.30)
RSRZ outliers	164620	1384 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>..</div> </div> </div>
1	B	361	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOSERINE AMINOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	358	Total	C	N	O	S	0	17	0
			2864	1814	490	545	15			
1	B	355	Total	C	N	O	S	0	13	0
			2831	1797	479	541	14			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula:  $C_8H_{10}NO_6P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Mg 2 2	0	0
3	B	3	Total Mg 4 4	0	1

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0
4	B	2	Total Cl 2 2	0	0

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 10 6 4	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	436	Total	O	0	0
			436	436		
7	B	440	Total	O	0	0
			440	440		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHOSPHOSERINE AMINOTRANSFERASE



- Molecule 1: PHOSPHOSERINE AMINOTRANSFERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.80Å 84.30Å 67.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.00 – 1.30 12.00 – 1.30	Depositor EDS
% Data completeness (in resolution range)	93.7 (12.00-1.30) 93.6 (12.00-1.30)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 1.30Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.117 , 0.156 0.128 , 0.137	Depositor DCC
$R_{free}$ test set	9397 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.6	Xtriage
Anisotropy	0.351	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.46 , 76.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6635	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PEG, MG, CL, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/2998	1.17	20/4051 (0.5%)
1	B	0.66	0/2941	1.20	20/3978 (0.5%)
All	All	0.66	0/5939	1.19	40/8029 (0.5%)

There are no bond length outliers.

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	42	ARG	NE-CZ-NH1	-11.54	114.53	120.30
1	B	248	ARG	NE-CZ-NH2	-9.49	115.56	120.30
1	B	248	ARG	NE-CZ-NH1	9.34	124.97	120.30
1	B	344	ASP	CB-CG-OD2	-9.02	110.18	118.30
1	B	42	ARG	NE-CZ-NH1	-8.94	115.83	120.30
1	B	233	ASP	CB-CG-OD2	-8.41	110.73	118.30
1	B	328	ARG	NE-CZ-NH1	-8.35	116.13	120.30
1	A	350[A]	ARG	NE-CZ-NH2	-8.16	116.22	120.30
1	A	350[B]	ARG	NE-CZ-NH2	-8.16	116.22	120.30
1	B	42	ARG	NE-CZ-NH2	7.90	124.25	120.30
1	B	132	ASP	CB-CG-OD2	7.81	125.33	118.30
1	A	328	ARG	NE-CZ-NH1	7.55	124.08	120.30
1	A	248	ARG	NE-CZ-NH2	-7.34	116.63	120.30
1	A	304	ARG	NE-CZ-NH2	-6.99	116.80	120.30
1	B	46	TYR	CB-CG-CD1	-6.89	116.87	121.00
1	A	44	GLN	CB-CG-CD	6.88	129.50	111.60
1	A	42	ARG	NH1-CZ-NH2	6.61	126.67	119.40
1	B	58	ARG	NE-CZ-NH1	6.51	123.55	120.30
1	A	218	GLU	O-C-N	6.31	132.79	122.70
1	B	42	ARG	CD-NE-CZ	5.85	131.79	123.60
1	B	257	LEU	CA-CB-CG	5.85	128.75	115.30
1	A	51	GLU	OE1-CD-OE2	-5.82	116.32	123.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307	GLU	OE1-CD-OE2	-5.82	116.32	123.30
1	A	50[A]	HIS	CA-CB-CG	5.81	123.47	113.60
1	A	50[B]	HIS	CA-CB-CG	5.81	123.47	113.60
1	B	338	TYR	CB-CG-CD2	5.75	124.45	121.00
1	B	328	ARG	NE-CZ-NH2	5.69	123.15	120.30
1	B	211	ASP	CB-CG-OD2	-5.59	113.27	118.30
1	B	302[A]	ASN	CA-CB-CG	5.51	125.53	113.40
1	B	302[B]	ASN	CA-CB-CG	5.51	125.53	113.40
1	A	66	ASP	CB-CG-OD2	5.46	123.22	118.30
1	A	328	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	164	ILE	CA-CB-CG2	5.45	121.80	110.90
1	B	46	TYR	CA-CB-CG	-5.38	103.17	113.40
1	B	83	LEU	CB-CG-CD1	-5.30	101.99	111.00
1	A	231	LYS	CD-CE-NZ	5.29	123.87	111.70
1	A	294	ARG	NE-CZ-NH1	-5.16	117.72	120.30
1	B	163	GLU	OE1-CD-OE2	-5.11	117.17	123.30
1	A	136	PHE	CB-CG-CD2	-5.03	117.28	120.80
1	A	51	GLU	CG-CD-OE1	5.02	128.35	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2864	0	2817	17	0
1	B	2831	0	2795	18	0
2	A	15	0	6	0	0
2	B	15	0	7	1	0
3	A	2	0	0	0	0
3	B	4	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	10	0	14	0	0
6	A	7	0	10	0	0
6	B	7	0	10	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	436	0	0	11	0
7	B	440	0	0	9	0
All	All	6635	0	5659	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266[B]:GLN:NE2	7:A:2309:HOH:O	2.02	0.91
1:A:266[B]:GLN:CD	7:A:2309:HOH:O	2.09	0.91
1:B:302[A]:ASN:HD22	1:B:309:ASN:HD21	1.25	0.81
1:B:328:ARG:HD2	7:B:2406:HOH:O	1.86	0.75
1:A:115:GLU:HG3	7:A:2170:HOH:O	1.87	0.74
1:B:302[A]:ASN:ND2	1:B:309:ASN:HD21	1.86	0.74
1:A:307:GLU:HA	1:A:310:GLN:OE1	1.92	0.69
1:B:200:PRO:HD2	7:B:2298:HOH:O	1.95	0.65
1:A:14[B]:LEU:HD23	7:A:2026:HOH:O	2.00	0.61
1:B:16:LYS:HE3	7:B:2036:HOH:O	2.04	0.58
1:A:200:PRO:HG3	1:A:243:SER:O	2.04	0.57
1:A:328:ARG:O	1:A:328:ARG:HG2	2.03	0.57
1:A:105:LYS:HE3	7:A:2145:HOH:O	2.05	0.57
1:A:266[B]:GLN:HG3	7:A:2309:HOH:O	2.06	0.56
1:A:266[B]:GLN:CG	7:A:2309:HOH:O	2.49	0.56
1:B:288[B]:HIS:HD2	7:B:2230:HOH:O	1.90	0.55
1:B:288[A]:HIS:HD2	7:B:2363:HOH:O	1.90	0.55
1:B:218:GLU:O	1:B:219:GLN:HB2	2.05	0.54
1:A:196:LYS:HG2	7:A:2245:HOH:O	2.07	0.54
1:A:14[B]:LEU:CD2	1:A:199:GLY:HA2	2.37	0.54
1:B:352:LEU:HD12	7:B:2431:HOH:O	2.07	0.54
1:A:3:GLN:HG2	1:A:5:PHE:CZ	2.43	0.53
1:A:196:LYS:HD3	7:A:2245:HOH:O	2.08	0.52
1:B:302[A]:ASN:HD22	1:B:309:ASN:ND2	2.02	0.51
1:B:196:LYS:NZ	2:B:1361:PLP:O3	2.48	0.46
1:B:266[A]:GLN:NE2	7:B:2324:HOH:O	2.48	0.46
1:B:266[A]:GLN:NE2	7:B:2327:HOH:O	2.48	0.46
1:A:123[A]:LYS:NZ	7:A:2184:HOH:O	2.49	0.45
1:B:67:TYR:OH	1:B:209:LYS:HE2	2.17	0.45
1:B:3:GLN:NE2	7:B:2001:HOH:O	2.49	0.45
1:A:315:LYS:NZ	7:A:2376:HOH:O	2.49	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:VAL:HG12	1:A:208:VAL:HG13	2.00	0.43
1:B:195:GLN:HA	1:B:199:GLY:O	2.19	0.42
1:B:288[B]:HIS:CD2	1:B:300:THR:HB	2.56	0.41
1:B:169:LEU:C	1:B:170[B]:ILE:HD13	2.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/361 (103%)	367 (99%)	5 (1%)	0	100	100
1	B	364/361 (101%)	357 (98%)	7 (2%)	0	100	100
All	All	736/722 (102%)	724 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/310 (103%)	317 (99%)	3 (1%)	75	50
1	B	316/310 (102%)	313 (99%)	3 (1%)	75	50
All	All	636/620 (103%)	630 (99%)	6 (1%)	73	50

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	196	LYS
1	A	328	ARG
1	B	90	LYS
1	B	210	LYS
1	B	328	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	30	ASN
1	A	44	GLN
1	A	68	GLN
1	A	319	GLN
1	B	68	GLN
1	B	282	ASN
1	B	319	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 10 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	PEG	B	1367	-	6,6,6	0.56	0	5,5,5	1.13	0
6	PEG	A	1367	-	6,6,6	0.50	0	5,5,5	1.31	1 (20%)
2	PLP	A	1361	1	15,15,16	1.14	1 (6%)	20,22,23	1.45	1 (5%)
5	PGE	A	1366	-	9,9,9	0.76	0	8,8,8	1.75	2 (25%)
2	PLP	B	1361	1	15,15,16	1.51	2 (13%)	20,22,23	1.32	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	B	1367	-	-	0/4/4/4	-
6	PEG	A	1367	-	-	0/4/4/4	-
2	PLP	A	1361	1	-	0/6/6/8	0/1/1/1
5	PGE	A	1366	-	-	3/7/7/7	-
2	PLP	B	1361	1	-	0/6/6/8	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1361	PLP	C3-C2	-3.26	1.37	1.40
2	A	1361	PLP	C3-C2	-2.48	1.38	1.40
2	B	1361	PLP	C5A-C5	2.04	1.56	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1361	PLP	C4A-C4-C5	-4.25	116.55	120.94
2	B	1361	PLP	C6-C5-C4	3.40	120.83	118.16
5	A	1366	PGE	C3-O2-C2	3.05	126.50	113.29
5	A	1366	PGE	C5-O3-C4	-2.89	100.75	113.29
6	A	1367	PEG	C3-O2-C2	2.67	124.85	113.29
2	B	1361	PLP	C5-C6-N1	-2.37	119.86	123.82
2	B	1361	PLP	C2A-C2-C3	2.04	123.41	120.89

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1366	PGE	O2-C3-C4-O3
5	A	1366	PGE	O3-C5-C6-O4
5	A	1366	PGE	C4-C3-O2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1361	PLP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	358/361 (99%)	-0.46	3 (0%) 82 85	7, 13, 30, 47	17 (4%)
1	B	355/361 (98%)	-0.41	3 (0%) 82 85	8, 13, 31, 54	13 (3%)
All	All	713/722 (98%)	-0.44	6 (0%) 82 85	7, 13, 30, 54	30 (4%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	213	LEU	4.0
1	B	217	VAL	3.7
1	B	218	GLU	3.0
1	A	257[B]	LEU	3.0
1	A	200	PRO	3.0
1	A	216	LYS	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	PGE	A	1366	10/10	0.89	0.10	40,43,48,52	0
6	PEG	A	1367	7/7	0.94	0.08	23,24,34,44	0
6	PEG	B	1367	7/7	0.94	0.08	21,25,34,35	0
3	MG	B	1363	1/1	0.97	0.09	29,29,29,29	1
3	MG	B	1364[A]	1/1	0.97	0.11	25,25,25,25	1
3	MG	B	1364[B]	1/1	0.97	0.11	23,23,23,23	1
2	PLP	A	1361	15/16	0.99	0.03	8,10,12,19	0
2	PLP	B	1361	15/16	0.99	0.03	9,10,14,20	0
3	MG	B	1362	1/1	0.99	0.03	11,11,11,11	1
4	CL	A	1365	1/1	1.00	0.05	17,17,17,17	0
4	CL	B	1365	1/1	1.00	0.01	10,10,10,10	0
4	CL	B	1366	1/1	1.00	0.07	19,19,19,19	0
3	MG	A	1362	1/1	1.00	0.08	16,16,16,16	0
3	MG	A	1363	1/1	1.00	0.07	15,15,15,15	0
4	CL	A	1364	1/1	1.00	0.01	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.