



wwPDB EM Validation Summary Report ⓘ

Oct 6, 2024 – 07:32 pm BST

PDB ID : 8C8M
EMDB ID : EMD-16484
Title : In vitro structure of the Nitrosopumilus maritimus S-layer - Composite map between two and six-fold symmetrised
Authors : von Kuegelgen, A.; Bharat, T.
Deposited on : 2023-01-20
Resolution : 2.87 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1734	
1	B	1734	
1	C	1734	
1	D	1734	
1	E	1734	
1	F	1734	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 70506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell surface protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1580	Total	C	N	O	S	0	0
			11751	7234	1888	2610	19		
1	B	1580	Total	C	N	O	S	0	0
			11751	7234	1888	2610	19		
1	C	1580	Total	C	N	O	S	0	0
			11751	7234	1888	2610	19		
1	D	1580	Total	C	N	O	S	0	0
			11751	7234	1888	2610	19		
1	E	1580	Total	C	N	O	S	0	0
			11751	7234	1888	2610	19		
1	F	1580	Total	C	N	O	S	0	0
			11751	7234	1888	2610	19		

SER LEU LEU TRP THR GLY SER SER SER GLN SER PHE SER PRO PRO LEU LEU SER TRP THR ILE PRO THR PRO GLU ALA ALA GLY THR TYR THR THR THR THR PHE VAL VAL TRP TRP GLU GLY VAL ASP ASN PRO PRO THR THR LEU LEU SER SER PRO PRO VAL SER THR THR VAL ASN VAL SER


- Molecule 1: Cell surface protein

Chain B: 73% 18% 9%

[illegible]

- Molecule 1: Cell surface protein

Chain C:  73% 18% 9%



ASN	V1369	
LEU		
SER	D1372	
GLY		
ARG	Y1376	
THR	S1377	
VAL	W1378	
ASP	T1379	
ALA		
PHE	V1382	
PRO	Y1383	
SER	I1384	
ASN	T1385	
LEU		
TRP	F1393	
ILE		
ASP	D1399	
SER		
VAL	E1403	
THR	T1404	
GLY		
ALA	I1409	
PHE	K1410	
PRO	V1411	
SER		
LEU	R1414	
THR		
GLU	K1422	
ASN	L1423	
ASP		
ASN	T1434	
PRO		
THR	I1438	
GLU	L1439	
ALA		
SER	D1446	
LEU		
THR	T1453	
PRO		
VAL	P1465	
THR		
SER	L1469	
ASN		
VAL	D1473	
SER	D1474	
THR		
GLY	V1480	
ASN		
VAL	I1490	
THR	V1491	
VAL		
PRO	N1499	
LEU	I1500	
ALA		
TRP	V1503	
ILE		
THR	E1507	
GLY		
SER	Y1510	
LEU		
ALA		

G1516	
V1517	
V1520	
D1524	
P1529	
V1532	
V1537	
S1541	
L1550	
T1551	
V1552	
T1553	
E1554	
T1555	
A1558	
T1559	
G1560	
I1561	
F1562	
E1563	
G1564	
T1565	
V1566	
F1567	
S1574	
R1578	
I1579	
R1580	
D1585	
M1594	
T1595	
L1596	
P1597	
E1605	
T1618	
VAL	
VAL	
PRO	
PRO	
LEU	
ALA	
TRP	
ILE	
THR	
GLY	
SER	
ALA	

SER
SER
GLY
GLN
SER
PHE
SER
PRO
ALA
LEU
SER
TRP
ILE
PRO
THR
GLU
ALA
GLY
THR
TYR
THR
ALA
THR
ALA
PHE
VAL
TRP
GLU
SER
VAL
ASP
ASN
PRO
THR
ALA
LEU
SER
PHE
PRO
PRO
VAL
THR
THR
THR
VAL
ASN
VAL
SER

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	354860	Depositor
Resolution determination method	OTHER	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; RELION refinement with in-built CTF correction. The function is similar to a Wiener filter, so amplitude correction included.	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48.5	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	10.250	Depositor
Minimum map value	-4.662	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.422	Depositor
Recommended contour level	1.05514	Depositor
Map size (\AA)	349.44, 349.44, 349.44	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.092, 1.092, 1.092	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	9/11941 (0.1%)	0.81	18/16339 (0.1%)
1	B	0.69	2/11941 (0.0%)	0.77	8/16339 (0.0%)
1	C	0.69	2/11941 (0.0%)	0.77	6/16339 (0.0%)
1	D	0.68	1/11941 (0.0%)	0.76	7/16339 (0.0%)
1	E	0.69	2/11941 (0.0%)	0.77	7/16339 (0.0%)
1	F	0.70	3/11941 (0.0%)	0.77	5/16339 (0.0%)
All	All	0.70	19/71646 (0.0%)	0.77	51/98034 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	793	GLU	CD-OE2	16.03	1.43	1.25
1	A	1255	GLU	CD-OE1	-15.38	1.08	1.25
1	A	793	GLU	CD-OE1	-13.35	1.10	1.25
1	A	196	GLU	CD-OE1	-10.86	1.13	1.25
1	A	1299	GLU	CD-OE1	-9.95	1.14	1.25

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	729	ASP	CB-CG-OD1	15.40	132.16	118.30
1	A	123	ASP	CB-CG-OD1	11.60	128.74	118.30
1	A	794	ASP	CB-CG-OD2	10.58	127.82	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	729	ASP	CB-CG-OD2	-9.00	110.20	118.30
1	E	1257	ASP	CB-CG-OD1	-8.15	110.96	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	196	GLU	Sidechain
1	A	729	ASP	Sidechain
1	B	55	SER	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11751	0	10951	245	0
1	B	11751	0	10951	221	0
1	C	11751	0	10951	248	0
1	D	11751	0	10951	252	0
1	E	11751	0	10949	221	0
1	F	11751	0	10951	248	0
All	All	70506	0	65704	1252	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 1252 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1217:SER:CB	1:D:1424:VAL:HG21	1.65	1.25
1:A:1424:VAL:HG21	1:F:1217:SER:CB	1.65	1.24
1:C:816:PRO:HD2	1:D:1083:ASN:HD21	1.01	1.14
1:A:1424:VAL:CG2	1:F:1217:SER:HB2	1.82	1.09
1:C:1217:SER:HB2	1:D:1424:VAL:CG2	1.83	1.08

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1578/1734 (91%)	1517 (96%)	61 (4%)	0	100	100
1	B	1578/1734 (91%)	1518 (96%)	60 (4%)	0	100	100
1	C	1578/1734 (91%)	1520 (96%)	58 (4%)	0	100	100
1	D	1578/1734 (91%)	1518 (96%)	60 (4%)	0	100	100
1	E	1578/1734 (91%)	1518 (96%)	60 (4%)	0	100	100
1	F	1578/1734 (91%)	1520 (96%)	58 (4%)	0	100	100
All	All	9468/10404 (91%)	9111 (96%)	357 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1312/1438 (91%)	1300 (99%)	12 (1%)	75	91
1	B	1312/1438 (91%)	1301 (99%)	11 (1%)	79	92
1	C	1312/1438 (91%)	1302 (99%)	10 (1%)	79	92
1	D	1312/1438 (91%)	1300 (99%)	12 (1%)	75	91
1	E	1312/1438 (91%)	1300 (99%)	12 (1%)	75	91
1	F	1312/1438 (91%)	1300 (99%)	12 (1%)	75	91
All	All	7872/8628 (91%)	7803 (99%)	69 (1%)	74	91

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	38	ASN
1	F	269	ARG
1	F	1047	PHE
1	C	274	ARG
1	C	269	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 75 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	740	ASN
1	F	956	GLN
1	E	853	ASN
1	F	97	ASN
1	B	1154	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-16484. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 156



Y Index: 175




Z Index: 186


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 1.05514. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 697 nm³; this corresponds to an approximate mass of 630 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.348 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-16484 and PDB model 8C8M. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 1.05514 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.05514).

9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 96% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (1.05514) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.9580	<div><div></div></div> 0.5560
A	<div><div></div></div> 0.9620	<div><div></div></div> 0.5630
B	<div><div></div></div> 0.9580	<div><div></div></div> 0.5530
C	<div><div></div></div> 0.9530	<div><div></div></div> 0.5520
D	<div><div></div></div> 0.9630	<div><div></div></div> 0.5620
E	<div><div></div></div> 0.9580	<div><div></div></div> 0.5530
F	<div><div></div></div> 0.9540	<div><div></div></div> 0.5520

1.0

0.0

<0.0