



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 01:07 AM EDT

PDB ID : 5CTM  
Title : Structure of BPu1 beta-lactamase  
Authors : Smith, C.A.; Vakulenko, S.B.  
Deposited on : 2015-07-24  
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1050 (1.06-0.94)
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)
RSRZ outliers	127900	1023 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div>3%</div> <div>94%</div> <div>6%</div> </div>
1	B	233	<div> <div>10%</div> <div>89%</div> <div>10%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FLC	A	301	-	-	X	-
4	PEG	B	302	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9158 atoms, of which 4254 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	233	Total	C	H	N	O	S	28	33	0
			4264	1367	2156	349	380	12			
1	B	232	Total	C	H	N	O	S	11	23	0
			4111	1316	2061	356	368	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	ALA	SER	conflict	UNP A8FFI9
B	59	ALA	SER	conflict	UNP A8FFI9

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula:  $C_6H_5O_7^-$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			18	6	5	7		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			18	6	5	7		
2	B	1	Total	C	H	O	0	0
			18	6	5	7		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	401	Total	O	0	17
			420	420		
5	B	268	Total	O	0	4
			272	272		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.66Å 79.87Å 65.01Å 90.00° 92.29° 90.00°	Depositor
Resolution (Å)	37.70 – 1.00 37.70 – 0.99	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.70-1.00) 99.9 (37.70-0.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 0.99Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.126 , 0.138 0.126 , 0.139	Depositor DCC
$R_{free}$ test set	13540 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.3	Xtriage
Anisotropy	0.476	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 46.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	9158	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, KCX, PEG, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2259	0.67	1/3036 (0.0%)
1	B	0.38	0/2146	0.60	0/2889
All	All	0.38	0/4405	0.64	1/5925 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	118	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	80[B]	VAL	Peptide
1	B	81[B]	LYS	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	2156	2137	14	0
1	B	2050	2061	2040	24	0
2	A	26	10	10	7	0
2	B	13	5	5	1	0
3	A	8	12	12	0	0
4	B	7	10	10	5	0
5	A	420	0	0	3	0
5	B	272	0	0	3	0
All	All	4904	4254	4214	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:135[A]:ASN:ND2	1:B:148:ASP:OD2	1.99	0.95
1:B:79[B]:GLU:O	1:B:81[B]:LYS:N	2.15	0.79
1:A:95[A]:GLU:OE2	5:A:401:HOH:O	2.01	0.76
1:A:163[B]:GLU:OE1	5:A:402:HOH:O	2.05	0.73
1:B:118[B]:ARG:NH1	1:B:119:ASP:OD2	2.22	0.72
1:B:194[A]:ARG:NH1	5:B:402:HOH:O	2.25	0.69
1:B:80[B]:VAL:HG23	1:B:205:LYS:HG3	1.76	0.68
1:A:149[B]:SER:HB3	2:A:301:FLC:HG2	1.76	0.66
1:B:284:TYR:HA	4:B:302:PEG:H21	1.77	0.66
1:A:163[A]:GLU:OE1	5:A:403:HOH:O	2.12	0.66
1:A:149[B]:SER:CB	2:A:301:FLC:HG2	2.27	0.63
1:A:149[A]:SER:HB3	2:A:301:FLC:HG2	1.85	0.58
1:B:275[B]:ASP:OD1	5:B:401:HOH:O	2.17	0.57
1:A:149[A]:SER:CB	2:A:301:FLC:HG2	2.35	0.56
1:B:149:SER:OG	2:B:301:FLC:OG1	2.16	0.56
1:B:80[B]:VAL:HG23	1:B:205:LYS:CG	2.38	0.53
1:B:79[B]:GLU:C	1:B:81[B]:LYS:N	2.62	0.53
1:B:283:LYS:HG3	4:B:302:PEG:H42	1.91	0.53
1:B:283:LYS:CG	4:B:302:PEG:H42	2.39	0.53
1:B:83[B]:LYS:O	1:B:84[B]:SER:OG	2.23	0.52
1:B:79[B]:GLU:H	1:B:82[B]:GLU:HG2	1.74	0.52
1:A:149[B]:SER:HB3	2:A:301:FLC:CG	2.39	0.52
1:A:159[B]:ARG:NE	1:A:185:ASP:OD2	2.43	0.52

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:GLN:HG2	1:B:79[B]:GLU:O	2.09	0.52
1:B:266[A]:ILE:HD11	1:B:272:GLU:HB2	1.91	0.51
1:B:79[B]:GLU:HB2	1:B:82[B]:GLU:N	2.29	0.47
1:B:275[A]:ASP:OD2	1:B:279[A]:ARG:NH2	2.47	0.47
1:B:279[A]:ARG:NH1	5:B:405:HOH:O	2.47	0.47
1:B:209[B]:GLU:HG3	1:B:211:THR:HG23	1.95	0.47
1:B:79[B]:GLU:N	1:B:82[B]:GLU:HG2	2.30	0.46
1:A:167[B]:ASN:OD1	1:A:171:LYS:HE2	2.17	0.45
1:A:242[B]:SER:HB2	2:A:301:FLC:HA1	1.99	0.44
1:B:284:TYR:HD1	4:B:302:PEG:H12	1.83	0.43
1:B:284:TYR:CD1	4:B:302:PEG:H12	2.54	0.43
1:A:245:ASP:OD2	1:A:267[B]:LYS:HG2	2.19	0.42
1:B:212:LEU:HB2	1:B:219[A]:MET:SD	2.59	0.42
1:A:285:HIS:HE2	2:A:302:FLC:CAC	2.33	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/233 (113%)	257 (98%)	6 (2%)	0	100	100
1	B	252/233 (108%)	240 (95%)	8 (3%)	4 (2%)	9	1
All	All	515/466 (110%)	497 (96%)	14 (3%)	4 (1%)	34	3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	80[A]	VAL
1	B	80[B]	VAL
1	B	81[A]	LYS
1	B	81[B]	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/201 (116%)	228 (97%)	6 (3%)	46	13
1	B	218/201 (108%)	218 (100%)	0	100	100
All	All	452/402 (112%)	446 (99%)	6 (1%)	81	36

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65[A]	ASP
1	A	65[B]	ASP
1	A	132[A]	GLU
1	A	132[B]	GLU
1	A	149[A]	SER
1	A	149[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KCX	A	104	1	9,11,12	1.24	1 (11%)	5,12,14	1.55	1 (20%)
1	KCX	B	104	1	9,11,12	0.92	1 (11%)	5,12,14	1.30	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	104	1	-	0/9/10/12	-
1	KCX	B	104	1	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	104	KCX	CE-NZ	2.44	1.51	1.46
1	B	104	KCX	CE-NZ	2.22	1.51	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	104	KCX	OQ1-CX-NZ	-3.02	120.27	124.96
1	B	104	KCX	CE-NZ-CX	2.67	126.17	121.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FLC	B	301	-	12,12,12	1.03	0	17,17,17	1.27	1 (5%)
2	FLC	A	302	-	12,12,12	1.03	0	17,17,17	5.27	7 (41%)
4	PEG	B	302	-	6,6,6	0.35	0	5,5,5	0.95	0
3	EDO	A	303	-	3,3,3	0.45	0	2,2,2	0.26	0
3	EDO	A	304	-	3,3,3	0.46	0	2,2,2	0.30	0
2	FLC	A	301	-	12,12,12	1.38	2 (16%)	17,17,17	1.65	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	B	301	-	-	9/16/16/16	-
2	FLC	A	302	-	-	8/16/16/16	-
4	PEG	B	302	-	-	2/4/4/4	-
3	EDO	A	303	-	-	0/1/1/1	-
3	EDO	A	304	-	-	0/1/1/1	-
2	FLC	A	301	-	-	5/16/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	FLC	OA2-CAC	-2.85	1.21	1.30
2	A	301	FLC	CG-CB	-2.29	1.51	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	FLC	OHB-CB-CA	-12.22	80.81	109.40
2	A	302	FLC	OHB-CB-CBC	-10.60	93.98	108.86
2	A	302	FLC	OHB-CB-CG	-6.73	93.65	109.40
2	A	302	FLC	OB2-CBC-CB	6.65	124.61	113.05
2	A	302	FLC	OB1-CBC-CB	-6.39	113.20	122.25
2	A	302	FLC	CA-CB-CBC	6.17	123.37	110.11

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	FLC	CG-CB-CA	5.29	122.95	109.16
2	A	301	FLC	OB2-CBC-CB	3.82	119.68	113.05
2	A	301	FLC	OHB-CB-CBC	-2.83	104.89	108.86
2	A	301	FLC	CB-CA-CAC	-2.22	108.44	113.81
2	B	301	FLC	OA2-CAC-CA	2.17	121.33	114.35
2	A	301	FLC	CA-CB-CBC	2.17	114.76	110.11

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	FLC	CG-CB-CBC-OB1
2	A	301	FLC	CG-CB-CBC-OB2
2	A	301	FLC	OHB-CB-CBC-OB1
2	A	301	FLC	OHB-CB-CBC-OB2
2	A	302	FLC	CAC-CA-CB-CBC
2	A	302	FLC	CA-CB-CG-CGC
2	B	301	FLC	CG-CB-CBC-OB1
2	B	301	FLC	CG-CB-CBC-OB2
2	B	301	FLC	OHB-CB-CBC-OB1
2	B	301	FLC	OHB-CB-CBC-OB2
2	B	301	FLC	CAC-CA-CB-OHB
2	B	301	FLC	OHB-CB-CG-CGC
4	B	302	PEG	O2-C3-C4-O4
2	B	301	FLC	CAC-CA-CB-CG
4	B	302	PEG	C4-C3-O2-C2
2	A	302	FLC	CBC-CB-CG-CGC
2	A	302	FLC	CA-CB-CBC-OB2
2	A	301	FLC	CAC-CA-CB-OHB
2	B	301	FLC	CA-CB-CG-CGC
2	A	302	FLC	CA-CB-CBC-OB1
2	A	302	FLC	CAC-CA-CB-OHB
2	A	302	FLC	CB-CA-CAC-OA1
2	A	302	FLC	CB-CA-CAC-OA2
2	B	301	FLC	CBC-CB-CG-CGC

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	FLC	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	302	FLC	1	0
4	B	302	PEG	5	0
2	A	301	FLC	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/233 (99%)	0.38	6 (2%) 56 47	7, 11, 21, 28	7 (3%)
1	B	231/233 (99%)	0.74	23 (9%) 7 10	8, 17, 28, 35	7 (3%)
All	All	463/466 (99%)	0.56	29 (6%) 20 19	7, 14, 27, 35	14 (3%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	290	SER	9.2
1	B	59	ALA	9.0
1	B	80[A]	VAL	5.4
1	B	289	ALA	5.2
1	A	59	ALA	4.8
1	B	81[A]	LYS	4.5
1	B	173	ILE	4.1
1	B	174	HIS	3.3
1	B	202	ARG	3.2
1	B	167	ASN	2.9
1	A	257	HIS	2.9
1	A	290	SER	2.8
1	B	83[A]	LYS	2.7
1	B	163	GLU	2.6
1	B	172	ALA	2.5
1	B	279[A]	ARG	2.5
1	A	232	LYS	2.4
1	B	178	LYS	2.3
1	B	258	LYS	2.3
1	B	255	HIS	2.3
1	B	60	ILE	2.3
1	B	171	LYS	2.2
1	A	233	HIS	2.2
1	B	118[A]	ARG	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	83	LYS	2.1
1	B	257	HIS	2.0
1	B	79[A]	GLU	2.0
1	B	84[A]	SER	2.0
1	B	194[A]	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	B	104	12/13	0.97	0.09	9,11,13,14	0
1	KCX	A	104	12/13	0.99	0.09	6,8,10,12	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PEG	B	302	7/7	0.74	0.29	31,37,38,39	17
2	FLC	B	301	13/13	0.78	0.36	10,23,29,29	18
2	FLC	A	301	13/13	0.84	0.27	7,17,21,22	18
3	EDO	A	303	4/4	0.90	0.11	23,28,28,28	0
2	FLC	A	302	13/13	0.91	0.15	21,24,29,29	0
3	EDO	A	304	4/4	0.93	0.13	20,24,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.