



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2025 – 02:05 PM EDT

PDB ID : 9CUB / pdb_00009cub
Title : Human STING G230A/R293Q variant bound to diABZI-a1
Authors : Critton, D.A.
Deposited on : 2024-07-26
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	<div> <div>13%</div> <div>77%</div> <div>5%</div> <div>18%</div> </div>
1	B	210	<div> <div>30%</div> <div>73%</div> <div>•</div> <div>24%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5217 atoms, of which 2495 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	172	Total	C	H	N	O	S	1323	1	0
			2684	868	1323	237	251	5			
1	B	159	Total	C	H	N	O	S	1132	0	0
			2333	774	1132	197	225	5			

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	132	MET	-	initiating methionine	UNP Q86WV6
A	133	GLY	-	expression tag	UNP Q86WV6
A	134	SER	-	expression tag	UNP Q86WV6
A	135	SER	-	expression tag	UNP Q86WV6
A	136	HIS	-	expression tag	UNP Q86WV6
A	137	HIS	-	expression tag	UNP Q86WV6
A	138	HIS	-	expression tag	UNP Q86WV6
A	139	HIS	-	expression tag	UNP Q86WV6
A	140	HIS	-	expression tag	UNP Q86WV6
A	141	HIS	-	expression tag	UNP Q86WV6
A	142	SER	-	expression tag	UNP Q86WV6
A	143	SER	-	expression tag	UNP Q86WV6
A	144	GLY	-	expression tag	UNP Q86WV6
A	145	GLU	-	expression tag	UNP Q86WV6
A	146	THR	-	expression tag	UNP Q86WV6
A	147	VAL	-	expression tag	UNP Q86WV6
A	148	ARG	-	expression tag	UNP Q86WV6
A	149	PHE	-	expression tag	UNP Q86WV6
A	150	GLN	-	expression tag	UNP Q86WV6
A	151	GLY	-	expression tag	UNP Q86WV6
A	152	HIS	-	expression tag	UNP Q86WV6
A	153	MET	-	expression tag	UNP Q86WV6
A	154	SER	-	expression tag	UNP Q86WV6
A	230	ALA	GLY	variant	UNP Q86WV6
A	232	ARG	HIS	conflict	UNP Q86WV6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	293	GLN	ARG	variant	UNP Q86WV6
B	132	MET	-	initiating methionine	UNP Q86WV6
B	133	GLY	-	expression tag	UNP Q86WV6
B	134	SER	-	expression tag	UNP Q86WV6
B	135	SER	-	expression tag	UNP Q86WV6
B	136	HIS	-	expression tag	UNP Q86WV6
B	137	HIS	-	expression tag	UNP Q86WV6
B	138	HIS	-	expression tag	UNP Q86WV6
B	139	HIS	-	expression tag	UNP Q86WV6
B	140	HIS	-	expression tag	UNP Q86WV6
B	141	HIS	-	expression tag	UNP Q86WV6
B	142	SER	-	expression tag	UNP Q86WV6
B	143	SER	-	expression tag	UNP Q86WV6
B	144	GLY	-	expression tag	UNP Q86WV6
B	145	GLU	-	expression tag	UNP Q86WV6
B	146	THR	-	expression tag	UNP Q86WV6
B	147	VAL	-	expression tag	UNP Q86WV6
B	148	ARG	-	expression tag	UNP Q86WV6
B	149	PHE	-	expression tag	UNP Q86WV6
B	150	GLN	-	expression tag	UNP Q86WV6
B	151	GLY	-	expression tag	UNP Q86WV6
B	152	HIS	-	expression tag	UNP Q86WV6
B	153	MET	-	expression tag	UNP Q86WV6
B	154	SER	-	expression tag	UNP Q86WV6
B	230	ALA	GLY	variant	UNP Q86WV6
B	232	ARG	HIS	conflict	UNP Q86WV6
B	293	GLN	ARG	variant	UNP Q86WV6

- Molecule 2 is 1,1'-[(2E)-but-2-ene-1,4-diyl]bis{2-[(1-ethyl-3-methyl-1H-pyrazole-5-carbon yl)amino]-7-methoxy-1H-1,3-benzimidazole-5-carboxamide} (CCD ID: A1A3A) (formula: C₃₆H₄₀N₁₂O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	94	36	40	12	6	40	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	70	Total	O	
			70	70	0
3	B	36	Total	O	
			36	36	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Stimulator of interferon genes protein



- Molecule 1: Stimulator of interferon genes protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.97Å 72.64Å 66.43Å 90.00° 98.02° 90.00°	Depositor
Resolution (Å)	25.43 – 1.87 25.43 – 1.87	Depositor EDS
% Data completeness (in resolution range)	68.7 (25.43-1.87) 69.1 (25.43-1.87)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 1.86Å)	Xtriage
Refinement program	BUSTER 2.11.8 (21-NOV-2022)	Depositor
R, R_{free}	0.263 , 0.287 0.252 , 0.278	Depositor DCC
R_{free} test set	9684 reflections (5.31%)	wwPDB-VP
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5217	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1A3A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/1390	0.94	1/1888 (0.1%)
1	B	0.70	0/1222	0.87	0/1663
All	All	0.70	0/2612	0.91	1/3551 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	VAL	N-CA-CB	5.13	116.62	110.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1361	1323	1323	5	0
1	B	1201	1132	1132	4	0
2	A	54	40	0	0	0
3	A	70	0	0	0	0
3	B	36	0	0	1	0
All	All	2722	2495	2455	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:MET:N	1:A:290:LEU:HD13	2.09	0.68
1:B:332:HIS:HE1	3:B:403:HOH:O	1.96	0.48
1:B:299:LEU:O	1:B:306:GLN:HB2	2.15	0.47
1:A:299:LEU:O	1:A:306:GLN:HB2	2.16	0.46
1:A:312:ILE:HG23	1:A:332:HIS:CE1	2.56	0.41
1:A:271:MET:HE3	1:B:157:HIS:CD2	2.56	0.41
1:A:271:MET:CE	1:B:157:HIS:CD2	3.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	167/210 (80%)	162 (97%)	5 (3%)	0	100	100
1	B	149/210 (71%)	146 (98%)	3 (2%)	0	100	100
All	All	316/420 (75%)	308 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/183 (78%)	139 (98%)	3 (2%)	48	34
1	B	120/183 (66%)	118 (98%)	2 (2%)	56	43
All	All	262/366 (72%)	257 (98%)	5 (2%)	52	38

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	212	LEU
1	A	223	ASP
1	A	296	GLU
1	B	165	ILE
1	B	301	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	218	ASN
1	A	266	GLN
1	A	308	ASN
1	B	157	HIS
1	B	332	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1A3A	A	401	-	49,59,59	0.73	0	56,86,86	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1A3A	A	401	-	-	8/27/39/39	0/6/6/6

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	A1A3A	C12-C14-O6-C32
2	A	401	A1A3A	C4-C14-O6-C32
2	A	401	A1A3A	O1-C23-C7-C3
2	A	401	A1A3A	N9-C23-C7-C3
2	A	401	A1A3A	N9-C23-C7-C1
2	A	401	A1A3A	O1-C23-C7-C1
2	A	401	A1A3A	C22-C21-C33-N5
2	A	401	A1A3A	N10-C24-C8-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/210 (81%)	0.88	28 (16%) 5 6	5, 17, 34, 49	1 (0%)
1	B	159/210 (75%)	1.86	63 (39%) 1 0	11, 29, 60, 89	0
All	All	331/420 (78%)	1.35	91 (27%) 2 1	5, 21, 49, 89	1 (0%)

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	MET	7.2
1	B	212	LEU	6.4
1	B	179	ILE	5.8
1	A	215	ALA	5.7
1	B	305	SER	5.6
1	B	185	HIS	5.3
1	B	222	LEU	5.2
1	B	182	TYR	5.2
1	B	300	ALA	5.2
1	B	248	LEU	4.7
1	B	153	MET	4.4
1	B	240	TYR	4.3
1	B	219	ILE	4.3
1	B	303	PRO	4.2
1	B	221	PHE	4.2
1	A	212	LEU	4.0
1	A	323	PHE	3.9
1	B	322	SER	3.8
1	B	250	ASN	3.8
1	B	225	LEU	3.7
1	B	238	ARG	3.6
1	A	305	SER	3.6
1	A	251	GLY	3.5
1	B	176	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	175	LEU	3.5
1	A	274	TYR	3.5
1	B	168	LEU	3.4
1	B	251	GLY	3.4
1	B	247	LEU	3.3
1	A	303	PRO	3.3
1	B	154	SER	3.3
1	B	323	PHE	3.3
1	B	256	THR	3.2
1	B	301	ASP	3.1
1	A	227	GLN	3.1
1	A	211	ASN	3.1
1	B	325	LEU	3.1
1	B	172	LEU	3.0
1	A	214	MET	3.0
1	B	299	LEU	2.9
1	A	241	SER	2.9
1	B	226	PRO	2.8
1	B	210	ASP	2.8
1	B	218	ASN	2.8
1	B	253	ARG	2.8
1	A	213	SER	2.8
1	B	183	ASN	2.7
1	A	275	SER	2.7
1	A	239	VAL	2.7
1	B	211	ASN	2.7
1	B	244	ILE	2.7
1	B	177	ALA	2.6
1	B	239	VAL	2.6
1	B	217	PRO	2.6
1	B	196	GLN	2.6
1	B	216	ASP	2.6
1	B	213	SER	2.6
1	B	181	THR	2.6
1	A	301	ASP	2.5
1	B	302	ALA	2.5
1	A	240	TYR	2.5
1	A	308	ASN	2.5
1	B	194	VAL	2.5
1	B	214	MET	2.4
1	A	333	LEU	2.4
1	A	273	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	220	ARG	2.4
1	A	304	GLU	2.4
1	A	280	SER	2.4
1	A	276	GLN	2.4
1	B	329	VAL	2.3
1	B	285	LEU	2.3
1	A	300	ALA	2.3
1	B	224	LYS	2.3
1	B	215	ALA	2.2
1	B	169	ARG	2.2
1	A	299	LEU	2.2
1	B	223	ASP	2.2
1	B	307	ASN	2.2
1	B	252	GLN	2.1
1	B	178	ARG	2.1
1	B	297	ASP	2.1
1	A	208	VAL	2.1
1	B	330	LEU	2.1
1	B	245	TYR	2.1
1	B	336	GLU	2.1
1	A	317	PRO	2.0
1	B	306	GLN	2.0
1	A	307	ASN	2.0
1	B	173	PRO	2.0
1	B	311	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	$Q < 0.9$
2	A1A3A	A	401	54/54	0.84	0.11	35,39,42,43	40

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.