



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 2, 2024 – 10:09 PM EDT

PDB ID : 4DK1
Title : Crystal Structure of MacA-MexA chimeric protein, containing the Pseudomonas aeruginosa MexA alpha-hairpin domain.
Authors : Xu, Y.; Ha, N.C.
Deposited on : 2012-02-03
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1094 (3.56-3.44)
Clashscore	180529	1045 (3.54-3.46)
Ramachandran outliers	177936	1032 (3.54-3.46)
Sidechain outliers	177891	1033 (3.54-3.46)
RSRZ outliers	164620	1093 (3.56-3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	<div> <div>16%</div> <div>32%</div> <div>23%</div> <div>10%</div> <div>19%</div> </div>
1	B	341	<div> <div>13%</div> <div>35%</div> <div>26%</div> <div>10%</div> <div>16%</div> </div>
1	C	341	<div> <div>18%</div> <div>31%</div> <div>24%</div> <div>8%</div> <div>18%</div> </div>
1	D	341	<div> <div>18%</div> <div>31%</div> <div>24%</div> <div>8%</div> <div>18%</div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative MacA, Multidrug resistance protein mexA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	Se	4	0	0
			2108	1316	355	435	2			
1	B	285	Total	C	N	O	Se	4	0	0
			2191	1369	369	451	2			
1	C	278	Total	C	N	O	Se	4	0	0
			2135	1336	358	439	2			
1	D	278	Total	C	N	O	Se	4	0	0
			2134	1334	359	439	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	GLY	-	expression tag	UNP Q2EHL9
A	27	ALA	-	expression tag	UNP Q2EHL9
A	28	MSE	-	expression tag	UNP Q2EHL9
A	29	ASP	-	expression tag	UNP Q2EHL9
A	68	ILE	VAL	SEE REMARK 999	UNP Q2EHL9
A	85	LEU	ILE	SEE REMARK 999	UNP Q2EHL9
B	26	GLY	-	expression tag	UNP Q2EHL9
B	27	ALA	-	expression tag	UNP Q2EHL9
B	28	MSE	-	expression tag	UNP Q2EHL9
B	29	ASP	-	expression tag	UNP Q2EHL9
B	68	ILE	VAL	SEE REMARK 999	UNP Q2EHL9
B	85	LEU	ILE	SEE REMARK 999	UNP Q2EHL9
C	26	GLY	-	expression tag	UNP Q2EHL9
C	27	ALA	-	expression tag	UNP Q2EHL9
C	28	MSE	-	expression tag	UNP Q2EHL9
C	29	ASP	-	expression tag	UNP Q2EHL9
C	68	ILE	VAL	SEE REMARK 999	UNP Q2EHL9
C	85	LEU	ILE	SEE REMARK 999	UNP Q2EHL9
D	26	GLY	-	expression tag	UNP Q2EHL9
D	27	ALA	-	expression tag	UNP Q2EHL9
D	28	MSE	-	expression tag	UNP Q2EHL9

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	ASP	-	expression tag	UNP Q2EHL9
D	68	ILE	VAL	SEE REMARK 999	UNP Q2EHL9
D	85	LEU	ILE	SEE REMARK 999	UNP Q2EHL9


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative MacA, Multidrug resistance protein mexA



- Molecule 1: Putative MacA, Multidrug resistance protein mexA





4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	151.71Å 151.71Å 216.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.95 – 3.50 19.95 – 3.50	Depositor EDS
% Data completeness (in resolution range)	91.3 (19.95-3.50) 91.4 (19.95-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.60 (at 3.52Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.314 , 0.368 0.287 , 0.341	Depositor DCC
R_{free} test set	1709 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	65.6	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.12 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.31$, $\langle L^2 \rangle = 0.15$	Xtriage
Estimated twinning fraction	0.277 for -h,-k,l	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	8568	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.23	11/2127 (0.5%)	1.47	15/2884 (0.5%)
1	B	1.22	7/2210 (0.3%)	1.44	20/2994 (0.7%)
1	C	0.97	1/2154 (0.0%)	1.35	13/2920 (0.4%)
1	D	1.00	2/2153 (0.1%)	1.35	14/2917 (0.5%)
All	All	1.11	21/8644 (0.2%)	1.41	62/11715 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	23
1	B	0	25
1	C	0	22
1	D	0	24
All	All	0	94

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	196	GLU	CG-CD	9.57	1.66	1.51
1	A	196	GLU	CG-CD	8.69	1.65	1.51
1	B	45	ASN	CG-ND2	-6.84	1.15	1.32
1	A	196	GLU	CB-CG	6.58	1.64	1.52
1	B	121	GLN	CD-NE2	-6.55	1.16	1.32

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	NE-CZ-NH2	16.43	128.51	120.30
1	C	146	ARG	NE-CZ-NH1	16.19	128.40	120.30
1	D	146	ARG	NE-CZ-NH1	16.14	128.37	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	146	ARG	NE-CZ-NH2	14.79	127.70	120.30
1	D	146	ARG	NE-CZ-NH2	-14.74	112.93	120.30

There are no chirality outliers.

5 of 94 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	37	VAL	Peptide
1	A	39	ARG	Peptide
1	A	56	ASN	Peptide
1	A	75	LEU	Peptide
1	A	85	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	0	2133	239	2
1	B	2191	0	2215	265	2
1	C	2135	0	2158	221	1
1	D	2134	0	2156	222	1
All	All	8568	0	8662	912	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 53.

The worst 5 of 912 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:LEU:HB2	1:A:319:VAL:HB	1.29	1.15
1:A:67:LYS:NZ	1:A:68:ILE:O	1.89	1.05
1:C:67:LYS:NZ	1:C:68:ILE:O	1.89	1.05
1:D:67:LYS:NZ	1:D:68:ILE:O	1.89	1.05
1:D:197:ILE:HD11	1:D:202:ILE:HD13	1.42	1.01

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:176:ASN:O	1:D:174:ASN:ND2[3_565]	1.97	0.23
1:A:199:GLU:O	1:B:217:SER:OG[2_665]	2.11	0.09
1:A:176:ASN:O	1:B:174:ASN:ND2[2_665]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/341 (79%)	163 (60%)	51 (19%)	56 (21%)	0	1
1	B	277/341 (81%)	170 (61%)	56 (20%)	51 (18%)	0	1
1	C	270/341 (79%)	164 (61%)	49 (18%)	57 (21%)	0	1
1	D	270/341 (79%)	161 (60%)	59 (22%)	50 (18%)	0	1
All	All	1087/1364 (80%)	658 (60%)	215 (20%)	214 (20%)	0	1

5 of 214 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	38	LYS
1	A	75	LEU
1	A	82	GLY
1	A	121	GLN
1	A	122	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/288 (82%)	132 (56%)	104 (44%)	0	0
1	B	245/288 (85%)	134 (55%)	111 (45%)	0	0
1	C	239/288 (83%)	141 (59%)	98 (41%)	0	1
1	D	238/288 (83%)	134 (56%)	104 (44%)	0	0
All	All	958/1152 (83%)	541 (56%)	417 (44%)	0	0

5 of 417 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	92	THR
1	C	250	ILE
1	D	267	ASN
1	C	121	GLN
1	C	181	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	121	GLN
1	D	266	ASN
1	C	266	ASN
1	D	281	ASN
1	D	63	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	274/341 (80%)	-0.72	0	100 100	7, 60, 178, 245	1 (0%)
1	B	283/341 (82%)	-0.67	0	100 100	7, 63, 166, 285	1 (0%)
1	C	276/341 (80%)	-0.46	1 (0%)	89 79	27, 99, 200, 277	1 (0%)
1	D	276/341 (80%)	-0.52	0	100 100	27, 93, 190, 275	1 (0%)
All	All	1109/1364 (81%)	-0.59	1 (0%)	92 89	7, 82, 185, 285	4 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	177	GLN	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.