



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 9, 2024 – 05:52 PM EST

PDB ID : 2DR0  
Title : Crystal structure of human carboxylesterase in complex with taurocholate  
Authors : Bencharit, S.; Redinbo, M.R.  
Deposited on : 2006-06-02  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	1497 (3.20-3.20)
Ramachandran outliers	177936	1479 (3.20-3.20)
Sidechain outliers	177891	1478 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	542	
1	B	542	
1	C	542	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	379	X	-	-	-
5	TCH	A	101	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TCH	A	102	X	-	X	-
5	TCH	B	201	X	-	-	-
5	TCH	B	202	X	-	X	-
5	TCH	C	301	X	-	-	-
5	TCH	C	302	X	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12988 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Liver carboxylesterase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	0	0
			4130	2662	685	763	20			
1	B	532	Total	C	N	O	S	0	0	0
			4130	2662	685	763	20			
1	C	532	Total	C	N	O	S	0	0	0
			4130	2662	685	763	20			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLN	deletion	UNP P23141
B	?	-	GLN	deletion	UNP P23141
C	?	-	GLN	deletion	UNP P23141

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula:  $C_{11}H_{19}NO_9$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			21	11	1	9		
3	B	1	Total	C	N	O	0	0
			21	11	1	9		
3	C	1	Total	C	N	O	0	0
			21	11	1	9		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is TAUROCHOLIC ACID (three-letter code: TCH) (formula:  $C_{26}H_{45}NO_7S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
5	A	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
5	B	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
5	B	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
5	C	1	Total	C	N	O	S	0	0
			35	26	1	7	1		
5	C	1	Total	C	N	O	S	0	0
			35	26	1	7	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	104	Total	O	0	0
			104	104		
6	B	72	Total	O	0	0
			72	72		
6	C	77	Total	O	0	0
			77	77		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: Liver carboxylesterase 1




#### • Molecule 1: Liver carboxylesterase 1







● Molecule 1: Liver carboxylesterase 1



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.42Å 179.95Å 201.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.61 – 3.20	Depositor
% Data completeness (in resolution range)	98.6 (30.61-3.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.14	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.219 , 0.255	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	12988	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, SIA, TCH, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.49	2/4236 (0.0%)	0.61	0/5754
1	B	0.44	0/4236	0.61	2/5754 (0.0%)
1	C	0.46	1/4236 (0.0%)	0.61	1/5754 (0.0%)
All	All	0.46	3/12708 (0.0%)	0.61	3/17262 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1549	ASN	CG-OD1	-9.65	1.02	1.24
1	A	1549	ASN	CG-ND2	-9.52	1.09	1.32
1	C	3549	ASN	CG-OD1	-5.01	1.12	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2303	PHE	N-CA-C	6.79	129.34	111.00
1	C	3304	LEU	CA-CB-CG	-6.53	100.29	115.30
1	B	2304	LEU	CA-CB-CG	-5.45	102.77	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4130	0	4130	179	0
1	B	4130	0	4131	197	0
1	C	4130	0	4130	210	0
2	A	14	0	13	1	0
2	B	14	0	13	0	0
2	C	14	0	13	0	0
3	A	21	0	18	2	0
3	B	21	0	18	6	0
3	C	21	0	18	5	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
4	C	10	0	0	1	0
5	A	70	0	87	31	0
5	B	70	0	87	39	0
5	C	70	0	87	48	0
6	A	104	0	0	11	0
6	B	72	0	0	13	0
6	C	77	0	0	11	0
All	All	12988	0	12745	589	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 589 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1304:LEU:HD13	5:A:102:TCH:C18	1.69	1.22
1:C:3252:THR:HG21	5:C:302:TCH:O3	1.33	1.22
1:C:3255:LEU:HG	5:C:302:TCH:H2	1.15	1.13
1:C:3301:MET:HB2	1:C:3303:PHE:CZ	1.87	1.08
1:C:3255:LEU:HG	5:C:302:TCH:C2	1.84	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/542 (98%)	473 (89%)	54 (10%)	3 (1%)	22	57
1	B	530/542 (98%)	473 (89%)	53 (10%)	4 (1%)	16	51
1	C	530/542 (98%)	467 (88%)	60 (11%)	3 (1%)	22	57
All	All	1590/1626 (98%)	1413 (89%)	167 (10%)	10 (1%)	22	57

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	3302	LYS
1	B	2185	SER
1	C	3185	SER
1	C	3304	LEU
1	A	1185	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	448/457 (98%)	423 (94%)	25 (6%)	17	50
1	B	448/457 (98%)	426 (95%)	22 (5%)	21	54
1	C	448/457 (98%)	425 (95%)	23 (5%)	20	53
All	All	1344/1371 (98%)	1274 (95%)	70 (5%)	19	52

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	3218	PHE
1	C	3242	ARG
1	C	3381	LEU
1	A	1551	PHE
1	A	1523	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1030	HIS
1	C	3030	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SIA	A	182	-	21,21,21	0.87	0	24,31,31	1.12	2 (8%)
2	NAG	B	279	1	14,14,15	0.87	1 (7%)	17,19,21	0.74	0
3	SIA	B	282	-	21,21,21	1.15	3 (14%)	24,31,31	1.47	3 (12%)
5	TCH	B	201	-	38,38,38	2.84	20 (52%)	59,60,60	3.67	36 (61%)
5	TCH	A	102	-	38,38,38	2.98	18 (47%)	59,60,60	7.49	41 (69%)
5	TCH	B	202	-	38,38,38	2.99	18 (47%)	59,60,60	7.51	43 (72%)
2	NAG	C	379	1	14,14,15	0.79	0	17,19,21	0.86	1 (5%)
5	TCH	A	101	-	38,38,38	2.81	19 (50%)	59,60,60	3.67	35 (59%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	B	285	-	4,4,4	0.41	0	6,6,6	0.17	0
4	SO4	C	185	-	4,4,4	0.42	0	6,6,6	0.14	0
4	SO4	A	284	-	4,4,4	0.42	0	6,6,6	0.21	0
4	SO4	A	184	-	4,4,4	0.38	0	6,6,6	0.09	0
4	SO4	B	385	-	4,4,4	0.44	0	6,6,6	0.18	0
4	SO4	C	384	-	4,4,4	0.44	0	6,6,6	0.17	0
5	TCH	C	302	-	38,38,38	3.32	19 (50%)	59,60,60	7.66	42 (71%)
2	NAG	A	179	1	14,14,15	0.63	0	17,19,21	0.68	1 (5%)
5	TCH	C	301	-	38,38,38	2.86	19 (50%)	59,60,60	3.68	38 (64%)
3	SIA	C	382	-	21,21,21	0.95	1 (4%)	24,31,31	0.88	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TCH	B	201	-	4/4/13/14	9/16/81/81	0/4/4/4
2	NAG	B	279	1	-	2/6/23/26	0/1/1/1
3	SIA	A	182	-	-	7/20/38/38	0/1/1/1
3	SIA	B	282	-	-	9/20/38/38	0/1/1/1
5	TCH	B	202	-	4/4/13/14	10/16/81/81	0/4/4/4
5	TCH	A	102	-	4/4/13/14	9/16/81/81	0/4/4/4
2	NAG	C	379	1	1/1/5/7	5/6/23/26	0/1/1/1
5	TCH	A	101	-	4/4/13/14	9/16/81/81	0/4/4/4
5	TCH	C	302	-	4/4/13/14	8/16/81/81	0/4/4/4
2	NAG	A	179	1	-	4/6/23/26	0/1/1/1
5	TCH	C	301	-	4/4/13/14	9/16/81/81	0/4/4/4
3	SIA	C	382	-	-	12/20/38/38	0/1/1/1

The worst 5 of 118 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	302	TCH	C23-C24	9.31	1.69	1.51
5	A	102	TCH	O24-C24	7.62	1.38	1.23
5	B	201	TCH	C13-C12	6.55	1.64	1.54
5	C	301	TCH	C13-C12	6.50	1.64	1.54
5	C	301	TCH	C11-C12	6.42	1.63	1.53

The worst 5 of 244 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	102	TCH	O2S-S26-C26	44.39	173.81	106.73
5	B	202	TCH	O2S-S26-C26	43.92	173.09	106.73
5	C	302	TCH	O2S-S26-C26	42.79	171.39	106.73
5	A	102	TCH	O3S-S26-O2S	-12.64	79.78	111.40
5	C	302	TCH	O3S-S26-O2S	-12.53	80.06	111.40

5 of 25 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	379	NAG	C1
5	A	101	TCH	C3
5	A	101	TCH	C9
5	A	101	TCH	C5
5	A	101	TCH	C20

5 of 93 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	179	NAG	C8-C7-N2-C2
2	A	179	NAG	O7-C7-N2-C2
2	B	279	NAG	O7-C7-N2-C2
2	C	379	NAG	C8-C7-N2-C2
2	C	379	NAG	O7-C7-N2-C2

There are no ring outliers.


11 monomers are involved in 133 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	182	SIA	2	0
3	B	282	SIA	6	0
5	B	201	TCH	5	0
5	A	102	TCH	26	0
5	B	202	TCH	34	0
5	A	101	TCH	5	0
4	C	185	SO4	1	0
5	C	302	TCH	41	0
2	A	179	NAG	1	0
5	C	301	TCH	7	0
3	C	382	SIA	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.








## Ligand SIA B 282




## Ligand TCH B 201









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.