



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2025 – 12:08 PM EDT

PDB ID : 5EIG
Title : Engineered human cystathionine gamma lyase (E59T, E339V) to deplet cysteine
Authors : Yan, W.; Zhang, J.
Deposited on : 2015-10-29
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	 2% 76% 19% 5%
1	C	405	 % 82% 13% 5%
1	D	405	 2% 82% 13% 5%
1	E	405	 % 81% 14% 5%
1	F	405	 % 81% 12% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	405	<div><div></div><div>83%</div><div>13%</div><div></div></div>
2	B	405	<div><div>4%</div><div></div><div>78%</div><div>17%</div><div>5%</div><div></div></div>
2	H	405	<div><div>%</div><div></div><div>82%</div><div>13%</div><div></div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 24750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cystathionine gamma-lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	P	S	0	0	0
			2995	1909	510	555	1	20			
1	C	388	Total	C	N	O	P	S	0	0	0
			3017	1922	514	560	1	20			
1	D	387	Total	C	N	O	P	S	0	0	0
			3013	1920	513	559	1	20			
1	E	385	Total	C	N	O	P	S	0	0	0
			3002	1914	511	556	1	20			
1	F	382	Total	C	N	O	P	S	0	0	0
			2985	1901	510	553	1	20			
1	G	388	Total	C	N	O	P	S	0	0	0
			3020	1924	515	560	1	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	THR	GLU	engineered mutation	UNP P32929
A	339	VAL	GLU	engineered mutation	UNP P32929
C	59	THR	GLU	engineered mutation	UNP P32929
C	339	VAL	GLU	engineered mutation	UNP P32929
D	59	THR	GLU	engineered mutation	UNP P32929
D	339	VAL	GLU	engineered mutation	UNP P32929
E	59	THR	GLU	engineered mutation	UNP P32929
E	339	VAL	GLU	engineered mutation	UNP P32929
F	59	THR	GLU	engineered mutation	UNP P32929
F	339	VAL	GLU	engineered mutation	UNP P32929
G	59	THR	GLU	engineered mutation	UNP P32929
G	339	VAL	GLU	engineered mutation	UNP P32929


- Molecule 2 is a protein called Cystathionine gamma-lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	386	Total	C	N	O	P	S	0	0	0
			3001	1912	513	556	1	19			
2	H	387	Total	C	N	O	P	S	0	0	0
			3005	1916	512	557	1	19			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	59	THR	GLU	engineered mutation	UNP P32929
B	339	VAL	GLU	engineered mutation	UNP P32929
H	59	THR	GLU	engineered mutation	UNP P32929
H	339	VAL	GLU	engineered mutation	UNP P32929

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CYSTEINE (three-letter code: CYS) (formula: $C_3H_7NO_2S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	S	0	0
			7	3	1	2	1		
4	E	1	Total	C	N	O	S	0	0
			7	3	1	2	1		
4	G	1	Total	C	N	O	S	0	0
			7	3	1	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	72	Total	O	0	0
			72	72		
5	B	57	Total	O	0	0
			57	57		
5	C	62	Total	O	0	0
			62	62		
5	D	67	Total	O	0	0
			67	67		
5	E	94	Total	O	0	0
			94	94		
5	F	85	Total	O	0	0
			85	85		
5	G	128	Total	O	0	0
			128	128		
5	H	101	Total	O	0	0
			101	101		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cystathionine gamma-lyase




• Molecule 1: Cystathionine gamma-lyase




• Molecule 1: Cystathionine gamma-lyase







• Molecule 1: Cystathionine gamma-lyase




• Molecule 1: Cystathionine gamma-lyase




• Molecule 1: Cystathionine gamma-lyase





• Molecule 2: Cystathionine gamma-lyase





● Molecule 2: Cystathionine gamma-lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	113.38Å 163.46Å 181.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.70 40.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.8 (40.00-2.70) 99.0 (40.00-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.191 , 0.233 0.193 , 0.233	Depositor DCC
R_{free} test set	4638 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	31.1	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 29.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	24750	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9187e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, SO4, 5OW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/3029	0.49	0/4109
1	C	0.25	0/3052	0.45	1/4141 (0.0%)
1	D	0.24	0/3048	0.46	0/4136
1	E	0.23	0/3037	0.45	0/4121
1	F	0.24	0/3019	0.45	0/4093
1	G	0.25	0/3055	0.46	0/4145
2	B	0.24	0/3042	0.46	0/4125
2	H	0.25	0/3047	0.46	0/4134
All	All	0.25	0/24329	0.46	1/33004 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	113	VAL	CB-CA-C	-5.23	101.47	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	212	5OW	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2995	0	2965	60	0
1	C	3017	0	2985	37	0
1	D	3013	0	2982	35	0
1	E	3002	0	2972	41	0
1	F	2985	0	2946	37	0
1	G	3020	0	2988	44	0
2	B	3001	0	2987	45	0
2	H	3005	0	2994	45	0
3	B	10	0	0	2	0
3	D	5	0	0	0	0
3	F	5	0	0	0	0
3	H	5	0	0	0	0
4	C	7	0	4	3	0
4	E	7	0	4	1	0
4	G	7	0	4	3	0
5	A	72	0	0	14	0
5	B	57	0	0	7	0
5	C	62	0	0	6	0
5	D	67	0	0	7	0
5	E	94	0	0	13	0
5	F	85	0	0	12	0
5	G	128	0	0	10	0
5	H	101	0	0	11	0
All	All	24750	0	23831	318	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 318 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:114:TYR:HH	4:C:501:CYS:N	1.57	1.01
1:A:364:ARG:NH1	5:A:505:HOH:O	2.01	0.94
1:F:10:GLY:N	5:F:605:HOH:O	2.02	0.91
1:C:384:GLU:OE2	5:C:601:HOH:O	1.88	0.90

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359:VAL:O	1:A:364:ARG:NE	2.06	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/405 (94%)	369 (97%)	8 (2%)	2 (0%)	25	49
1	C	383/405 (95%)	370 (97%)	12 (3%)	1 (0%)	37	61
1	D	382/405 (94%)	369 (97%)	11 (3%)	2 (0%)	25	49
1	E	380/405 (94%)	369 (97%)	10 (3%)	1 (0%)	37	61
1	F	375/405 (93%)	364 (97%)	11 (3%)	0	100	100
1	G	383/405 (95%)	371 (97%)	11 (3%)	1 (0%)	37	61
2	B	379/405 (94%)	368 (97%)	10 (3%)	1 (0%)	37	61
2	H	380/405 (94%)	366 (96%)	13 (3%)	1 (0%)	37	61
All	All	3041/3240 (94%)	2946 (97%)	86 (3%)	9 (0%)	37	61

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	180	GLY
1	A	181	ASP
1	D	181	ASP
1	D	192	SER
2	H	192	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/343 (95%)	324 (99%)	2 (1%)	84	94
1	C	328/343 (96%)	326 (99%)	2 (1%)	84	94
1	D	328/343 (96%)	324 (99%)	4 (1%)	67	86
1	E	327/343 (95%)	327 (100%)	0	100	100
1	F	325/343 (95%)	322 (99%)	3 (1%)	75	90
1	G	328/343 (96%)	327 (100%)	1 (0%)	91	97
2	B	327/343 (95%)	325 (99%)	2 (1%)	84	94
2	H	328/343 (96%)	327 (100%)	1 (0%)	91	97
All	All	2617/2744 (95%)	2602 (99%)	15 (1%)	84	94

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	140	ILE
1	G	102	LYS
1	D	175	ILE
2	H	256	ASN
1	F	256	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	27	GLN
1	F	398	HIS
1	G	324	HIS
1	G	65	ASN
1	D	27	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	LLP	B	212	2	23,24,25	2.51	6 (26%)	25,32,34	1.22	3 (12%)
2	LLP	H	212	2	23,24,25	2.51	6 (26%)	25,32,34	1.24	3 (12%)
1	5OW	C	212	1	30,31,32	0.97	1 (3%)	31,42,44	1.27	4 (12%)
1	5OW	A	212	1	30,31,32	0.95	1 (3%)	31,42,44	1.17	3 (9%)
1	5OW	E	212	1	30,31,32	0.94	1 (3%)	31,42,44	1.22	3 (9%)
1	5OW	D	212	1	30,31,32	1.01	1 (3%)	31,42,44	1.24	3 (9%)
1	5OW	G	212	1	30,31,32	0.94	1 (3%)	31,42,44	1.24	2 (6%)
1	5OW	F	212	1	30,31,32	0.96	1 (3%)	31,42,44	1.11	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LLP	B	212	2	-	8/16/17/19	0/1/1/1
2	LLP	H	212	2	-	7/16/17/19	0/1/1/1
1	5OW	C	212	1	-	8/29/30/32	0/1/1/1
1	5OW	A	212	1	-	10/29/30/32	0/1/1/1
1	5OW	E	212	1	-	11/29/30/32	0/1/1/1
1	5OW	D	212	1	-	10/29/30/32	0/1/1/1
1	5OW	G	212	1	-	11/29/30/32	0/1/1/1
1	5OW	F	212	1	-	11/29/30/32	0/1/1/1

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	212	LLP	C4-C4'	7.23	1.62	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	212	LLP	C4-C4'	7.21	1.62	1.46
2	B	212	LLP	C4'-NZ	5.03	1.44	1.27
2	H	212	LLP	C4'-NZ	5.00	1.43	1.27
2	B	212	LLP	C2'-C2	3.76	1.56	1.50

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	212	LLP	C4-C4'-NZ	-3.32	108.71	124.04
2	H	212	LLP	C4-C4'-NZ	-3.23	109.15	124.04
1	C	212	5OW	C4-C4'-NZ	2.85	122.38	115.53
1	G	212	5OW	C1-C10-S11	-2.79	111.08	114.21
1	G	212	5OW	C4-C4'-NZ	2.73	122.07	115.53

There are no chirality outliers.

5 of 76 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	212	5OW	C-CA-CB-CG
1	A	212	5OW	O-C-CA-CB
1	A	212	5OW	N4-C4'-NZ-CE
1	A	212	5OW	C5'-OP4-P-OP3
1	A	212	5OW	C5'-OP4-P-OP2

There are no ring outliers.

7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	212	LLP	1	0
2	H	212	LLP	4	0
1	C	212	5OW	2	0
1	A	212	5OW	1	0
1	E	212	5OW	1	0
1	G	212	5OW	4	0
1	F	212	5OW	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	D	501	-	4,4,4	0.24	0	6,6,6	0.07	0
4	CYS	E	501	-	5,6,6	1.08	1 (20%)	3,7,7	1.56	1 (33%)
4	CYS	G	501	-	5,6,6	1.07	1 (20%)	3,7,7	1.46	1 (33%)
4	CYS	C	501	-	5,6,6	1.08	1 (20%)	3,7,7	1.59	1 (33%)
3	SO4	B	501	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	B	502	-	4,4,4	0.22	0	6,6,6	0.09	0
3	SO4	F	501	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	H	501	-	4,4,4	0.25	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CYS	G	501	-	-	4/6/6/6	-
4	CYS	E	501	-	-	4/6/6/6	-
4	CYS	C	501	-	-	3/6/6/6	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	501	CYS	OXT-C	-2.29	1.23	1.30
4	G	501	CYS	OXT-C	-2.28	1.23	1.30
4	C	501	CYS	OXT-C	-2.27	1.23	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	501	CYS	OXT-C-O	-2.71	117.94	124.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	501	CYS	OXT-C-O	-2.68	118.01	124.08
4	G	501	CYS	OXT-C-O	-2.52	118.36	124.08

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	501	CYS	O-C-CA-CB
4	E	501	CYS	OXT-C-CA-CB
4	G	501	CYS	O-C-CA-N
4	G	501	CYS	C-CA-CB-SG
4	G	501	CYS	OXT-C-CA-N

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	501	CYS	1	0
4	G	501	CYS	3	0
4	C	501	CYS	3	0
3	B	501	SO4	1	0
3	B	502	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/405 (94%)	0.00	9 (2%) 61 60	17, 32, 73, 96	0
1	C	387/405 (95%)	0.10	5 (1%) 74 74	16, 36, 67, 87	0
1	D	386/405 (95%)	0.04	9 (2%) 61 60	16, 34, 73, 91	0
1	E	384/405 (94%)	-0.37	1 (0%) 90 89	13, 23, 47, 72	0
1	F	381/405 (94%)	-0.14	6 (1%) 70 70	12, 29, 67, 90	0
1	G	387/405 (95%)	-0.49	1 (0%) 90 89	11, 21, 43, 65	0
2	B	385/405 (95%)	0.37	17 (4%) 39 38	21, 43, 78, 98	0
2	H	386/405 (95%)	-0.13	6 (1%) 70 70	13, 28, 57, 76	0
All	All	3079/3240 (95%)	-0.08	54 (1%) 67 67	11, 30, 68, 98	0

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	51	ALA	5.1
2	H	51	ALA	4.2
1	A	104	GLY	4.1
2	H	16	GLN	3.7
1	C	140	ILE	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	5OW	C	212	31/32	0.91	0.13	23,37,45,53	3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	5OW	A	212	31/32	0.93	0.10	18,28,36,37	4
1	5OW	D	212	31/32	0.93	0.10	15,32,39,50	2
1	5OW	E	212	31/32	0.93	0.11	12,23,28,34	3
1	5OW	F	212	31/32	0.93	0.10	16,24,43,52	3
2	LLP	B	212	24/25	0.95	0.09	29,35,44,46	0
1	5OW	G	212	31/32	0.95	0.09	11,19,31,52	3
2	LLP	H	212	24/25	0.95	0.09	18,23,27,28	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CYS	E	501	7/7	0.79	0.18	33,34,51,100	0
3	SO4	H	501	5/5	0.82	0.11	38,41,60,72	0
4	CYS	G	501	7/7	0.82	0.15	23,25,36,67	0
4	CYS	C	501	7/7	0.83	0.11	43,46,57,70	0
3	SO4	D	501	5/5	0.83	0.11	54,56,61,73	0
3	SO4	B	502	5/5	0.83	0.11	54,58,70,76	0
3	SO4	B	501	5/5	0.89	0.12	47,52,69,75	0
3	SO4	F	501	5/5	0.89	0.10	35,37,55,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.