



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2024 – 12:49 AM EDT

PDB ID : 3ETD
Title : Structure of glutamate dehydrogenase complexed with bithionol
Authors : Li, M.; Smith, T.J.
Deposited on : 2008-10-07
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	<div> <div>11%</div> <div> <div></div> <div>70%</div> <div>26%</div> <div>.</div> </div> </div>
1	B	501	<div> <div>7%</div> <div> <div></div> <div>70%</div> <div>26%</div> <div>.</div> </div> </div>
1	C	501	<div> <div>7%</div> <div> <div></div> <div>69%</div> <div>27%</div> <div>.</div> </div> </div>
1	D	501	<div> <div>8%</div> <div> <div></div> <div>71%</div> <div>26%</div> <div>.</div> </div> </div>
1	E	501	<div> <div>8%</div> <div> <div></div> <div>70%</div> <div>26%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	501	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLU	D	550	-	-	X	-
2	GLU	E	550	-	-	X	-
4	GTP	A	553	-	-	-	X
4	GTP	B	553	-	-	-	X
4	GTP	E	553	X	-	-	-
4	GTP	F	553	-	-	-	X
5	B1T	A	552	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 24226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUD1 protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	501	Total	C	N	O	S	0	0	0
			3915	2473	687	736	19			
1	B	501	Total	C	N	O	S	0	0	0
			3915	2473	687	736	19			
1	C	501	Total	C	N	O	S	0	0	0
			3915	2473	687	736	19			
1	D	501	Total	C	N	O	S	0	0	0
			3915	2473	687	736	19			
1	E	501	Total	C	N	O	S	0	0	0
			3915	2473	687	736	19			
1	F	501	Total	C	N	O	S	0	0	0
			3915	2473	687	736	19			

- Molecule 2 is GLUTAMIC ACID (three-letter code: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 9	C 5	N 1	O 3	0	0
2	B	1	Total 9	C 5	N 1	O 3	0	0
2	C	1	Total 9	C 5	N 1	O 3	0	0
2	D	1	Total 9	C 5	N 1	O 3	0	0
2	E	1	Total 9	C 5	N 1	O 3	0	0
2	F	1	Total 9	C 5	N 1	O 3	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	C	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	D	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	E	1	Total 48	C 21	N 7	O 17	P 3	0	0


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).







Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	O	S	0	0
			19	12	4	2	1		
5	B	1	Total	C	Cl	O	S	0	0
			19	12	4	2	1		
5	C	1	Total	C	Cl	O	S	0	0
			19	12	4	2	1		
5	D	1	Total	C	Cl	O	S	0	0
			19	12	4	2	1		
5	E	1	Total	C	Cl	O	S	0	0
			19	12	4	2	1		
5	F	1	Total	C	Cl	O	S	0	0
			19	12	4	2	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	13	Total	O	0	0
			13	13		
6	B	17	Total	O	0	0
			17	17		
6	C	19	Total	O	0	0
			19	19		
6	D	20	Total	O	0	0
			20	20		
6	E	13	Total	O	0	0
			13	13		
6	F	6	Total	O	0	0
			6	6		



• Molecule 1: GLUD1 protein




• Molecule 1: GLUD1 protein



• Molecule 1: GLUD1 protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	122.38Å 101.30Å 166.59Å 90.00° 102.46° 90.00°	Depositor
Resolution (Å)	48.34 – 2.50 48.36 – 2.49	Depositor EDS
% Data completeness (in resolution range)	94.7 (48.34-2.50) 94.1 (48.36-2.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.48Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.239 , 0.257 0.230 , 0.251	Depositor DCC
R_{free} test set	6984 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	24226	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B1T, GTP, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/3998	0.74	3/5396 (0.1%)
1	B	0.44	0/3998	0.91	6/5396 (0.1%)
1	C	0.44	0/3998	0.73	4/5396 (0.1%)
1	D	0.43	0/3998	0.79	3/5396 (0.1%)
1	E	0.43	0/3998	0.73	4/5396 (0.1%)
1	F	0.43	0/3998	0.86	6/5396 (0.1%)
All	All	0.44	0/23988	0.80	26/32376 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	419	ARG	NE-CZ-NH1	-22.85	108.87	120.30
1	F	419	ARG	NE-CZ-NH2	21.56	131.08	120.30
1	B	35	ARG	NE-CZ-NH1	-21.50	109.55	120.30
1	B	35	ARG	NE-CZ-NH2	21.09	130.85	120.30
1	B	44	ARG	NE-CZ-NH1	-19.57	110.51	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	500	PHE	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3915	0	3880	124	0
1	B	3915	0	3880	127	0
1	C	3915	0	3880	125	0
1	D	3915	0	3880	124	0
1	E	3915	0	3880	131	0
1	F	3915	0	3880	123	0
2	A	9	0	5	1	0
2	B	9	0	5	3	0
2	C	9	0	5	0	0
2	D	9	0	5	5	0
2	E	9	0	5	5	0
2	F	9	0	5	2	0
3	A	48	0	26	8	0
3	B	48	0	26	5	0
3	C	48	0	26	6	0
3	D	48	0	26	8	0
3	E	48	0	26	8	0
3	F	48	0	26	9	0
4	A	32	0	12	2	0
4	B	32	0	10	1	0
4	C	32	0	12	1	0
4	D	32	0	12	1	0
4	E	32	0	12	0	0
4	F	32	0	12	0	0
5	A	19	0	4	2	0
5	B	19	0	4	2	0
5	C	19	0	4	3	0
5	D	19	0	4	4	0
5	E	19	0	4	1	0
5	F	19	0	4	2	0
6	A	13	0	0	1	0
6	B	17	0	0	0	0
6	C	19	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	20	0	0	2	0
6	E	13	0	0	1	0
6	F	6	0	0	1	0
All	All	24226	0	23560	724	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 724 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:5:ASP:OD2	1:F:332:THR:HB	1.59	1.00
1:B:3:ARG:HG3	1:B:3:ARG:HH11	1.26	1.00
1:D:332:THR:H	1:D:335:ASN:HD21	1.12	0.98
1:C:150:MET:HG3	5:C:552:B1T:CLAE	2.03	0.95
1:B:150:MET:HG3	5:B:552:B1T:CLAE	2.04	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	499/501 (100%)	464 (93%)	28 (6%)	7 (1%)	11	20
1	B	499/501 (100%)	463 (93%)	30 (6%)	6 (1%)	13	24
1	C	499/501 (100%)	459 (92%)	33 (7%)	7 (1%)	11	20
1	D	499/501 (100%)	462 (93%)	32 (6%)	5 (1%)	15	28
1	E	499/501 (100%)	462 (93%)	33 (7%)	4 (1%)	19	35
1	F	499/501 (100%)	462 (93%)	30 (6%)	7 (1%)	11	20
All	All	2994/3006 (100%)	2772 (93%)	186 (6%)	36 (1%)	13	24

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	SER
1	A	496	ALA
1	B	3	ARG
1	B	313	SER
1	C	2	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/417 (100%)	394 (94%)	23 (6%)	21	41
1	B	417/417 (100%)	393 (94%)	24 (6%)	20	38
1	C	417/417 (100%)	392 (94%)	25 (6%)	19	37
1	D	417/417 (100%)	393 (94%)	24 (6%)	20	38
1	E	417/417 (100%)	393 (94%)	24 (6%)	20	38
1	F	417/417 (100%)	395 (95%)	22 (5%)	22	43
All	All	2502/2502 (100%)	2360 (94%)	142 (6%)	20	39

5 of 142 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	373	LEU
1	E	494	ASN
1	F	279	SER
1	C	4	GLU
1	C	2	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 76 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	335	ASN
1	F	364	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	388	ASN
1	F	132	ASN
1	F	494	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NDP	F	551	-	47,52,52	1.66	8 (17%)	61,80,80	2.63	22 (36%)
3	NDP	E	551	-	47,52,52	1.54	7 (14%)	61,80,80	2.61	18 (29%)
4	GTP	B	553	-	29,34,34	1.96	7 (24%)	35,54,54	5.40	13 (37%)
2	GLU	A	550	-	7,8,9	0.89	0	4,9,11	1.33	0
4	GTP	E	553	-	29,34,34	1.52	7 (24%)	35,54,54	4.27	15 (42%)
2	GLU	F	550	-	7,8,9	1.71	2 (28%)	4,9,11	1.65	1 (25%)
5	B1T	B	552	-	20,20,20	0.76	0	29,29,29	0.85	0
5	B1T	A	552	-	20,20,20	0.78	0	29,29,29	0.99	0
2	GLU	E	550	-	7,8,9	0.99	0	4,9,11	0.88	0
5	B1T	D	552	-	20,20,20	0.82	0	29,29,29	0.86	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLU	D	550	-	7,8,9	0.98	0	4,9,11	0.79	0
4	GTP	F	553	-	29,34,34	2.87	5 (17%)	35,54,54	4.55	14 (40%)
3	NDP	B	551	-	47,52,52	1.89	7 (14%)	61,80,80	2.68	22 (36%)
3	NDP	A	551	-	47,52,52	1.89	9 (19%)	61,80,80	2.79	27 (44%)
3	NDP	D	551	-	47,52,52	1.92	9 (19%)	61,80,80	2.71	23 (37%)
4	GTP	C	553	-	29,34,34	2.24	6 (20%)	35,54,54	1.50	7 (20%)
5	B1T	F	552	-	20,20,20	0.83	1 (5%)	29,29,29	0.96	0
4	GTP	D	553	-	29,34,34	2.51	6 (20%)	35,54,54	3.13	13 (37%)
2	GLU	B	550	-	7,8,9	0.98	0	4,9,11	1.08	0
3	NDP	C	551	-	47,52,52	2.04	8 (17%)	61,80,80	2.67	22 (36%)
5	B1T	C	552	-	20,20,20	0.76	0	29,29,29	0.86	0
5	B1T	E	552	-	20,20,20	0.79	0	29,29,29	0.95	0
2	GLU	C	550	-	7,8,9	1.03	0	4,9,11	1.33	0
4	GTP	A	553	-	29,34,34	1.87	4 (13%)	35,54,54	3.15	14 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '–' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	F	551	-	-	2/30/77/77	0/5/5/5
3	NDP	E	551	-	-	2/30/77/77	0/5/5/5
4	GTP	B	553	-	-	2/18/38/38	0/3/3/3
2	GLU	A	550	-	-	2/6/7/9	-
4	GTP	E	553	-	1/1/7/7	2/18/38/38	0/3/3/3
2	GLU	F	550	-	-	2/6/7/9	-
5	B1T	B	552	-	-	0/4/4/4	0/2/2/2
5	B1T	A	552	-	-	0/4/4/4	0/2/2/2
2	GLU	E	550	-	-	2/6/7/9	-
5	B1T	D	552	-	-	0/4/4/4	0/2/2/2
2	GLU	D	550	-	-	5/6/7/9	-
4	GTP	F	553	-	-	1/18/38/38	0/3/3/3
3	NDP	B	551	-	-	2/30/77/77	0/5/5/5
3	NDP	A	551	-	-	5/30/77/77	0/5/5/5
3	NDP	D	551	-	-	3/30/77/77	0/5/5/5
4	GTP	C	553	-	-	1/18/38/38	0/3/3/3
5	B1T	F	552	-	-	0/4/4/4	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	D	553	-	-	4/18/38/38	0/3/3/3
2	GLU	B	550	-	-	3/6/7/9	-
3	NDP	C	551	-	-	5/30/77/77	0/5/5/5
5	B1T	C	552	-	-	0/4/4/4	0/2/2/2
5	B1T	E	552	-	-	0/4/4/4	0/2/2/2
2	GLU	C	550	-	-	1/6/7/9	-
4	GTP	A	553	-	-	3/18/38/38	0/3/3/3

The worst 5 of 86 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	553	GTP	O4'-C1'	12.07	1.56	1.40
3	C	551	NDP	PN-O3	-9.56	1.49	1.59
4	D	553	GTP	PB-O3A	9.12	1.69	1.59
3	D	551	NDP	PN-O3	-7.86	1.51	1.59
3	A	551	NDP	PN-O3	-7.47	1.51	1.59

The worst 5 of 211 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	553	GTP	C4'-O4'-C1'	-25.78	86.32	109.92
4	E	553	GTP	C5'-C4'-C3'	14.60	167.75	115.21
4	F	553	GTP	O4'-C1'-N9	13.54	126.70	108.75
4	F	553	GTP	O4'-C4'-C3'	-12.04	81.25	105.15
4	F	553	GTP	O4'-C4'-C5'	-10.91	74.38	109.33

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	E	553	GTP	C4'

5 of 47 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	550	GLU	N-CA-CB-CG
2	A	550	GLU	C-CA-CB-CG
2	C	550	GLU	O-C-CA-CB
2	D	550	GLU	O-C-CA-CB
2	D	550	GLU	N-CA-CB-CG


There are no ring outliers.

21 monomers are involved in 73 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	551	NDP	9	0
3	E	551	NDP	8	0
4	B	553	GTP	1	0
2	A	550	GLU	1	0
2	F	550	GLU	2	0
5	B	552	B1T	2	0
5	A	552	B1T	2	0
2	E	550	GLU	5	0
5	D	552	B1T	4	0
2	D	550	GLU	5	0
3	B	551	NDP	5	0
3	A	551	NDP	8	0
3	D	551	NDP	8	0
4	C	553	GTP	1	0
5	F	552	B1T	2	0
4	D	553	GTP	1	0
2	B	550	GLU	3	0
3	C	551	NDP	6	0
5	C	552	B1T	3	0
5	E	552	B1T	1	0
4	A	553	GTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand GTP B 553




Ligand GTP E 553





Ligand B1T A 552




Ligand GTP F 553













Ligand GTP C 553




Ligand B1T F 552









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	501/501 (100%)	0.74	56 (11%) 5 4	36, 60, 90, 105	0
1	B	501/501 (100%)	0.57	34 (6%) 17 17	36, 61, 90, 104	0
1	C	501/501 (100%)	0.51	33 (6%) 18 19	34, 59, 90, 107	0
1	D	501/501 (100%)	0.63	42 (8%) 11 11	36, 60, 91, 105	0
1	E	501/501 (100%)	0.66	42 (8%) 11 11	36, 60, 90, 105	0
1	F	501/501 (100%)	0.79	79 (15%) 2 1	37, 61, 90, 105	0
All	All	3006/3006 (100%)	0.65	286 (9%) 8 8	34, 60, 91, 107	0

The worst 5 of 286 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	11.8
1	C	1	ALA	11.1
1	E	1	ALA	10.1
1	D	1	ALA	9.4
1	D	424	HIS	9.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GTP	F	553	32/32	0.52	0.54	116,119,134,135	0
4	GTP	B	553	32/32	0.53	0.48	105,112,119,119	0
5	B1T	E	552	19/19	0.56	0.38	105,108,114,116	0
5	B1T	B	552	19/19	0.58	0.37	103,109,112,113	0
5	B1T	A	552	19/19	0.58	0.49	106,110,114,117	0
4	GTP	D	553	32/32	0.64	0.31	105,111,117,118	0
4	GTP	E	553	32/32	0.64	0.30	100,110,124,125	0
4	GTP	A	553	32/32	0.65	0.42	111,116,125,126	0
4	GTP	C	553	32/32	0.66	0.33	104,113,129,131	0
5	B1T	C	552	19/19	0.67	0.33	104,108,114,115	0
5	B1T	D	552	19/19	0.69	0.32	104,107,112,113	0
2	GLU	E	550	9/10	0.78	0.39	68,73,74,75	0
5	B1T	F	552	19/19	0.80	0.24	93,95,99,100	0
3	NDP	B	551	48/48	0.84	0.22	64,71,80,85	0
3	NDP	E	551	48/48	0.84	0.23	67,81,86,87	0
3	NDP	D	551	48/48	0.85	0.26	52,63,74,79	0
3	NDP	F	551	48/48	0.85	0.22	74,83,85,87	0
3	NDP	A	551	48/48	0.87	0.20	59,66,73,77	0
2	GLU	A	550	9/10	0.88	0.27	58,61,62,63	0
3	NDP	C	551	48/48	0.90	0.21	43,55,66,72	0
2	GLU	F	550	9/10	0.91	0.20	67,68,70,71	0
2	GLU	B	550	9/10	0.92	0.24	54,58,60,60	0
2	GLU	D	550	9/10	0.93	0.21	41,46,50,51	0
2	GLU	C	550	9/10	0.94	0.19	52,55,56,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GTP F 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP B 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B1T A 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP D 553:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP E 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP C 553:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B1T F 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP B 551:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NDP E 551:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP D 551:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NDP F 551:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP A 551:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP C 551:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.