



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 07:32 PM EDT

PDB ID : 1F06
Title : THREE DIMENSIONAL STRUCTURE OF THE TERNARY COMPLEX
OF CORYNEBACTERIUM GLUTAMICUM DIAMINOPIMELATE DEHY-
DROGENASE NADPH-L-2-AMINO-6-METHYLENE-PIMELATE
Authors : Cirilli, M.; Scapin, G.; Sutherland, A.; Caplan, J.F.; Vederas, J.C.; Blanchard,
J.S.
Deposited on : 2000-05-14
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>18%</div> <div>.</div> </div> </div>
1	B	320	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>.</div> </div> </div>

2 Entry composition ⓘ


There are 4 unique types of molecules in this entry. The entry contains 5175 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MESO-DIAMINOPIMELATE D-DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	0	0
			2475	1543	442	480	10			
1	B	320	Total	C	N	O	S	0	0	0
			2475	1543	442	480	10			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			13	8	1	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total	O	0	0
			54	54		
4	B	62	Total	O	0	0
			62	62		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MESO-DIAMINOPIMELATE D-DEHYDROGENASE



• Molecule 1: MESO-DIAMINOPIMELATE D-DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.55Å 65.59Å 84.28Å 90.00° 106.52° 90.00°	Depositor
Resolution (Å)	48.60 – 2.10 48.62 – 2.13	Depositor EDS
% Data completeness (in resolution range)	(Not available) (48.60-2.10) 88.2 (48.62-2.13)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.12Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.179 , 0.209 0.185 , 0.217	Depositor DCC
R_{free} test set	1027 reflections (2.60%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5175	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, 2NP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2528	0.63	0/3432
1	B	0.36	0/2528	0.64	1/3432 (0.0%)
All	All	0.36	0/5056	0.64	1/6864 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	648	LEU	CA-CB-CG	5.33	127.55	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2475	0	2378	38	0
1	B	2475	0	2375	44	0
2	A	48	0	26	1	0
2	B	48	0	26	2	0
3	B	13	0	11	1	0
4	A	54	0	0	1	0
4	B	62	0	0	1	0
All	All	5175	0	4816	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:HIS:HB3	1:A:167:GLN:HG2	1.52	0.90
1:A:311:ASN:ND2	1:A:311:ASN:H	1.74	0.86
1:B:811:ASN:H	1:B:811:ASN:ND2	1.73	0.83
1:B:648:LEU:HD21	1:B:666:VAL:HG22	1.63	0.80
1:B:667:GLN:HE22	1:B:695:ARG:HE	1.32	0.78
1:B:667:GLN:HE21	1:B:695:ARG:HB3	1.52	0.75
1:B:628:ARG:HE	1:B:642:THR:HG21	1.51	0.74
1:B:667:GLN:NE2	1:B:695:ARG:HE	1.85	0.74
1:B:531:VAL:HG21	1:B:559:VAL:HG22	1.69	0.73
1:A:128:ARG:HE	1:A:142:THR:HG21	1.52	0.72
1:A:31:VAL:HG21	1:A:59:VAL:HG22	1.73	0.70
1:B:525:GLN:NE2	1:B:783:ARG:HH22	1.94	0.65
1:A:12:GLY:O	1:A:16:ARG:HG3	1.97	0.65
1:A:78:LYS:O	1:A:81:GLN:HG2	1.97	0.64
1:A:320:VAL:HA	1:B:625:SER:HB3	1.79	0.64
1:A:25:GLN:HE21	1:A:283:ARG:HH22	1.46	0.64
1:A:25:GLN:NE2	1:A:283:ARG:HH22	1.96	0.63
1:B:525:GLN:HE21	1:B:783:ARG:HH22	1.46	0.63
1:B:578:LYS:O	1:B:581:GLN:HG2	1.98	0.63
1:A:311:ASN:H	1:A:311:ASN:HD22	1.46	0.62
1:B:811:ASN:H	1:B:811:ASN:HD22	1.46	0.61
1:B:512:GLY:O	1:B:516:ARG:HG3	1.99	0.61
1:B:754:THR:OG1	1:B:759:HIS:HE1	1.83	0.61
1:A:86:VAL:HG21	1:A:282:GLY:HA2	1.83	0.59
1:A:254:THR:OG1	1:A:259:HIS:HE1	1.85	0.59
1:A:125:SER:HB3	1:B:820:VAL:HA	1.84	0.58
1:B:586:VAL:HG21	1:B:782:GLY:HA2	1.86	0.56
1:A:216:MET:O	1:A:221:VAL:HG23	2.06	0.56
1:A:6:VAL:O	1:A:31:VAL:HG22	2.10	0.51
1:B:506:VAL:O	1:B:531:VAL:HG22	2.11	0.51
1:B:649:SER:OG	1:B:652:HIS:HD2	1.93	0.51
1:B:709:ILE:O	1:B:713:ILE:HG13	2.11	0.51
1:A:59:VAL:HG11	1:A:62:LEU:HD13	1.92	0.50
1:B:559:VAL:HG11	1:B:562:LEU:HD13	1.94	0.50
1:B:717:PRO:HA	1:B:721:VAL:HG22	1.93	0.49
1:B:695:ARG:NH2	3:B:950:2NP:O12	2.44	0.49
1:A:152:HIS:HB3	1:A:167:GLN:CG	2.34	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:VAL:HG12	1:A:61:VAL:HG13	1.96	0.47
1:B:660:PRO:O	1:B:708:ARG:NH2	2.46	0.47
1:B:774:THR:OG1	2:B:901:NDP:H41N	2.14	0.47
1:A:7:ALA:HB3	1:A:62:LEU:HD12	1.97	0.47
1:B:593:ARG:NH2	4:B:1033:HOH:O	2.46	0.47
1:A:201:ALA:O	1:A:206:HIS:HE1	1.97	0.47
1:B:506:VAL:HG12	1:B:561:VAL:HG13	1.96	0.47
1:B:684:ALA:O	1:B:687:LEU:HB2	2.15	0.47
1:B:525:GLN:HE21	1:B:783:ARG:NH2	2.13	0.47
1:B:710:GLU:O	1:B:714:ARG:HG3	2.14	0.46
1:B:620:ASP:CG	1:B:628:ARG:HH22	2.19	0.46
1:A:310:GLU:HB2	1:A:315:LEU:HD13	1.97	0.46
1:B:513:ASN:HB2	2:B:901:NDP:O1N	2.16	0.46
1:B:717:PRO:HA	1:B:721:VAL:CG2	2.46	0.46
1:A:286:HIS:O	1:A:290:GLN:HG2	2.16	0.45
1:B:507:ALA:HB3	1:B:562:LEU:HD12	1.98	0.45
1:A:216:MET:C	1:A:221:VAL:HG23	2.37	0.45
1:A:51:ASP:O	1:A:55:HIS:HD2	2.00	0.45
1:A:143:PHE:HB3	1:A:171:PRO:HB3	1.98	0.44
1:A:186:ASP:O	1:A:187:LEU:HD23	2.17	0.44
1:A:205:ASP:O	1:A:209:ILE:HG13	2.17	0.44
1:A:72:ILE:HB	1:A:73:PRO:HD3	1.98	0.44
1:A:88:THR:O	2:A:401:NDP:H2N	2.18	0.44
1:B:572:ILE:HB	1:B:573:PRO:HD3	1.99	0.44
1:B:810:GLU:HB2	1:B:815:LEU:HD13	1.99	0.43
1:A:120:ASP:CG	1:A:128:ARG:HH22	2.22	0.43
1:A:144:TRP:O	1:A:171:PRO:HD3	2.19	0.43
1:A:25:GLN:HE21	1:A:283:ARG:NH2	2.14	0.43
1:B:551:ASP:O	1:B:555:HIS:HD2	2.01	0.43
1:B:663:GLN:HE21	1:B:701:ALA:HA	1.84	0.43
1:A:311:ASN:HD22	1:A:311:ASN:N	2.16	0.42
1:A:145:GLY:HA2	1:A:146:PRO:C	2.40	0.42
1:B:695:ARG:HG3	1:B:723:TYR:HB3	2.00	0.42
1:A:20:LYS:HG2	1:A:21:LEU:HD13	2.01	0.42
1:A:156:LEU:HD12	1:A:156:LEU:HA	1.86	0.42
1:B:659:ILE:HA	1:B:660:PRO:HD2	1.92	0.42
1:B:786:HIS:O	1:B:790:GLN:HG2	2.19	0.42
1:B:811:ASN:ND2	1:B:811:ASN:N	2.50	0.42
1:A:142:THR:HG23	4:A:1118:HOH:O	2.20	0.41
1:B:520:LYS:HG2	1:B:521:LEU:HD13	2.01	0.41
1:A:159:ILE:HD11	1:A:216:MET:CE	2.51	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:667:GLN:NE2	1:B:695:ARG:HB3	2.27	0.40
1:B:648:LEU:HD21	1:B:666:VAL:CG2	2.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/320 (99%)	305 (96%)	13 (4%)	0	100	100
1	B	318/320 (99%)	306 (96%)	12 (4%)	0	100	100
All	All	636/640 (99%)	611 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/261 (100%)	229 (88%)	31 (12%)	5	2
1	B	260/261 (100%)	233 (90%)	27 (10%)	7	4
All	All	520/522 (100%)	462 (89%)	58 (11%)	6	3

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	18	VAL
1	A	21	LEU
1	A	30	LEU
1	A	40	LEU
1	A	49	VAL
1	A	61	VAL
1	A	62	LEU
1	A	66	MET
1	A	70	THR
1	A	74	GLU
1	A	78	LYS
1	A	100	GLN
1	A	114	LEU
1	A	125	SER
1	A	142	THR
1	A	156	LEU
1	A	158	ARG
1	A	167	GLN
1	A	186	ASP
1	A	190	LYS
1	A	191	GLN
1	A	202	ASP
1	A	207	GLU
1	A	224	GLU
1	A	237	SER
1	A	238	GLU
1	A	244	HIS
1	A	307	LEU
1	A	311	ASN
1	A	315	LEU
1	B	502	THR
1	B	518	VAL
1	B	521	LEU
1	B	530	LEU
1	B	540	LEU
1	B	549	VAL
1	B	561	VAL
1	B	562	LEU
1	B	566	MET
1	B	570	THR
1	B	574	GLU
1	B	578	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	600	GLN
1	B	614	LEU
1	B	625	SER
1	B	642	THR
1	B	648	LEU
1	B	666	VAL
1	B	676	LEU
1	B	677	GLU
1	B	686	ASP
1	B	695	ARG
1	B	702	ASP
1	B	740	THR
1	B	744	HIS
1	B	811	ASN
1	B	815	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	25	GLN
1	A	138	HIS
1	A	206	HIS
1	A	259	HIS
1	A	286	HIS
1	A	311	ASN
1	B	503	ASN
1	B	525	GLN
1	B	555	HIS
1	B	638	HIS
1	B	650	GLN
1	B	652	HIS
1	B	663	GLN
1	B	667	GLN
1	B	759	HIS
1	B	811	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	2NP	B	950	-	10,12,12	1.91	2 (20%)	11,15,15	2.17	3 (27%)
2	NDP	A	401	-	47,52,52	2.02	7 (14%)	61,80,80	2.22	11 (18%)
2	NDP	B	901	-	47,52,52	2.15	9 (19%)	61,80,80	2.17	9 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2NP	B	950	-	-	0/14/14/14	-
2	NDP	A	401	-	-	10/30/77/77	0/5/5/5
2	NDP	B	901	-	-	4/30/77/77	0/5/5/5

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NDP	O7N-C7N	8.70	1.44	1.24
2	B	901	NDP	O7N-C7N	8.03	1.43	1.24
2	B	901	NDP	C4N-C3N	-7.14	1.36	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NDP	C4N-C3N	-6.22	1.38	1.50
3	B	950	2NP	C7-C6	5.11	1.52	1.49
2	B	901	NDP	C7N-C3N	-4.96	1.38	1.48
2	A	401	NDP	C7N-C3N	-3.88	1.40	1.48
2	B	901	NDP	C4N-C5N	-3.78	1.39	1.49
2	A	401	NDP	C4N-C5N	-3.10	1.41	1.49
3	B	950	2NP	O11-C1	2.97	1.30	1.22
2	A	401	NDP	C2A-N3A	2.84	1.36	1.32
2	A	401	NDP	O4B-C1B	2.55	1.44	1.40
2	B	901	NDP	C6N-C5N	2.48	1.40	1.33
2	A	401	NDP	C6N-C5N	2.45	1.40	1.33
2	B	901	NDP	O4B-C1B	2.34	1.44	1.40
2	B	901	NDP	P2B-O2X	-2.22	1.46	1.54
2	B	901	NDP	C2A-N3A	2.14	1.35	1.32
2	B	901	NDP	P2B-O3X	-2.09	1.47	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NDP	C1B-N9A-C4A	10.99	145.95	126.64
2	B	901	NDP	C1B-N9A-C4A	10.76	145.54	126.64
2	B	901	NDP	O4B-C1B-N9A	7.15	118.23	108.75
2	A	401	NDP	O4B-C1B-N9A	7.03	118.06	108.75
3	B	950	2NP	O72-C7-C6	4.85	120.30	114.89
2	A	401	NDP	C2B-C1B-N9A	4.83	123.28	112.56
2	B	901	NDP	C2B-C1B-N9A	4.65	122.89	112.56
2	A	401	NDP	O2B-C2B-C1B	3.48	122.27	110.05
2	B	901	NDP	O2B-C2B-C1B	3.47	122.25	110.05
2	B	901	NDP	C4A-C5A-N7A	3.43	112.96	109.34
2	A	401	NDP	O5B-C5B-C4B	3.40	120.56	108.99
2	B	901	NDP	O5B-C5B-C4B	3.26	120.09	108.99
2	A	401	NDP	C4A-C5A-N7A	3.19	112.71	109.34
2	A	401	NDP	O5D-C5D-C4D	3.06	119.40	108.99
2	B	901	NDP	O5D-C5D-C4D	2.97	119.11	108.99
3	B	950	2NP	C4-C5-C6	-2.87	108.71	113.35
3	B	950	2NP	C5-C6-C7	2.52	120.07	116.79
2	B	901	NDP	O3B-C3B-C4B	-2.41	104.16	111.08
2	A	401	NDP	C3D-C2D-C1D	2.39	105.98	101.46
2	A	401	NDP	O3B-C3B-C4B	-2.23	104.67	111.08
2	A	401	NDP	O2B-C2B-C3B	2.09	119.17	111.68
2	B	901	NDP	O3B-C3B-C2B	-2.08	105.36	111.19
2	A	401	NDP	C6N-N1N-C2N	2.04	121.50	119.32

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NDP	C5D-O5D-PN-O3
2	A	401	NDP	C5D-O5D-PN-O2N
2	B	901	NDP	O4D-C1D-N1N-C2N
2	A	401	NDP	PN-O3-PA-O5B
2	B	901	NDP	C1B-C2B-O2B-P2B
2	A	401	NDP	PA-O3-PN-O2N
2	A	401	NDP	O4D-C1D-N1N-C2N
2	A	401	NDP	C3B-C4B-C5B-O5B
2	B	901	NDP	C3B-C4B-C5B-O5B
2	A	401	NDP	C2N-C3N-C7N-N7N
2	B	901	NDP	C2N-C3N-C7N-N7N
2	A	401	NDP	O4D-C4D-C5D-O5D
2	A	401	NDP	C1B-C2B-O2B-P2B
2	A	401	NDP	PA-O3-PN-O1N


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	950	2NP	1	0
2	A	401	NDP	1	0
2	B	901	NDP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/320 (100%)	-0.31	2 (0%) 89 91	14, 26, 46, 67	0
1	B	320/320 (100%)	-0.37	4 (1%) 77 80	13, 23, 44, 74	0
All	All	640/640 (100%)	-0.34	6 (0%) 84 86	13, 24, 45, 74	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	501	MET	6.4
1	A	1	MET	2.8
1	B	686	ASP	2.6
1	B	570	THR	2.6
1	A	186	ASP	2.5
1	B	502	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NDP	A	401	48/48	0.95	0.10	18,33,37,39	0
3	2NP	B	950	13/13	0.96	0.10	14,17,24,29	0
2	NDP	B	901	48/48	0.98	0.09	12,18,27,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.