



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 04:52 PM EDT

PDB ID : 6F0Q  
Title : Crystal structure of Pizza6-AYW  
Authors : Noguchi, H.; De Zitter, E.; Van Meervelt, L.; Voet, A.R.D.  
Deposited on : 2017-11-20  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>18%</div> <div>.</div> </div> </div>
1	B	256	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4604 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pizza6-AYW.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	0	27	0
			1962	1214	350	398			
1	B	251	Total	C	N	O	0	28	0
			1979	1224	352	403			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	322	Total	O	0	0
			322	322		
2	B	341	Total	O	0	0
			341	341		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Pizza6-AYW



#### • Molecule 1: Pizza6-AYW



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.01Å 70.35Å 58.21Å 90.00° 100.60° 90.00°	Depositor
Resolution (Å)	26.50 – 1.30 26.50 – 1.30	Depositor EDS
% Data completeness (in resolution range)	94.7 (26.50-1.30) 94.7 (26.50-1.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.00 (at 1.30Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.155 , 0.192 0.156 , 0.192	Depositor DCC
$R_{free}$ test set	4205 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	7.7	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 41.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4604	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8106e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	2/2003 (0.1%)	0.89	3/2766 (0.1%)
1	B	0.82	4/2014 (0.2%)	0.95	5/2783 (0.2%)
All	All	0.77	6/4017 (0.1%)	0.92	8/5549 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	226[A]	SER	C-N	-7.90	1.18	1.33
1	B	226[B]	SER	C-N	-7.90	1.18	1.33
1	B	16[A]	SER	C-N	-6.55	1.21	1.33
1	B	16[B]	SER	C-N	-6.55	1.21	1.33
1	A	100[A]	SER	C-N	-5.88	1.22	1.33
1	A	100[B]	SER	C-N	-5.88	1.22	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	188[A]	VAL	CG1-CB-CG2	8.57	124.62	110.90
1	B	188[B]	VAL	CG1-CB-CG2	8.57	124.62	110.90
1	B	245	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	B	77	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	161	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	B	119	ARG	NE-CZ-NH2	-5.31	117.65	120.30
1	A	77	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	A	203	ARG	NE-CZ-NH2	-5.13	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1962	0	1928	25	0
1	B	1979	0	1940	34	0
2	A	322	0	0	2	0
2	B	341	0	0	2	0
All	All	4604	0	3868	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16[B]:SER:O	2:B:301:HOH:O	1.79	1.00
1:A:104[B]:VAL:HG12	1:A:110:VAL:HG22	1.66	0.76
1:B:146[B]:VAL:HG12	1:B:152:VAL:HG22	1.75	0.69
1:B:78[B]:VAL:HG11	1:B:112[B]:VAL:HG11	1.79	0.65
1:B:120:VAL:HG21	1:B:154:VAL:HG21	1.79	0.63
1:A:230[B]:VAL:HG12	1:A:236:VAL:HG22	1.82	0.61
1:A:37[B]:VAL:CG1	1:A:46:GLN:HG3	2.32	0.59
1:A:36:VAL:HG21	1:A:70[A]:VAL:HG21	1.86	0.58
1:A:97:ASN:HB2	2:A:554:HOH:O	2.02	0.57
1:A:204:VAL:HG21	1:A:238[A]:VAL:HG21	1.87	0.57
1:B:104[B]:VAL:HG22	1:B:110:VAL:HG22	1.87	0.56
1:B:29[B]:THR:HG23	1:B:60:VAL:HB	1.88	0.55
1:B:184[A]:SER:OG	1:B:226[A]:SER:HA	2.06	0.55
1:B:78[B]:VAL:HG23	1:B:91:LEU:HB2	1.89	0.55
1:A:29[A]:THR:HG23	1:A:60:VAL:HB	1.90	0.53
1:B:1:SER:N	1:B:251:ALA:O	2.42	0.53
1:B:162[B]:VAL:HG23	1:B:175:LEU:HB2	1.91	0.52
1:B:20[B]:VAL:HG22	1:B:26:VAL:HG22	1.92	0.52
1:A:62[B]:VAL:HG22	1:A:68:VAL:HG22	1.90	0.52
1:B:28[A]:VAL:HG21	1:B:246:VAL:HG21	1.92	0.52
1:A:78:VAL:HG21	1:A:112:VAL:HG21	1.91	0.51
1:B:162[B]:VAL:CG2	1:B:175:LEU:HB2	2.41	0.51
1:B:162[A]:VAL:HG21	1:B:196:VAL:HG21	1.93	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:VAL:HG21	1:B:70[B]:VAL:HG11	1.93	0.49
1:A:104[B]:VAL:HG23	2:A:489:HOH:O	2.12	0.49
1:B:37[B]:VAL:HG12	1:B:48:VAL:HG22	1.95	0.49
1:B:182:THR:O	1:B:199:HIS:HB3	2.12	0.49
1:B:78[B]:VAL:CG2	1:B:91:LEU:HB2	2.43	0.48
1:A:219:PHE:CZ	1:A:238[B]:VAL:HG22	2.48	0.48
1:A:56:THR:O	1:A:73:HIS:HB3	2.13	0.48
1:A:82:ALA:O	1:A:85:SER:HB3	2.13	0.48
1:B:197:THR:HG23	1:B:228:VAL:HB	1.96	0.47
1:A:120:VAL:HG21	1:A:154:VAL:HG21	1.96	0.47
1:B:140:THR:O	1:B:157:HIS:HB3	2.15	0.47
1:A:37[B]:VAL:CG1	1:A:46:GLN:CG	2.93	0.46
1:A:197[B]:THR:OG1	1:A:226[B]:SER:O	2.23	0.46
1:A:28:VAL:HG21	1:A:246:VAL:HG21	1.97	0.46
1:A:224:THR:O	1:A:241:HIS:HB3	2.15	0.46
1:A:140:THR:O	1:A:157:HIS:HB3	2.17	0.45
1:A:98:THR:O	1:A:115:HIS:HB3	2.16	0.45
1:B:155[A]:THR:HG23	1:B:186:VAL:HB	1.99	0.45
1:B:153:TRP:CZ3	1:B:188[B]:VAL:HG21	2.52	0.45
1:B:153:TRP:CE3	1:B:188[A]:VAL:HG21	2.51	0.45
1:A:188[B]:VAL:HG22	1:A:194:VAL:HG22	2.00	0.44
1:B:104[B]:VAL:HG12	1:B:105:ASP:O	2.18	0.44
1:B:14:THR:O	1:B:31:HIS:HB3	2.17	0.43
1:B:64:SER:HB2	2:B:540:HOH:O	2.18	0.43
1:B:188[A]:VAL:HG22	1:B:194:VAL:HG22	2.01	0.42
1:B:153:TRP:CH2	1:B:188[B]:VAL:HG11	2.53	0.42
1:A:18:VAL:HB	1:A:239[A]:THR:HG23	2.01	0.42
1:B:28[B]:VAL:HG12	1:B:37[B]:VAL:HG23	2.01	0.42
1:B:71[A]:THR:HG23	1:B:102:VAL:HB	2.00	0.42
1:B:78[B]:VAL:HG21	1:B:93:PHE:CE2	2.55	0.42
1:A:155:THR:HG23	1:A:186:VAL:HB	2.02	0.41
1:B:224:THR:O	1:B:241:HIS:HB3	2.21	0.41
1:A:197[A]:THR:CG2	1:A:225:PRO:HB2	2.50	0.41
1:B:184[B]:SER:O	1:B:184[B]:SER:OG	2.38	0.41
1:B:56:THR:O	1:B:73:HIS:HB3	2.20	0.41
1:A:104[B]:VAL:CG1	1:A:110:VAL:HG22	2.45	0.41

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/256 (108%)	267 (97%)	9 (3%)	0	100	100
1	B	277/256 (108%)	268 (97%)	9 (3%)	0	100	100
All	All	553/512 (108%)	535 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/201 (111%)	223 (100%)	0	100	100
1	B	225/201 (112%)	222 (99%)	3 (1%)	69	35
All	All	448/402 (111%)	445 (99%)	3 (1%)	88	61

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	64	SER
1	B	188[A]	VAL
1	B	188[B]	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	226[A]:SER	C	227:GLY	N	1.18

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	251/256 (98%)	0.24	16 (6%) 19 16	4, 9, 24, 32	0
1	B	251/256 (98%)	0.23	11 (4%) 34 32	4, 8, 21, 30	0
All	All	502/512 (98%)	0.24	27 (5%) 25 23	4, 9, 23, 32	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	251	ALA	5.7
1	B	233	ALA	5.3
1	B	142[A]	SER	5.2
1	B	65	ALA	5.2
1	B	232	SER	5.0
1	B	234	GLY	4.4
1	A	168	GLY	4.0
1	A	210	GLY	3.9
1	B	190	SER	3.8
1	A	64	SER	3.7
1	A	65	ALA	3.7
1	A	85	SER	3.6
1	A	82	ALA	3.3
1	A	126	GLY	3.2
1	A	84	GLY	3.2
1	B	64	SER	3.0
1	A	125	ALA	2.9
1	A	171	THR	2.9
1	A	83	ALA	2.9
1	A	170	ASN	2.6
1	A	209	ALA	2.6
1	B	42	GLY	2.5
1	B	250	ALA	2.4
1	A	66	GLY	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	168	GLY	2.2
1	A	22	SER	2.1
1	A	23	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.