



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2025 – 01:47 pm BST

PDB ID : 9FSC / pdb_00009fsc
Title : The structure of ornithine decarboxylase from Leishmania infantum
Authors : Fiorillo, A.; Ilari, A.; Antonelli, L.
Deposited on : 2024-06-20
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	8487 (2.30-2.26)
Clashscore	180529	9437 (2.30-2.26)
Ramachandran outliers	177936	9341 (2.30-2.26)
Sidechain outliers	177891	9342 (2.30-2.26)
RSRZ outliers	164620	8487 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	727	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ornithine decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	0	0
			3471	2214	585	651	21			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP E9AGB5
A	-18	GLY	-	expression tag	UNP E9AGB5
A	-17	SER	-	expression tag	UNP E9AGB5
A	-16	SER	-	expression tag	UNP E9AGB5
A	-15	HIS	-	expression tag	UNP E9AGB5
A	-14	HIS	-	expression tag	UNP E9AGB5
A	-13	HIS	-	expression tag	UNP E9AGB5
A	-12	HIS	-	expression tag	UNP E9AGB5
A	-11	HIS	-	expression tag	UNP E9AGB5
A	-10	HIS	-	expression tag	UNP E9AGB5
A	-9	SER	-	expression tag	UNP E9AGB5
A	-8	SER	-	expression tag	UNP E9AGB5
A	-7	GLY	-	expression tag	UNP E9AGB5
A	-6	LEU	-	expression tag	UNP E9AGB5
A	-5	VAL	-	expression tag	UNP E9AGB5
A	-4	PRO	-	expression tag	UNP E9AGB5
A	-3	ARG	-	expression tag	UNP E9AGB5
A	-2	GLY	-	expression tag	UNP E9AGB5
A	-1	SER	-	expression tag	UNP E9AGB5
A	0	HIS	-	expression tag	UNP E9AGB5

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	104	Total	O	0	0
			104	104		

i

- Molecule 1: ornithine decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	119.22Å 119.22Å 85.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	65.84 – 2.27 65.84 – 2.27	Depositor EDS
% Data completeness (in resolution range)	70.6 (65.84-2.27) 70.6 (65.84-2.27)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.235 , 0.273 0.224 , 0.255	Depositor DCC
R_{free} test set	1119 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	66.4	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.010 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3580	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/3551	1.09	8/4820 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	670	PHE	CA-CB-CG	6.38	120.18	113.80
1	A	356	ASN	CA-CB-CG	5.73	118.33	112.60
1	A	398	VAL	CA-C-N	5.46	127.54	120.44
1	A	398	VAL	C-N-CA	5.46	127.54	120.44
1	A	562	ASP	CA-CB-CG	5.29	117.89	112.60
1	A	327	ASP	CA-CB-CG	5.23	117.83	112.60
1	A	267	VAL	CA-C-N	5.00	127.05	120.60
1	A	267	VAL	C-N-CA	5.00	127.05	120.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3471	0	3376	26	0
2	A	5	0	0	0	0
3	A	104	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3580	0	3376	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:THR:HG21	1:A:371:HIS:CE1	2.20	0.77
1:A:143:VAL:O	1:A:146:LEU:HB2	2.01	0.60
1:A:336:LYS:HE2	1:A:341:LEU:HD21	1.85	0.58
1:A:156:TRP:NE1	1:A:324:ALA:HB2	2.19	0.57
1:A:522:LEU:HB2	1:A:541:PRO:HB2	1.87	0.55
1:A:350:THR:HG22	1:A:371:HIS:CG	2.43	0.53
1:A:406:ARG:NH2	1:A:446:TYR:CD2	2.78	0.51
1:A:430:VAL:HG13	1:A:483:VAL:HG21	1.91	0.51
1:A:484:LEU:HD22	1:A:495:ILE:HD13	1.94	0.50
1:A:371:HIS:HB3	1:A:413:CYS:SG	2.52	0.49
1:A:367:MET:HB3	1:A:370:ALA:HB2	1.95	0.48
1:A:516:ALA:HB3	1:A:547:TYR:HB2	1.94	0.48
1:A:641:LEU:HD11	1:A:644:GLN:HB2	1.97	0.47
1:A:350:THR:HG21	1:A:371:HIS:ND1	2.30	0.47
1:A:350:THR:CG2	1:A:371:HIS:CG	2.99	0.46
1:A:313:GLU:HA	1:A:316:MET:HE3	1.98	0.45
1:A:350:THR:CG2	1:A:371:HIS:ND1	2.80	0.44
1:A:354:VAL:HG23	1:A:359:GLU:HB3	1.99	0.43
1:A:512:MET:HE3	1:A:512:MET:HB2	1.95	0.43
1:A:216:TRP:HB3	1:A:656:LEU:HD21	2.00	0.43
1:A:424:ASP:HB3	1:A:471:THR:HB	1.99	0.43
1:A:307:ASP:OD1	1:A:332:ALA:HB3	2.19	0.43
1:A:331:PHE:HB3	1:A:352:VAL:HG23	2.01	0.41
1:A:169:VAL:HG11	1:A:298:VAL:HG23	2.02	0.41
1:A:350:THR:CG2	1:A:371:HIS:CE1	2.99	0.41
1:A:284:TYR:O	1:A:498:GLU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/727 (59%)	410 (95%)	19 (4%)	1 (0%)	44	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	687	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/592 (63%)	367 (98%)	7 (2%)	52	67

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	LEU
1	A	176	VAL
1	A	389	THR
1	A	522	LEU
1	A	540	GLU
1	A	672	ASN
1	A	686	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	441	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	801	-	4,4,4	0.54	0	6,6,6	0.57	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/727 (60%)	0.83	32 (7%) 23 24	46, 70, 92, 108	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	689	ALA	4.1
1	A	252	ALA	3.9
1	A	209	ILE	3.7
1	A	412	VAL	3.6
1	A	684	SER	3.3
1	A	670	PHE	3.0
1	A	402	LEU	2.8
1	A	470	ASN	2.7
1	A	573	ASP	2.6
1	A	288	LYS	2.6
1	A	429	TYR	2.6
1	A	446	TYR	2.6
1	A	440	PHE	2.5
1	A	351	TYR	2.5
1	A	523	SER	2.5
1	A	366	LEU	2.4
1	A	408	PHE	2.4
1	A	225	GLN	2.4
1	A	405	ALA	2.4
1	A	350	THR	2.3
1	A	180	PRO	2.3
1	A	681	TRP	2.3
1	A	291	PRO	2.3
1	A	354	VAL	2.2
1	A	398	VAL	2.2
1	A	453	LEU	2.2
1	A	448	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	672	ASN	2.1
1	A	415	VAL	2.1
1	A	222	LEU	2.0
1	A	401	LEU	2.0
1	A	541	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PO4	A	801	5/5	0.93	0.10	64,64,64,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.