



wwPDB EM Validation Summary Report ⓘ

May 6, 2025 – 05:43 AM EDT

PDB ID : 8G5H / pdb_00008g5h
EMDB ID : EMD-29743
Title : Native GABA-A receptor from the mouse brain, ortho-alpha1-alpha3-beta2-gamma2 subtype, in complex with GABA, Zolpidem, and endogenous neurosteroid
Authors : Sun, C.; Gouaux, E.
Deposited on : 2023-02-13
Resolution : 2.89 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0rc1
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 210492 | 15764 |
| Ramachandran outliers | 207382 | 16835 |
| Sidechain outliers | 206894 | 16415 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | B | 512 | |
| 1 | E | 512 | |
| 2 | C | 455 | |
| 3 | D | 474 | |
| 4 | J | 223 | |
| 5 | K | 213 | |
| 6 | A | 492 | |
| 7 | F | 5 | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 7 | G | 5 | <div><div></div><div>40%</div><div></div><div>100%</div></div> |
| 7 | H | 5 | <div><div></div><div>80%</div><div></div><div>20%</div></div> |
| 7 | L | 5 | <div><div></div><div>20%</div><div></div><div>40%</div><div></div><div>40%</div><div></div><div>20%</div></div> |
| 7 | M | 5 | <div><div></div><div>80%</div><div></div><div>20%</div></div> |
| 8 | I | 7 | <div><div></div><div>43%</div><div></div><div>57%</div></div> |
| 9 | N | 3 | <div><div></div><div>33%</div><div></div><div>67%</div><div></div><div>33%</div></div> |

2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 15875 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric acid receptor subunit beta-2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 1 | B | 332 | Total | C | N | O | S | 0 | 0 |
| | | | 2718 | 1782 | 435 | 485 | 16 | | |
| 1 | E | 331 | Total | C | N | O | S | 0 | 0 |
| | | | 2711 | 1778 | 434 | 483 | 16 | | |

- Molecule 2 is a protein called Gamma-aminobutyric acid receptor subunit alpha-1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 2 | C | 344 | Total | C | N | O | S | 0 | 0 |
| | | | 2783 | 1800 | 466 | 501 | 16 | | |

- Molecule 3 is a protein called Gamma-aminobutyric acid receptor subunit gamma-2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 3 | D | 318 | Total | C | N | O | S | 0 | 0 |
| | | | 2605 | 1698 | 428 | 465 | 14 | | |

- Molecule 4 is a protein called Heavy Chain of 8E3 Fab.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 4 | J | 117 | Total | C | N | O | S | 0 | 0 |
| | | | 920 | 587 | 152 | 177 | 4 | | |

- Molecule 5 is a protein called Light Chain of 8E3 Fab.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 5 | K | 105 | Total | C | N | O | S | 0 | 0 |
| | | | 796 | 505 | 132 | 155 | 4 | | |

- Molecule 6 is a protein called Gamma-aminobutyric acid receptor subunit alpha-3.


| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 6 | A | 340 | Total | C | N | O | S | 0 | 0 |
| | | | 2748 | 1792 | 451 | 491 | 14 | | |

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|-------|
| 7 | F | 5 | Total | C | N | O | 0 | 0 |
| | | | 61 | 34 | 2 | 25 | | |
| 7 | G | 5 | Total | C | N | O | 0 | 0 |
| | | | 61 | 34 | 2 | 25 | | |
| 7 | H | 5 | Total | C | N | O | 0 | 0 |
| | | | 61 | 34 | 2 | 25 | | |
| 7 | L | 5 | Total | C | N | O | 0 | 0 |
| | | | 61 | 34 | 2 | 25 | | |
| 7 | M | 5 | Total | C | N | O | 0 | 0 |
| | | | 61 | 34 | 2 | 25 | | |

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|-------|
| 8 | I | 7 | Total | C | N | O | 0 | 0 |
| | | | 82 | 46 | 2 | 34 | | |

- Molecule 9 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|-------|
| 9 | N | 3 | Total | C | N | O | 0 | 0 |
| | | | 39 | 22 | 2 | 15 | | |

- Molecule 10 is GAMMA-AMINO-BUTANOIC ACID (CCD ID: ABU) (formula: $C_4H_9NO_2$).




| Mol | Chain | Residues | Atoms | | | | AltConf |
|-----|-------|----------|-------|---|---|---|---------|
| 10 | B | 1 | Total | C | N | O | 0 |
| | | | 7 | 4 | 1 | 2 | |
| 10 | E | 1 | Total | C | N | O | 0 |
| | | | 7 | 4 | 1 | 2 | |

- Molecule 11 is [(2R)-2-octanoyloxy-3-[oxidanyl-[(1R,2R,3S,4R,5R,6S)-2,3,6-tris(oxidanyl)-4,5-diphosphonoxy-cyclohexyl]oxy-phosphoryl]oxy-propyl] octanoate (CCD ID: PIO) (formula: $C_{25}H_{49}O_{19}P_3$).



| Mol | Chain | Residues | Atoms | | | | AltConf |
|-----|-------|----------|-------|----|----|---|---------|
| 11 | C | 1 | Total | C | O | P | 0 |
| | | | 47 | 25 | 19 | 3 | |
| 11 | A | 1 | Total | C | O | P | 0 |
| | | | 47 | 25 | 19 | 3 | |


- Molecule 12 is Zolpidem (CCD ID: R5R) (formula: C₁₉H₂₁N₃O).



| Mol | Chain | Residues | Atoms | | | | AltConf |
|-----|-------|----------|-------|----|---|---|---------|
| 12 | C | 1 | Total | C | N | O | 0 |
| | | | 23 | 19 | 3 | 1 | |


- Molecule 13 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆).



| Mol | Chain | Residues | Atoms | | | | AltConf |
|-----|-------|----------|-------|---|---|---|---------|
| 13 | D | 1 | Total | C | N | O | 0 |
| | | | 14 | 8 | 1 | 5 | |

- Molecule 14 is allopregnanolone (CCD ID: Y4B) (formula: C₂₁H₃₄O₂).




| Mol | Chain | Residues | Atoms | | | AltConf |
|-----|-------|----------|-------|----|---|---------|
| 14 | A | 1 | Total | C | O | 0 |
| | | | 23 | 21 | 2 | |


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-aminobutyric acid receptor subunit beta-2



- Molecule 1: Gamma-aminobutyric acid receptor subunit beta-2



THR
THR
VAL
ASP
LYS
LYS
SER
ALA
ALA
GLU
LEU
VAL
PHE
GLN

- Molecule 5: Light Chain of 8E3 Fab

Chain K:  46% 51%

Y1 Q6 K9 T20 Q37 L47 Y65 T74 T101 I105 LYS ARG ALA ASP ASP ALA ALA PRO THR THR VAL SER SER ILE PHE PRO PRO SER SER GLU GLN LEU THR THR GLY GLY ALA SER VAL VAL CYS PHE LEU LEU ASN ASN PHE THR PRO THR LYS ASP ILE VAL VAL LYS

TRP LYS ILE ASP GLY SER GLU ARG GLN ASP LYS ASP ASP SER THR THR MET SER SER THR LEU THR THR SER THR LYS ASP GLU TYR GLU ARG HIS SER TYR THR CYS GLU ALA THR HIS LYS THR THR SER THR PRO ILE ASN VAL LYS

SER PHE ASN ARG ASN GLU CYS

- Molecule 6: Gamma-aminobutyric acid receptor subunit alpha-3

Chain A:  55% 13% 31%

MET ILE THR THR MET TRP HIS PHE THR TYR VAL THR ARG VAL VAL LEU LEU LEU ILE SER ILE LEU LEU PRO GLY THR THR SER GLN SER THR LEU ARG GLN GLN GLY ASP PHE VAL LYS ASN PRO ILE LYS HIS ALA THR THR ASP ILE PRO ASP SER

THR ASP ASN ILE T37 L43 D48 D51 R55 F56 G57 A61 D68 I69 S78 D79 T80 D81 M82 T85 L110 L114 F125 M136 L142 L143 R144 M155 M167 E168 D169 M172 L179 Q207 Q214 Y215 I227 V236

R245 I262 P267 T261 V262 L264 S265 Q266 L271 N272 R273 V276 T280 V281 T285 L288 T289 P290 T291 T292 T308 D311 V316 C317 Y318 V321 F322 E327 V331 T335 K336 R337 G343 LYS LYS VAL PRO GLU ALA LEU GLU MET LYS

LYS LYS THR PRO ALA PRO THR LYS LYS ASN THR PHE ASN ILE VAL GLY THR THR TYR PRO ILE LEU LEU ALA ASP THR GLY PHE SER THR ILE ALA ALA PRO SER ALA SER SER THR PRO THR THR ALA ILE ALA SER PRO VAL PRO LYS ALA THR TYR VAL GLN MET ASP

SER PRO ALA GLU THR K419 T420 Y421 R422 S423 V424 S425 I430 S431 I434 L438 F442 V451 ASN ARG GLU GLU SER SER ALA ILE LYS GLY MET ILE ARG LYS GLN

- Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  60% 20% 20%


MAG1
MAG2
BNA3
MAN4
MAN5

- Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  40% 100%

MAG1
MAG2
BNA3
MAN4
MAN5


- Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  80% 20%




- Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  20% 40% 40% 20%



- Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  80% 20%



- Molecule 8: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  43% 57%



- Molecule 9: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  33% 67% 33%



4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, C1 | Depositor |
| Number of particles used | 91747 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 50.00 | Depositor |
| Minimum defocus (nm) | 800 | Depositor |
| Maximum defocus (nm) | 2100 | Depositor |
| Magnification | Not provided | |
| Image detector | GATAN K3 BIOQUANTUM (6k x 4k) | Depositor |
| Maximum map value | 0.631 | Depositor |
| Minimum map value | -0.212 | Depositor |
| Average map value | 0.003 | Depositor |
| Map value standard deviation | 0.017 | Depositor |
| Recommended contour level | 0.075 | Depositor |
| Map size (Å) | 297.36, 297.36, 297.36 | wwPDB |
| Map dimensions | 360, 360, 360 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 0.826, 0.826, 0.826 | Depositor |

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, Y4B, ABU, R5R, BMA, MAN, PIO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-------------|-------------|----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | B | 0.15 | 0/2788 | 0.32 | 0/3795 |
| 1 | E | 0.16 | 0/2781 | 0.33 | 2/3785 (0.1%) |
| 2 | C | 0.13 | 0/2854 | 0.29 | 2/3878 (0.1%) |
| 3 | D | 0.14 | 0/2676 | 0.30 | 0/3646 |
| 4 | J | 0.15 | 0/946 | 0.28 | 0/1284 |
| 5 | K | 0.15 | 0/815 | 0.31 | 0/1108 |
| 6 | A | 0.18 | 0/2822 | 0.35 | 2/3842 (0.1%) |
| All | All | 0.15 | 0/15682 | 0.32 | 6/21338 (0.0%) |

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^{\circ}$) | Ideal($^{\circ}$) |
|-----|-------|-----|------|--------|------|------------------------|---------------------|
| 6 | A | 136 | MET | CA-C-N | 5.36 | 131.34 | 121.70 |
| 6 | A | 136 | MET | C-N-CA | 5.36 | 131.34 | 121.70 |
| 1 | E | 109 | VAL | CA-C-N | 5.18 | 131.03 | 121.70 |
| 1 | E | 109 | VAL | C-N-CA | 5.18 | 131.03 | 121.70 |
| 2 | C | 111 | MET | CA-C-N | 5.15 | 130.97 | 121.70 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | B | 2718 | 0 | 2721 | 46 | 0 |
| 1 | E | 2711 | 0 | 2714 | 30 | 0 |
| 2 | C | 2783 | 0 | 2763 | 46 | 0 |
| 3 | D | 2605 | 0 | 2581 | 34 | 0 |
| 4 | J | 920 | 0 | 863 | 9 | 0 |
| 5 | K | 796 | 0 | 756 | 5 | 0 |
| 6 | A | 2748 | 0 | 2744 | 36 | 0 |
| 7 | F | 61 | 0 | 52 | 1 | 0 |
| 7 | G | 61 | 0 | 52 | 1 | 0 |
| 7 | H | 61 | 0 | 52 | 1 | 0 |
| 7 | L | 61 | 0 | 52 | 1 | 0 |
| 7 | M | 61 | 0 | 52 | 2 | 0 |
| 8 | I | 82 | 0 | 70 | 0 | 0 |
| 9 | N | 39 | 0 | 34 | 0 | 0 |
| 10 | B | 7 | 0 | 0 | 1 | 0 |
| 10 | E | 7 | 0 | 0 | 0 | 0 |
| 11 | A | 47 | 0 | 44 | 1 | 0 |
| 11 | C | 47 | 0 | 44 | 4 | 0 |
| 12 | C | 23 | 0 | 0 | 0 | 0 |
| 13 | D | 14 | 0 | 13 | 0 | 0 |
| 14 | A | 23 | 0 | 0 | 0 | 0 |
| All | All | 15875 | 0 | 15607 | 196 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 196 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|-----------------|--------------------------|-------------------|
| 1:B:68:ARG:NH1 | 1:B:121:ASP:O | 2.18 | 0.77 |
| 2:C:204:SER:OG | 2:C:207:GLY:O | 2.03 | 0.77 |
| 6:A:257:PRO:O | 6:A:261:THR:OG1 | 2.03 | 0.77 |
| 2:C:266:THR:HG1 | 2:C:293:TYR:HH | 1.27 | 0.76 |
| 1:E:269:ARG:NH1 | 1:E:282:ASP:OD2 | 2.22 | 0.72 |

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|---------|----------|-------------|-----|
| 1 | B | 328/512 (64%) | 315 (96%) | 13 (4%) | 0 | 100 | 100 |
| 1 | E | 327/512 (64%) | 316 (97%) | 11 (3%) | 0 | 100 | 100 |
| 2 | C | 340/455 (75%) | 335 (98%) | 5 (2%) | 0 | 100 | 100 |
| 3 | D | 314/474 (66%) | 301 (96%) | 13 (4%) | 0 | 100 | 100 |
| 4 | J | 115/223 (52%) | 112 (97%) | 3 (3%) | 0 | 100 | 100 |
| 5 | K | 103/213 (48%) | 95 (92%) | 8 (8%) | 0 | 100 | 100 |
| 6 | A | 336/492 (68%) | 324 (96%) | 12 (4%) | 0 | 100 | 100 |
| All | All | 1863/2881 (65%) | 1798 (96%) | 65 (4%) | 0 | 100 | 100 |

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|-----|
| 1 | B | 302/457 (66%) | 291 (96%) | 11 (4%) | 30 | 65 |
| 1 | E | 301/457 (66%) | 294 (98%) | 7 (2%) | 45 | 77 |
| 2 | C | 306/404 (76%) | 301 (98%) | 5 (2%) | 58 | 84 |
| 3 | D | 292/436 (67%) | 286 (98%) | 6 (2%) | 48 | 78 |
| 4 | J | 98/195 (50%) | 98 (100%) | 0 | 100 | 100 |
| 5 | K | 83/188 (44%) | 83 (100%) | 0 | 100 | 100 |
| 6 | A | 304/435 (70%) | 291 (96%) | 13 (4%) | 25 | 57 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|-----------------|------------|----------|-------------|
| All | All | 1686/2572 (66%) | 1644 (98%) | 42 (2%) | 43 75 |

5 of 42 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | E | 301 | LEU |
| 6 | A | 263 | ILE |
| 6 | A | 37 | THR |
| 6 | A | 85 | THR |
| 6 | A | 271 | LEU |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 5 | K | 37 | GLN |
| 5 | K | 87 | HIS |
| 6 | A | 266 | GLN |
| 3 | D | 258 | ASN |
| 3 | D | 239 | GLN |

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

35 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 7 | NAG | F | 1 | 7,2 | 14,14,15 | 0.44 | 0 | 17,19,21 | 0.49 | 0 |
| 7 | NAG | F | 2 | 7 | 14,14,15 | 0.21 | 0 | 17,19,21 | 0.50 | 0 |
| 7 | BMA | F | 3 | 7 | 11,11,12 | 0.65 | 0 | 15,15,17 | 0.74 | 0 |
| 7 | MAN | F | 4 | 7 | 11,11,12 | 0.73 | 1 (9%) | 15,15,17 | 1.14 | 2 (13%) |
| 7 | MAN | F | 5 | 7 | 11,11,12 | 0.60 | 0 | 15,15,17 | 0.88 | 2 (13%) |
| 7 | NAG | G | 1 | 1,7 | 14,14,15 | 0.65 | 0 | 17,19,21 | 0.79 | 1 (5%) |
| 7 | NAG | G | 2 | 7 | 14,14,15 | 0.23 | 0 | 17,19,21 | 0.46 | 0 |
| 7 | BMA | G | 3 | 7 | 11,11,12 | 0.50 | 0 | 15,15,17 | 0.77 | 0 |
| 7 | MAN | G | 4 | 7 | 11,11,12 | 0.68 | 0 | 15,15,17 | 1.17 | 2 (13%) |
| 7 | MAN | G | 5 | 7 | 11,11,12 | 0.62 | 0 | 15,15,17 | 0.90 | 1 (6%) |
| 7 | NAG | H | 1 | 1,7 | 14,14,15 | 0.93 | 1 (7%) | 17,19,21 | 0.79 | 1 (5%) |
| 7 | NAG | H | 2 | 7 | 14,14,15 | 0.39 | 0 | 17,19,21 | 1.34 | 2 (11%) |
| 7 | BMA | H | 3 | 7 | 11,11,12 | 1.04 | 1 (9%) | 15,15,17 | 1.49 | 2 (13%) |
| 7 | MAN | H | 4 | 7 | 11,11,12 | 0.78 | 1 (9%) | 15,15,17 | 1.17 | 2 (13%) |
| 7 | MAN | H | 5 | 7 | 11,11,12 | 0.77 | 1 (9%) | 15,15,17 | 1.02 | 2 (13%) |
| 8 | NAG | I | 1 | 6,8 | 14,14,15 | 0.16 | 0 | 17,19,21 | 0.58 | 0 |
| 8 | NAG | I | 2 | 8 | 14,14,15 | 0.24 | 0 | 17,19,21 | 0.68 | 1 (5%) |
| 8 | BMA | I | 3 | 8 | 11,11,12 | 0.70 | 0 | 15,15,17 | 0.82 | 0 |
| 8 | MAN | I | 4 | 8 | 11,11,12 | 0.90 | 1 (9%) | 15,15,17 | 1.21 | 2 (13%) |
| 8 | MAN | I | 5 | 8 | 11,11,12 | 0.64 | 0 | 15,15,17 | 1.23 | 2 (13%) |
| 8 | MAN | I | 6 | 8 | 11,11,12 | 0.61 | 0 | 15,15,17 | 0.97 | 2 (13%) |
| 8 | FUC | I | 7 | 8 | 10,10,11 | 0.64 | 0 | 14,14,16 | 0.81 | 0 |
| 7 | NAG | L | 1 | 1,7 | 14,14,15 | 0.46 | 0 | 17,19,21 | 0.53 | 0 |
| 7 | NAG | L | 2 | 7 | 14,14,15 | 0.41 | 0 | 17,19,21 | 1.26 | 1 (5%) |
| 7 | BMA | L | 3 | 7 | 11,11,12 | 0.41 | 0 | 15,15,17 | 0.72 | 0 |
| 7 | MAN | L | 4 | 7 | 11,11,12 | 0.70 | 0 | 15,15,17 | 0.93 | 2 (13%) |
| 7 | MAN | L | 5 | 7 | 11,11,12 | 0.70 | 0 | 15,15,17 | 0.95 | 1 (6%) |
| 7 | NAG | M | 1 | 1,7 | 14,14,15 | 0.44 | 0 | 17,19,21 | 0.43 | 0 |
| 7 | NAG | M | 2 | 7 | 14,14,15 | 0.31 | 0 | 17,19,21 | 1.43 | 2 (11%) |
| 7 | BMA | M | 3 | 7 | 11,11,12 | 0.84 | 1 (9%) | 15,15,17 | 1.21 | 2 (13%) |
| 7 | MAN | M | 4 | 7 | 11,11,12 | 0.79 | 1 (9%) | 15,15,17 | 1.20 | 2 (13%) |
| 7 | MAN | M | 5 | 7 | 11,11,12 | 0.84 | 1 (9%) | 15,15,17 | 1.21 | 3 (20%) |
| 9 | NAG | N | 1 | 9,3 | 14,14,15 | 0.26 | 0 | 17,19,21 | 0.73 | 1 (5%) |
| 9 | NAG | N | 2 | 9 | 14,14,15 | 0.22 | 0 | 17,19,21 | 0.49 | 0 |
| 9 | BMA | N | 3 | 9 | 11,11,12 | 0.63 | 0 | 15,15,17 | 0.73 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 7 | NAG | F | 1 | 7,2 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | F | 2 | 7 | - | 2/6/23/26 | 0/1/1/1 |
| 7 | BMA | F | 3 | 7 | - | 0/2/19/22 | 0/1/1/1 |
| 7 | MAN | F | 4 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 7 | MAN | F | 5 | 7 | - | 0/2/19/22 | 0/1/1/1 |
| 7 | NAG | G | 1 | 1,7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | G | 2 | 7 | - | 1/6/23/26 | 0/1/1/1 |
| 7 | BMA | G | 3 | 7 | - | 0/2/19/22 | 0/1/1/1 |
| 7 | MAN | G | 4 | 7 | - | 1/2/19/22 | 1/1/1/1 |
| 7 | MAN | G | 5 | 7 | - | 1/2/19/22 | 0/1/1/1 |
| 7 | NAG | H | 1 | 1,7 | - | 2/6/23/26 | 0/1/1/1 |
| 7 | NAG | H | 2 | 7 | - | 6/6/23/26 | 0/1/1/1 |
| 7 | BMA | H | 3 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 7 | MAN | H | 4 | 7 | - | 0/2/19/22 | 0/1/1/1 |
| 7 | MAN | H | 5 | 7 | - | 2/2/19/22 | 1/1/1/1 |
| 8 | NAG | I | 1 | 6,8 | - | 0/6/23/26 | 0/1/1/1 |
| 8 | NAG | I | 2 | 8 | - | 4/6/23/26 | 0/1/1/1 |
| 8 | BMA | I | 3 | 8 | - | 2/2/19/22 | 0/1/1/1 |
| 8 | MAN | I | 4 | 8 | - | 1/2/19/22 | 1/1/1/1 |
| 8 | MAN | I | 5 | 8 | - | 0/2/19/22 | 1/1/1/1 |
| 8 | MAN | I | 6 | 8 | - | 1/2/19/22 | 0/1/1/1 |
| 8 | FUC | I | 7 | 8 | - | - | 0/1/1/1 |
| 7 | NAG | L | 1 | 1,7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | L | 2 | 7 | - | 6/6/23/26 | 0/1/1/1 |
| 7 | BMA | L | 3 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 7 | MAN | L | 4 | 7 | - | 0/2/19/22 | 1/1/1/1 |
| 7 | MAN | L | 5 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 7 | NAG | M | 1 | 1,7 | - | 0/6/23/26 | 0/1/1/1 |
| 7 | NAG | M | 2 | 7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | BMA | M | 3 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 7 | MAN | M | 4 | 7 | - | 0/2/19/22 | 0/1/1/1 |
| 7 | MAN | M | 5 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 9 | NAG | N | 1 | 9,3 | - | 4/6/23/26 | 0/1/1/1 |
| 9 | NAG | N | 2 | 9 | - | 0/6/23/26 | 0/1/1/1 |
| 9 | BMA | N | 3 | 9 | - | 0/2/19/22 | 0/1/1/1 |

The worst 5 of 9 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 7 | H | 1 | NAG | O5-C1 | -3.16 | 1.38 | 1.43 |
| 7 | H | 4 | MAN | C1-C2 | 2.46 | 1.58 | 1.52 |
| 7 | M | 5 | MAN | C1-C2 | 2.43 | 1.58 | 1.52 |
| 7 | M | 4 | MAN | C1-C2 | 2.42 | 1.58 | 1.52 |
| 8 | I | 4 | MAN | C1-C2 | 2.40 | 1.57 | 1.52 |

The worst 5 of 38 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|------|-------------|----------|
| 7 | H | 3 | BMA | C1-O5-C5 | 4.64 | 118.41 | 112.19 |
| 7 | M | 2 | NAG | C2-N2-C7 | 4.37 | 128.76 | 122.90 |
| 7 | H | 2 | NAG | C2-N2-C7 | 4.26 | 128.61 | 122.90 |
| 7 | L | 2 | NAG | C2-N2-C7 | 4.25 | 128.60 | 122.90 |
| 8 | I | 5 | MAN | C1-O5-C5 | 3.84 | 117.33 | 112.19 |

There are no chirality outliers.

5 of 61 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 7 | L | 3 | BMA | O5-C5-C6-O6 |
| 7 | F | 4 | MAN | O5-C5-C6-O6 |
| 7 | L | 5 | MAN | O5-C5-C6-O6 |
| 7 | H | 3 | BMA | C4-C5-C6-O6 |
| 7 | F | 1 | NAG | O5-C5-C6-O6 |

All (5) ring outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------------|
| 7 | L | 4 | MAN | C1-C2-C3-C4-C5-O5 |
| 8 | I | 5 | MAN | C1-C2-C3-C4-C5-O5 |
| 7 | G | 4 | MAN | C1-C2-C3-C4-C5-O5 |
| 7 | H | 5 | MAN | C1-C2-C3-C4-C5-O5 |
| 8 | I | 4 | MAN | C1-C2-C3-C4-C5-O5 |

7 monomers are involved in 6 short contacts:


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 7 | G | 2 | NAG | 1 | 0 |
| 7 | M | 1 | NAG | 1 | 0 |
| 7 | H | 2 | NAG | 1 | 0 |
| 7 | L | 2 | NAG | 1 | 0 |
| 7 | F | 4 | MAN | 1 | 0 |


Continued on next page...


Continued from previous page...


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 7 | G | 3 | BMA | 1 | 0 |
| 7 | M | 2 | NAG | 1 | 0 |


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 12 | R5R | C | 602 | - | 22,25,25 | 2.56 | 5 (22%) | 21,36,36 | 1.65 | 4 (19%) |
| 14 | Y4B | A | 502 | - | 26,26,26 | 1.35 | 3 (11%) | 42,42,42 | 2.38 | 19 (45%) |
| 11 | PIO | C | 601 | - | 47,47,47 | 1.40 | 10 (21%) | 62,65,65 | 1.24 | 7 (11%) |
| 11 | PIO | A | 501 | - | 47,47,47 | 1.41 | 11 (23%) | 62,65,65 | 1.30 | 9 (14%) |
| 13 | NAG | D | 501 | 3 | 14,14,15 | 0.26 | 0 | 17,19,21 | 0.74 | 1 (5%) |
| 10 | ABU | E | 501 | - | 6,6,6 | 0.87 | 0 | 6,6,6 | 1.29 | 0 |
| 10 | ABU | B | 501 | - | 6,6,6 | 0.92 | 0 | 6,6,6 | 1.33 | 1 (16%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-------------|---------|
| 12 | R5R | C | 602 | - | - | 3/12/12/12 | 0/3/3/3 |
| 14 | Y4B | A | 502 | - | - | 4/4/62/62 | 0/4/4/4 |
| 11 | PIO | C | 601 | - | - | 17/44/68/68 | 0/1/1/1 |
| 11 | PIO | A | 501 | - | - | 11/44/68/68 | 0/1/1/1 |
| 13 | NAG | D | 501 | 3 | - | 2/6/23/26 | 0/1/1/1 |
| 10 | ABU | E | 501 | - | - | 3/4/4/4 | - |
| 10 | ABU | B | 501 | - | - | 2/4/4/4 | - |

The worst 5 of 29 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 12 | C | 602 | R5R | C19-N21 | 10.22 | 1.46 | 1.34 |
| 14 | A | 502 | Y4B | C04-C05 | 3.69 | 1.60 | 1.53 |
| 12 | C | 602 | R5R | C05-C08 | 3.29 | 1.52 | 1.49 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 11 | A | 501 | PIO | P4-O4 | 3.26 | 1.65 | 1.59 |
| 11 | C | 601 | PIO | P4-O4 | 3.26 | 1.65 | 1.59 |

The worst 5 of 41 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|------|-------------|----------|
| 12 | C | 602 | R5R | C18-C19-N21 | 5.19 | 124.39 | 117.39 |
| 14 | A | 502 | Y4B | C10-C06-C12 | 4.63 | 121.67 | 116.11 |
| 14 | A | 502 | Y4B | C12-C06-C05 | 4.25 | 104.16 | 99.72 |
| 14 | A | 502 | Y4B | C23-C22-C12 | 4.15 | 124.83 | 117.65 |
| 11 | C | 601 | PIO | O2C-C1A-C2A | 4.09 | 120.32 | 111.48 |

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-----------------|
| 10 | B | 501 | ABU | CD-CB-CG-C |
| 11 | C | 601 | PIO | C1C-O13-P1-O1 |
| 11 | C | 601 | PIO | C1C-O13-P1-O11 |
| 11 | C | 601 | PIO | C1C-O13-P1-O12 |
| 11 | C | 601 | PIO | C2A-C1A-O2C-C2C |


There are no ring outliers.

3 monomers are involved in 6 short contacts:


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 11 | C | 601 | PIO | 4 | 0 |
| 11 | A | 501 | PIO | 1 | 0 |
| 10 | B | 501 | ABU | 1 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand R5R C 602




Bond lengths



Bond angles




Torsions




Rings


Ligand Y4B A 502




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-29743. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 187




Y Index: 204




Z Index: 203


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.075. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_29743_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 217 nm³; this corresponds to an approximate mass of 196 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.346 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.346 \AA^{-1}

8.2 Resolution estimates [i](#)


| Resolution estimate (Å) | Estimation criterion (FSC cut-off) | | |
|---------------------------|------------------------------------|------|----------|
| | 0.143 | 0.5 | Half-bit |
| Reported by author | 2.89 | - | - |
| Author-provided FSC curve | 2.89 | 3.43 | 2.94 |
| Unmasked-calculated* | 4.03 | 8.01 | 4.13 |

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.03 differs from the reported value 2.89 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-29743 and PDB model 8G5H. Per-residue inclusion information can be found in [section 3](#) on [page 9](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.075 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.075).































9.4 Atom inclusion ⓘ



At the recommended contour level, 97% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.075) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|---|---|
| All |  0.9150 |  0.4770 |
| A |  0.9310 |  0.4820 |
| B |  0.9360 |  0.4900 |
| C |  0.9370 |  0.5000 |
| D |  0.8180 |  0.4130 |
| E |  0.9370 |  0.4870 |
| F |  0.8690 |  0.4700 |
| G |  0.5740 |  0.3700 |
| H |  0.8690 |  0.3870 |
| I |  0.9020 |  0.4610 |
| J |  0.9650 |  0.5050 |
| K |  0.9680 |  0.5170 |
| L |  0.6560 |  0.3530 |
| M |  0.8850 |  0.4120 |
| N |  0.5900 |  0.4470 |

