



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2024 – 08:54 PM EST

PDB ID : 4G68  
Title : Biochemical and structural insights into xylan utilization by the thermophilic bacterium *Caldanaerobius polysaccharolyticus*  
Authors : Agarwal, V.; Nair, S.K.  
Deposited on : 2012-07-18  
Resolution : 1.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	
2	B	432	
3	C	432	
4	D	3	
4	E	3	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	Se	0	2	0
			3058	1957	494	597	4	6			

- Molecule 2 is a protein called ABC transporter.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	392	Total	C	N	O	S	Se	0	0	0
			3050	1951	494	596	4	5			

- Molecule 3 is a protein called ABC transporter.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	392	Total	C	N	O	S	Se	0	0	0
			3050	1951	494	596	8	1			

- Molecule 4 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-alpha-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	3	Total	C	O	0	0	0
			28	15	13			
4	E	3	Total	C	O	0	0	0
			28	15	13			

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	364	Total 364	O 364	0	0
5	B	363	Total 363	O 363	0	0
5	C	269	Total 269	O 269	0	0



- Molecule 1: ABC transporter

[illegible]

- Molecule 2: ABC transporter


Q138	Q146 GLY	K147	HIS	S162	Y193	D194	M195	L198	N218	L244	A251	Q255	E284	A285	V286	R287	D297	P298	T299	E300	K316	Y317	K318	D319	E320	R323	K330	D336	Y337	L343	W346	N350	V356	Q362	I363	K364	K372	E406	G412		
MET	ALA	HIS	HIS	HIS	HIS	HIS	VAL	ASP	ASP	ASP	LYS	MET	CYS	SER	SER	SER	ASN	ASN	SER	SER	SER	LYS	THR	SER	SER	SER	LYS	LYS	I25	F29	L32	K41	E60	S61	V62	T63	E64	N79	K101	T126	I131

- Molecule 3: ABC transporter

M195	P82	MET
L198	G90	ALA
R199	G91	HIS
R207	F92	HIS
K213	K101	HIS
F216	V102	HIS
D217	L103	VAL
T222	L109	ASP
A225	N110	ASP
Q226	K114	LYS
K227	D115	MET
L228	M124	CYS
G240	M128	SER
L244	I131	ASN
F253	Q138	LEU
M254	Q139	SER
M260	L143	THR
Y261	I144	ASN
V279	I145	ASN
E284	M154	SER
A285	V155	SER
V286	K156	SER
R287	V157	SER
F288	F161	SER
P289	S162	LYS
I291	D166	LYS
G294	A167	THR
K295	I168	LYS
P298	K169	LYS
E300	K172	LYS
G303	P178	LYS
K316	A180	LYS
V322	L181	LYS
K326	G182	LYS
Y327	E183	LYS
A329	M187	LYS
K330	M190	LYS
Q331	W191	LYS
R332	Y192	LYS
	Y193	LYS
	Y194	LYS
	D66	LYS
	K69	LYS
	A78	LYS
	N79	LYS
	E81	LYS



- Molecule 4: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-alpha-D-xylopyranose



- Molecule 4: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-alpha-D-xylopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.33Å 150.86Å 150.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.80 25.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.00-1.80) 99.7 (25.00-1.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.42 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.5.0056	Depositor
R, $R_{free}$	0.230 , 0.261 0.215 , 0.257	Depositor DCC
$R_{free}$ test set	6314 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.8	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 20.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.478 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10210	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XYS, XYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.11	4/3125 (0.1%)	0.99	6/4224 (0.1%)
2	B	1.08	5/3112 (0.2%)	0.97	9/4209 (0.2%)
3	C	0.91	5/3116 (0.2%)	0.87	3/4221 (0.1%)
All	All	1.04	14/9353 (0.1%)	0.94	18/12654 (0.1%)

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320	GLU	CD-OE1	6.94	1.33	1.25
2	B	406	GLU	CD-OE2	-6.76	1.18	1.25
2	B	346	TRP	CD2-CE2	6.33	1.49	1.41
2	B	320	GLU	CD-OE1	6.33	1.32	1.25
2	B	64	GLU	CD-OE1	6.33	1.32	1.25

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	195	MET	CG-SD-CE	-11.49	81.82	100.20
1	A	323	ARG	NE-CZ-NH1	8.62	124.61	120.30
1	A	323	ARG	NE-CZ-NH2	-7.25	116.67	120.30
3	C	207	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	A	195	MSE	CG-SE-CE	-6.64	84.30	98.90

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3058	0	2995	32	1
2	B	3050	0	2981	34	1
3	C	3050	0	2981	165	2
4	D	28	0	9	0	0
4	E	28	0	9	0	0
5	A	364	0	0	14	0
5	B	363	0	0	14	0
5	C	269	0	0	126	0
All	All	10210	0	8975	229	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:218:ASN:HB3	5:B:1245:HOH:O	1.21	1.29
3:C:69:LYS:HD2	5:C:701:HOH:O	1.25	1.27
3:C:363:ILE:HG22	5:C:764:HOH:O	1.13	1.27
3:C:198:LEU:HG	5:C:765:HOH:O	1.28	1.27
3:C:279:VAL:HG21	5:C:584:HOH:O	1.26	1.26

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:162:SER:O	3:C:154:ASN:ND2[1_455]	2.02	0.18
1:A:347:LYS:NZ	3:C:295:LYS:NZ[1_455]	2.13	0.07

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/456 (86%)	386 (98%)	5 (1%)	1 (0%)	37	25
2	B	390/432 (90%)	382 (98%)	8 (2%)	0	100	100
3	C	390/432 (90%)	379 (97%)	10 (3%)	1 (0%)	37	25
All	All	1172/1320 (89%)	1147 (98%)	23 (2%)	2 (0%)	44	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	154	ASN
1	A	65	ASN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/375 (86%)	317 (99%)	4 (1%)	67	62
2	B	319/352 (91%)	315 (99%)	4 (1%)	65	59
3	C	319/356 (90%)	312 (98%)	7 (2%)	47	36
All	All	959/1083 (89%)	944 (98%)	15 (2%)	58	50

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	372	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	C	350	ASN
3	C	25	ILE
3	C	372	LYS
3	C	198	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	229	GLN
2	B	362	GLN
2	B	353	GLN
3	C	57	GLN
1	A	350	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	XYS	D	1	4	10,10,10	2.01	3 (30%)	14,14,14	2.63	8 (57%)
4	XYP	D	2	4	9,9,10	0.97	0	10,12,14	2.09	4 (40%)
4	XYP	D	3	4	9,9,10	1.00	0	10,12,14	1.31	1 (10%)
4	XYS	E	1	4	10,10,10	1.93	3 (30%)	14,14,14	2.45	7 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	XYP	E	2	4	9,9,10	1.19	1 (11%)	10,12,14	2.06	2 (20%)
4	XYP	E	3	4	9,9,10	0.92	0	10,12,14	1.03	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XYS	D	1	4	-	-	0/1/1/1
4	XYP	D	2	4	-	-	0/1/1/1
4	XYP	D	3	4	-	-	0/1/1/1
4	XYS	E	1	4	-	-	0/1/1/1
4	XYP	E	2	4	-	-	0/1/1/1
4	XYP	E	3	4	-	-	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	XYS	O5-C5	-4.87	1.35	1.43
4	E	1	XYS	O5-C5	-4.30	1.36	1.43
4	E	1	XYS	O5-C1	-2.69	1.38	1.43
4	E	2	XYP	C2-C3	2.46	1.56	1.52
4	E	1	XYS	O2-C2	-2.32	1.37	1.43

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	XYS	C5-O5-C1	6.29	125.54	112.46
4	E	1	XYS	C5-O5-C1	5.63	124.17	112.46
4	E	2	XYP	C4-C3-C2	-4.77	105.26	110.92
4	D	2	XYP	C4-C3-C2	-4.40	105.70	110.92
4	D	1	XYS	O5-C5-C4	3.44	118.99	110.79


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	386/456 (84%)	-1.76	0 100 100	4, 9, 19, 36	2 (0%)
2	B	387/432 (89%)	-1.77	0 100 100	4, 9, 18, 35	0
3	C	391/432 (90%)	-1.45	0 100 100	15, 28, 41, 54	0
All	All	1164/1320 (88%)	-1.66	0 100 100	4, 13, 36, 54	2 (0%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	XYS	D	1	10/10	0.99	0.02	8,10,15,17	0
4	XYP	D	2	9/10	1.00	0.01	5,5,6,6	0
4	XYP	D	3	9/10	1.00	0.01	4,4,4,4	0
4	XYS	E	1	10/10	1.00	0.02	9,11,14,19	0
4	XYP	E	2	9/10	1.00	0.01	4,5,5,5	0
4	XYP	E	3	9/10	1.00	0.01	3,3,4,4	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.