



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 12:38 AM EDT

PDB ID : 3G8R  
Title : Crystal structure of putative spore coat polysaccharide biosynthesis protein E from *Chromobacterium violaceum* ATCC 12472  
Authors : Malashkevich, V.N.; Toro, R.; Morano, C.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2009-02-12  
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	 63% 26% 6% . .
1	B	350	 67% 22% 6% . .

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable spore coat polysaccharide biosynthesis protein E.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	337	Total	C	N	O	S	Se	0	0	0
			2646	1671	464	497	4	10			
1	B	336	Total	C	N	O	S	Se	0	1	0
			2647	1671	466	496	4	10			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP Q7NR94
A	2	SER	-	expression tag	UNP Q7NR94
A	3	LEU	-	expression tag	UNP Q7NR94
A	343	GLU	-	expression tag	UNP Q7NR94
A	344	GLY	-	expression tag	UNP Q7NR94
A	345	HIS	-	expression tag	UNP Q7NR94
A	346	HIS	-	expression tag	UNP Q7NR94
A	347	HIS	-	expression tag	UNP Q7NR94
A	348	HIS	-	expression tag	UNP Q7NR94
A	349	HIS	-	expression tag	UNP Q7NR94
A	350	HIS	-	expression tag	UNP Q7NR94
B	1	MSE	-	expression tag	UNP Q7NR94
B	2	SER	-	expression tag	UNP Q7NR94
B	3	LEU	-	expression tag	UNP Q7NR94
B	343	GLU	-	expression tag	UNP Q7NR94
B	344	GLY	-	expression tag	UNP Q7NR94
B	345	HIS	-	expression tag	UNP Q7NR94
B	346	HIS	-	expression tag	UNP Q7NR94
B	347	HIS	-	expression tag	UNP Q7NR94
B	348	HIS	-	expression tag	UNP Q7NR94
B	349	HIS	-	expression tag	UNP Q7NR94
B	350	HIS	-	expression tag	UNP Q7NR94

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	125	Total 125	O 125	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable spore coat polysaccharide biosynthesis protein E



- Molecule 1: Probable spore coat polysaccharide biosynthesis protein E



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.67Å 107.67Å 164.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.49 19.88 – 2.49	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.49) 90.4 (19.88-2.49)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, $R_{free}$	0.208 , 0.278 0.212 , 0.272	Depositor DCC
$R_{free}$ test set	1575 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.1	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 30.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5420	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.83	1/2694 (0.0%)	0.85	4/3633 (0.1%)
1	B	0.86	3/2698 (0.1%)	0.86	4/3637 (0.1%)
All	All	0.84	4/5392 (0.1%)	0.86	8/7270 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
All	All	0	6

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	258	MSE	CG-SE	-5.61	1.76	1.95
1	B	212	MSE	CG-SE	-5.33	1.77	1.95
1	A	212	MSE	CG-SE	-5.21	1.77	1.95
1	B	17	MSE	CG-SE	-5.04	1.78	1.95

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ASP	C-N-CD	6.37	141.78	128.40
1	B	63	ASP	N-CA-CB	-5.62	100.48	110.60
1	A	308	GLY	N-CA-C	-5.54	99.25	113.10
1	A	205	PRO	N-CA-C	-5.31	98.30	112.10
1	B	50	LEU	CA-CB-CG	5.28	127.45	115.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	160	ARG	Peptide
1	A	203	GLU	Peptide
1	A	204	ASP	Peptide
1	A	307	GLU	Peptide
1	B	203	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2646	0	2599	91	0
1	B	2647	0	2603	92	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	125	0	0	6	0
All	All	5420	0	5202	180	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 180 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:THR:HG22	1:B:184:ARG:NH2	1.62	1.14
1:A:189:ARG:HH11	1:A:189:ARG:HG3	1.05	1.08
1:B:142:THR:HG22	1:B:184:ARG:HH22	1.14	1.06
1:A:208:MSE:HE1	1:A:247:ARG:C	1.77	1.04
1:B:208:MSE:HE1	1:B:247:ARG:C	1.89	0.92

There are no symmetry-related clashes.



### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/350 (96%)	309 (92%)	20 (6%)	6 (2%)	8	14
1	B	335/350 (96%)	315 (94%)	18 (5%)	2 (1%)	25	43
All	All	670/700 (96%)	624 (93%)	38 (6%)	8 (1%)	13	24

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	ASP
1	B	178	ASP
1	A	206	ASP
1	A	298	ASN
1	A	208	MSE

#### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/279 (100%)	243 (87%)	35 (13%)	4	8
1	B	278/279 (100%)	242 (87%)	36 (13%)	4	8
All	All	556/558 (100%)	485 (87%)	71 (13%)	4	8

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	230	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	232	GLN
1	B	306	VAL
1	A	231	ASP
1	A	230	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	232	GLN
1	B	236	ASN
1	B	313	ASN
1	A	236	ASN
1	A	244	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	327/350 (93%)	-0.15	4 (1%) 79 80	27, 44, 63, 86	0
1	B	326/350 (93%)	-0.10	5 (1%) 73 75	28, 42, 62, 79	0
All	All	653/700 (93%)	-0.13	9 (1%) 75 77	27, 43, 63, 86	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	232	GLN	6.4
1	B	266	ALA	3.7
1	A	263	GLU	2.9
1	B	205	PRO	2.8
1	B	297	ASP	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	501	1/1	0.84	0.05	88,88,88,88	0
2	ZN	B	501	1/1	0.93	0.09	81,81,81,81	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.