



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2024 – 11:06 AM EDT

PDB ID : 3GLY  
Title : REFINED CRYSTAL STRUCTURES OF GLUCOAMYLASE FROM AS-PERGILLUS AWAMORI VAR. X100  
Authors : Aleshin, A.E.; Hoffman, C.; Firsov, L.M.; Honzatko, R.B.  
Deposited on : 1994-06-03  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	470	
2	B	5	
3	C	8	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLUCOAMYLASE-471.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	470	Total	C	N	O	S	0	5	0
			3578	2243	586	741	8			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	58	LEU	ILE	conflict	UNP P22832
A	60	ILE	LEU	conflict	UNP P22832
A	117	THR	ALA	conflict	UNP P22832

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 4 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	1
			12	6	6		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	673	Total 673	O 673	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: GLUCOAMYLASE-471



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.70Å 104.30Å 48.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.141 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4517	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.01	2/3693 (0.1%)	1.37	32/5051 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	397	SER	CB-OG	23.77	1.73	1.42
1	A	397	SER	CA-CB	10.07	1.68	1.52

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	ARG	NE-CZ-NH2	-11.88	114.36	120.30
1	A	273	ARG	NE-CZ-NH1	10.28	125.44	120.30
1	A	397	SER	N-CA-CB	-9.74	95.88	110.50
1	A	273	ARG	NE-CZ-NH2	-9.18	115.71	120.30
1	A	68	ARG	NE-CZ-NH1	8.81	124.71	120.30
1	A	397	SER	CA-CB-OG	-8.76	87.55	111.20
1	A	241	ARG	NE-CZ-NH1	8.47	124.53	120.30
1	A	194	ARG	NE-CZ-NH2	-8.34	116.13	120.30
1	A	241	ARG	NE-CZ-NH2	-8.22	116.19	120.30
1	A	428	ARG	NE-CZ-NH2	-8.07	116.26	120.30
1	A	293	ASP	CB-CG-OD1	7.90	125.41	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	414	ASP	CB-CG-OD2	7.43	124.99	118.30
1	A	160	ARG	NE-CZ-NH2	-7.39	116.61	120.30
1	A	131	ARG	NE-CZ-NH1	6.83	123.71	120.30
1	A	289	TYR	CB-CG-CD1	6.39	124.83	121.00
1	A	305	ARG	NE-CZ-NH2	-6.27	117.16	120.30
1	A	341	LEU	CA-CB-CG	6.10	129.34	115.30
1	A	429	ARG	NE-CZ-NH1	5.97	123.28	120.30
1	A	15	ARG	CD-NE-CZ	5.96	131.95	123.60
1	A	257	ASP	CB-CG-OD2	5.83	123.55	118.30
1	A	194	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	283	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	418	SER	N-CA-CB	5.45	118.67	110.50
1	A	286	ARG	NE-CZ-NH1	5.31	122.96	120.30
1	A	406	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	297	ASP	CB-CG-OD2	-5.21	113.61	118.30
1	A	254	HIS	CA-CB-CG	-5.19	104.78	113.60
1	A	396	GLY	C-N-CA	-5.17	108.78	121.70
1	A	374	VAL	CB-CA-C	5.16	121.20	111.40
1	A	396	GLY	O-C-N	5.12	130.90	122.70
1	A	398	LEU	C-N-CA	5.07	134.37	121.70
1	A	413	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	25[A]	ASP	Sidechain
1	A	25[B]	ASP	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3578	0	3322	39	0
2	B	61	0	52	0	0
3	C	94	0	79	4	0
4	A	111	0	96	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	673	0	0	4	0
All	All	4517	0	3549	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397:SER:CB	1:A:397:SER:OG	1.73	1.36
1:A:397:SER:OG	1:A:397:SER:CA	2.35	0.73
1:A:43:THR:HG23	3:C:5:MAN:H61	1.77	0.67
1:A:395:ASN:OD1	1:A:397:SER:HB2	2.03	0.58
1:A:337[A]:LYS:HE2	5:A:1024:HOH:O	2.03	0.58
1:A:43:THR:CG2	3:C:5:MAN:H61	2.34	0.57
3:C:6:MAN:H61	3:C:8:MAN:C5	2.34	0.57
1:A:217:ALA:HB3	1:A:218:PRO:HD3	1.86	0.57
1:A:3:LEU:HD11	1:A:385:VAL:HG12	1.87	0.56
1:A:316:PRO:HD2	1:A:402:PHE:O	2.07	0.54
1:A:325:ALA:HB2	1:A:384:PHE:HB2	1.91	0.53
1:A:2:THR:HG23	1:A:3:LEU:H	1.75	0.52
1:A:397:SER:CB	1:A:397:SER:HG	2.16	0.51
1:A:312:TYR:O	1:A:313:ASN:HB2	2.11	0.51
1:A:21:ASN:HD21	1:A:28:TRP:HE3	1.62	0.48
1:A:21:ASN:ND2	1:A:28:TRP:HE3	2.12	0.47
3:C:6:MAN:H61	3:C:8:MAN:H5	1.94	0.47
1:A:6:TRP:CH2	1:A:10:GLU:HG2	2.50	0.47
1:A:59:VAL:HG21	1:A:416:THR:HG22	1.96	0.47
1:A:136:ILE:O	1:A:140:GLN:HG3	2.16	0.46
1:A:60:ILE:HG21	1:A:81:TYR:CD2	2.51	0.45
1:A:189:ILE:HG22	1:A:224:LEU:HD13	1.99	0.45
1:A:428:ARG:CZ	1:A:434:PRO:HD3	2.47	0.44
1:A:341:LEU:HD11	1:A:373:ILE:HG21	1.99	0.44
1:A:293:ASP:HB3	5:A:1112:HOH:O	2.18	0.44
1:A:44:ASP:HB3	5:A:1037:HOH:O	2.17	0.43
1:A:352:LYS:HE2	5:A:1045:HOH:O	2.16	0.43
1:A:315:ASN:HB3	1:A:316:PRO:HD2	2.00	0.43
1:A:369:THR:HG22	1:A:373:ILE:HD12	2.00	0.43
1:A:445:VAL:HA	1:A:446:PRO:HD3	1.85	0.42
1:A:309:ASP:O	1:A:314:GLY:HA3	2.19	0.42
1:A:424:THR:O	1:A:428:ARG:HG2	2.19	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:ILE:HG23	1:A:220:ILE:HG23	2.01	0.41
1:A:427:ASN:HD22	1:A:427:ASN:HA	1.73	0.41
1:A:235:ALA:HB2	1:A:245:ASP:HB3	2.03	0.41
1:A:81:TYR:O	1:A:85:GLN:HG2	2.21	0.41
1:A:2:THR:HG23	1:A:3:LEU:N	2.34	0.40
1:A:193:HIS:O	1:A:197:VAL:HG23	2.21	0.40
1:A:365:SER:HA	1:A:370:TYR:CG	2.56	0.40
1:A:55[B]:ASP:OD1	1:A:417:TRP:HB2	2.21	0.40
1:A:106:GLU:HA	1:A:107:PRO:HD3	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/470 (101%)	454 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	390/385 (101%)	381 (98%)	9 (2%)	45	59

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ASP
1	A	247	ASN
1	A	341	LEU
1	A	397	SER
1	A	399	SER
1	A	410	LEU
1	A	422	LEU
1	A	441	SER
1	A	464	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	168	GLN
1	A	426	ASN
1	A	427	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	1.31	2 (14%)	17,19,21	1.73	2 (11%)
2	NAG	B	2	2	14,14,15	0.98	1 (7%)	17,19,21	1.45	2 (11%)
2	BMA	B	3	2	11,11,12	0.59	0	15,15,17	1.32	1 (6%)
2	MAN	B	4	2	11,11,12	0.78	1 (9%)	15,15,17	0.99	1 (6%)
2	MAN	B	5	2	11,11,12	0.90	0	15,15,17	1.43	3 (20%)
3	NAG	C	1	3,1	14,14,15	1.08	1 (7%)	17,19,21	1.41	3 (17%)
3	NAG	C	2	3	14,14,15	1.01	1 (7%)	17,19,21	1.30	2 (11%)
3	BMA	C	3	3	11,11,12	0.54	0	15,15,17	1.08	1 (6%)
3	MAN	C	4	3	11,11,12	0.41	0	15,15,17	1.18	1 (6%)
3	MAN	C	5	3	11,11,12	0.63	0	15,15,17	1.21	2 (13%)
3	MAN	C	6	3	11,11,12	0.53	0	15,15,17	1.08	1 (6%)
3	MAN	C	7	3	11,11,12	0.51	0	15,15,17	1.14	2 (13%)
3	MAN	C	8	3	11,11,12	0.68	0	15,15,17	1.02	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	2/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	1/2/19/22	0/1/1/1
3	MAN	C	5	3	-	2/2/19/22	0/1/1/1
3	MAN	C	6	3	-	2/2/19/22	0/1/1/1
3	MAN	C	7	3	-	0/2/19/22	0/1/1/1
3	MAN	C	8	3	-	2/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C8-C7	3.23	1.57	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	NAG	C8-C7	2.98	1.56	1.50
2	B	2	NAG	C8-C7	2.85	1.56	1.50
2	B	1	NAG	C2-N2	2.78	1.50	1.46
3	C	1	NAG	C8-C7	2.74	1.56	1.50
2	B	4	MAN	O5-C5	2.15	1.47	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	4.90	118.75	112.19
2	B	3	BMA	C1-O5-C5	4.04	117.60	112.19
2	B	1	NAG	C2-N2-C7	-4.03	117.49	122.90
3	C	2	NAG	O4-C4-C5	-3.75	100.09	109.32
3	C	4	MAN	C1-O5-C5	3.75	117.21	112.19
2	B	2	NAG	O5-C1-C2	-3.49	105.88	111.29
3	C	1	NAG	C6-C5-C4	3.21	120.89	113.02
3	C	6	MAN	C1-O5-C5	3.14	116.39	112.19
2	B	2	NAG	O5-C5-C4	-3.11	103.26	110.83
3	C	5	MAN	C1-O5-C5	3.00	116.21	112.19
3	C	1	NAG	O5-C1-C2	-2.93	106.76	111.29
3	C	5	MAN	O6-C6-C5	2.90	121.21	111.33
3	C	3	BMA	C1-O5-C5	2.71	115.82	112.19
2	B	5	MAN	C6-C5-C4	2.58	119.37	113.02
2	B	5	MAN	C3-C4-C5	-2.55	105.61	110.23
2	B	5	MAN	O5-C5-C4	-2.54	104.64	110.83
3	C	1	NAG	O5-C5-C4	-2.44	104.89	110.83
2	B	4	MAN	C6-C5-C4	2.24	118.53	113.02
3	C	8	MAN	C1-O5-C5	2.14	115.05	112.19
3	C	7	MAN	O6-C6-C5	2.07	118.39	111.33
3	C	2	NAG	O5-C5-C4	-2.04	105.86	110.83
3	C	7	MAN	C1-O5-C5	2.02	114.90	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	8	MAN	C4-C5-C6-O6
3	C	8	MAN	O5-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6
3	C	6	MAN	C4-C5-C6-O6
3	C	6	MAN	O5-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	B	3	BMA	C4-C5-C6-O6
3	C	5	MAN	C4-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
3	C	5	MAN	O5-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	6	MAN	2	0
3	C	8	MAN	2	0
3	C	5	MAN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MAN	A	488(A)	1	11,11,12	0.62	0	15,15,17	0.77	0
4	MAN	A	485(A)	1	11,11,12	0.55	0	15,15,17	0.90	1 (6%)
4	MAN	A	494(A)	1	11,11,12	0.85	1 (9%)	15,15,17	1.35	2 (13%)
4	MAN	A	493(A)	1	11,11,12	0.58	0	15,15,17	1.56	1 (6%)
4	MAN	A	489(A)	1	11,11,12	0.68	0	15,15,17	1.21	1 (6%)
4	MAN	A	486(A)[B]	-	11,11,12	0.91	1 (9%)	15,15,17	1.22	3 (20%)
4	MAN	A	491(A)	1	11,11,12	0.70	0	15,15,17	1.26	2 (13%)
4	MAN	A	492(A)	1	11,11,12	0.94	0	15,15,17	1.07	1 (6%)
4	MAN	A	487(A)	1	11,11,12	0.82	1 (9%)	15,15,17	0.96	1 (6%)
4	MAN	A	486(A)[A]	-	11,11,12	0.91	1 (9%)	15,15,17	1.15	2 (13%)
4	MAN	A	490(A)	1	11,11,12	0.90	1 (9%)	15,15,17	1.28	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	488(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	485(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	494(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	493(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	489(A)	1	-	2/2/19/22	0/1/1/1
4	MAN	A	486(A)[B]	-	-	2/2/19/22	0/1/1/1
4	MAN	A	491(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	492(A)	1	-	0/2/19/22	0/1/1/1
4	MAN	A	487(A)	1	-	2/2/19/22	0/1/1/1
4	MAN	A	486(A)[A]	-	-	0/2/19/22	0/1/1/1
4	MAN	A	490(A)	1	-	2/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	486(A)[A]	MAN	O5-C5	2.75	1.48	1.43
4	A	486(A)[B]	MAN	O5-C5	2.75	1.48	1.43
4	A	494(A)	MAN	O5-C1	2.45	1.47	1.43
4	A	487(A)	MAN	O5-C5	2.29	1.47	1.43
4	A	490(A)	MAN	O5-C1	2.21	1.47	1.43

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	493(A)	MAN	C1-O5-C5	5.50	119.55	112.19
4	A	494(A)	MAN	C1-O5-C5	4.18	117.79	112.19
4	A	491(A)	MAN	C1-O5-C5	3.95	117.48	112.19
4	A	489(A)	MAN	C1-O5-C5	3.38	116.71	112.19
4	A	490(A)	MAN	C1-O5-C5	3.32	116.63	112.19
4	A	492(A)	MAN	O3-C3-C4	-2.68	104.05	110.38
4	A	486(A)[A]	MAN	O5-C5-C4	-2.49	104.76	110.83
4	A	486(A)[B]	MAN	O5-C5-C4	-2.49	104.76	110.83
4	A	486(A)[B]	MAN	O6-C6-C5	2.41	119.53	111.33
4	A	494(A)	MAN	O5-C1-C2	2.39	116.49	110.79
4	A	486(A)[A]	MAN	C6-C5-C4	2.31	118.70	113.02
4	A	486(A)[B]	MAN	C6-C5-C4	2.31	118.70	113.02
4	A	491(A)	MAN	O3-C3-C4	-2.13	105.35	110.38
4	A	487(A)	MAN	O3-C3-C4	-2.09	105.45	110.38
4	A	485(A)	MAN	C1-O5-C5	2.05	114.93	112.19

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	486(A)[B]	MAN	C4-C5-C6-O6
4	A	487(A)	MAN	O5-C5-C6-O6
4	A	489(A)	MAN	O5-C5-C6-O6
4	A	490(A)	MAN	O5-C5-C6-O6
4	A	487(A)	MAN	C4-C5-C6-O6
4	A	489(A)	MAN	C4-C5-C6-O6
4	A	486(A)[B]	MAN	O5-C5-C6-O6
4	A	490(A)	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.