



# wwPDB EM Validation Summary Report ⓘ

Jan 16, 2025 – 12:34 am GMT

PDB ID : 9H3O  
EMDB ID : EMD-51832  
Title : 50S subunit precursor C\_(L22)-\_GAC  
Authors : Lauer, S.; Nikolay, R.; Spahn, C.M.T.  
Deposited on : 2024-10-17  
Resolution : 4.54 Å(reported)  
Based on initial model : 8RPY

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	2	46	
2	A	2904	
3	D	209	
4	E	201	
5	J	142	
6	K	122	
7	L	143	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
8	N	120	
9	P	114	
10	Q	117	
11	R	103	
12	T	93	
13	U	102	
14	Y	63	
15	4	38	
16	G	176	
17	Z	58	

## 2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 60003 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Large ribosomal subunit protein bL34.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	2	39	Total	C	N	O	S	0	0
			312	187	77	47	1		

- Molecule 2 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	A	2166	Total	C	N	O	P	0	0
			46524	20754	8595	15009	2166		

- Molecule 3 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	175	Total	C	N	O	S	0	0
			1306	820	235	247	4		

- Molecule 4 is a protein called Large ribosomal subunit protein uL4.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	165	Total	C	N	O	S	0	0
			1282	809	225	243	5		

- Molecule 5 is a protein called Large ribosomal subunit protein uL13.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	J	142	Total	C	N	O	S	0	0
			1129	714	212	199	4		

- Molecule 6 is a protein called Large ribosomal subunit protein uL14.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	K	122	Total	C	N	O	S	0	0
			939	587	180	166	6		

- Molecule 7 is a protein called Large ribosomal subunit protein uL15.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	L	109	Total	C	N	O	0	0
			778	483	149	146		

- Molecule 8 is a protein called Large ribosomal subunit protein bL17.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	N	120	Total	C	N	O	S	0	0
			961	593	196	167	5		

- Molecule 9 is a protein called Large ribosomal subunit protein bL19.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	P	114	Total	C	N	O	S	0	0
			917	574	179	163	1		

- Molecule 10 is a protein called Large ribosomal subunit protein bL20.

Mol	Chain	Residues	Atoms				AltConf	Trace
10	Q	117	Total	C	N	O	0	0
			947	604	192	151		

- Molecule 11 is a protein called Large ribosomal subunit protein bL21.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	R	103	Total	C	N	O	S	0	0
			816	516	153	145	2		

- Molecule 12 is a protein called Large ribosomal subunit protein uL23.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	T	93	Total	C	N	O	S	0	0
			739	466	139	132	2		

- Molecule 13 is a protein called Large ribosomal subunit protein uL24.

Mol	Chain	Residues	Atoms				AltConf	Trace
13	U	102	Total	C	N	O	0	0
			780	492	146	142		

- Molecule 14 is a protein called Large ribosomal subunit protein uL29.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	Y	63	Total	C	N	O	S	0	0
			509	313	99	95	2		

- Molecule 15 is a protein called Large ribosomal subunit protein bL36A.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	4	38	Total	C	N	O	S	0	0
			302	185	65	48	4		

- Molecule 16 is a protein called Large ribosomal subunit protein uL6.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	G	176	Total	C	N	O	S	0	0
			1323	832	243	246	2		

- Molecule 17 is a protein called Large ribosomal subunit protein uL30.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Z	57	Total	C	N	O	S	0	0
			439	276	86	75	2		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Large ribosomal subunit protein bL34

Chain 2: 




- Molecule 2: 23S ribosomal RNA

Chain A: 












• Molecule 3: 50S ribosomal protein L3



• Molecule 4: Large ribosomal subunit protein uL4





• Molecule 5: Large ribosomal subunit protein uL13

Chain J:  80% 20%




- Molecule 6: Large ribosomal subunit protein uL14

Chain K:  29% 81% 19%



- Molecule 7: Large ribosomal subunit protein uL15

Chain L:  19% 66% 10% 24%





- Molecule 8: Large ribosomal subunit protein bL17

Chain N:  87% 13%




- Molecule 9: Large ribosomal subunit protein bL19

Chain P:  31% 89% 11%




- Molecule 10: Large ribosomal subunit protein bL20



Chain Q:  82% 18%





- Molecule 11: Large ribosomal subunit protein bL21





- Molecule 12: Large ribosomal subunit protein uL23




- Molecule 13: Large ribosomal subunit protein uL24



- Molecule 14: Large ribosomal subunit protein uL29



- Molecule 15: Large ribosomal subunit protein bL36A




- Molecule 16: Large ribosomal subunit protein uL6





• Molecule 17: Large ribosomal subunit protein uL30



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	9154	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	45	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	1.490	Depositor
Minimum map value	-0.285	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.075	Depositor
Recommended contour level	0.268	Depositor
Map size (Å)	424.0, 424.0, 424.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.4133333, 1.4133333, 1.4133333	Depositor

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	2	0.23	0/314	0.65	0/413
2	A	0.50	7/52108 (0.0%)	0.76	29/81280 (0.0%)
3	D	0.24	0/1320	0.51	0/1776
4	E	0.24	0/1294	0.46	0/1741
5	J	0.24	0/1152	0.50	0/1551
6	K	0.24	0/948	0.54	0/1268
7	L	0.24	0/780	0.54	0/1041
8	N	0.24	0/974	0.58	0/1301
9	P	0.24	0/929	0.54	0/1242
10	Q	0.24	0/960	0.51	0/1278
11	R	0.25	0/829	0.52	0/1107
12	T	0.24	0/745	0.50	0/994
13	U	0.25	0/788	0.50	0/1051
14	Y	0.25	0/510	0.51	0/677
15	4	0.25	0/303	0.55	0/397
16	G	1.04	1/1343 (0.1%)	0.65	3/1816 (0.2%)
17	Z	0.80	0/443	0.88	1/593 (0.2%)
All	All	0.48	8/65740 (0.0%)	0.72	33/99526 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
16	G	0	7
17	Z	0	3
All	All	0	10

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2657	A	N3-C4	52.27	1.66	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2657	A	C6-N1	51.56	1.71	1.35
2	A	2657	A	N1-C2	40.02	1.70	1.34
2	A	2657	A	C5-C4	37.45	1.65	1.38
2	A	2657	A	C2-N3	36.53	1.66	1.33

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2657	A	N1-C2-N3	-27.29	115.66	129.30
2	A	2657	A	C2-N3-C4	21.84	121.52	110.60
2	A	929	U	OP1-P-O3'	-13.80	74.84	105.20
2	A	929	U	O3'-P-O5'	10.84	124.59	104.00
2	A	2657	A	N7-C8-N9	9.62	118.61	113.80

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
16	G	131	VAL	Peptide
16	G	86	LEU	Peptide
16	G	88	LEU	Peptide
16	G	89	VAL	Peptide
16	G	90	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	312	0	344	10	0
2	A	46524	0	23411	443	0
3	D	1306	0	1355	24	0
4	E	1282	0	1336	14	0
5	J	1129	0	1162	23	0
6	K	939	0	1012	16	0
7	L	778	0	841	10	0
8	N	961	0	1000	15	0
9	P	917	0	965	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	Q	947	0	1022	20	0
11	R	816	0	839	14	0
12	T	739	0	807	10	0
13	U	780	0	834	9	0
14	Y	509	0	543	5	0
15	4	302	0	343	1	0
16	G	1323	0	1374	147	0
17	Z	439	0	485	55	0
All	All	60003	0	37673	701	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 701 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2657:A:N1	2:A:2657:A:C6	1.71	1.55
2:A:2657:A:N1	2:A:2657:A:C2	1.70	1.53
2:A:2657:A:C2	16:G:176:LYS:C	1.92	1.42
2:A:2657:A:C2	16:G:176:LYS:CA	2.05	1.39
16:G:129:GLU:O	16:G:163:TYR:C	1.64	1.33

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	2	37/46 (80%)	37 (100%)	0	0	100	100
3	D	171/209 (82%)	166 (97%)	5 (3%)	0	100	100
4	E	161/201 (80%)	156 (97%)	5 (3%)	0	100	100

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	J	140/142 (99%)	137 (98%)	3 (2%)	0	100	100
6	K	120/122 (98%)	115 (96%)	5 (4%)	0	100	100
7	L	105/143 (73%)	96 (91%)	9 (9%)	0	100	100
8	N	118/120 (98%)	110 (93%)	8 (7%)	0	100	100
9	P	112/114 (98%)	112 (100%)	0	0	100	100
10	Q	115/117 (98%)	112 (97%)	3 (3%)	0	100	100
11	R	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
12	T	91/93 (98%)	84 (92%)	7 (8%)	0	100	100
13	U	100/102 (98%)	94 (94%)	6 (6%)	0	100	100
14	Y	61/63 (97%)	56 (92%)	5 (8%)	0	100	100
15	4	36/38 (95%)	35 (97%)	1 (3%)	0	100	100
16	G	174/176 (99%)	131 (75%)	39 (22%)	4 (2%)	5	29
17	Z	55/58 (95%)	47 (86%)	6 (11%)	2 (4%)	3	20
All	All	1697/1847 (92%)	1588 (94%)	103 (6%)	6 (0%)	32	68

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
16	G	153	PRO
16	G	142	GLN
16	G	163	TYR
17	Z	13	ILE
17	Z	42	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2	31/38 (82%)	30 (97%)	1 (3%)	34	55
3	D	136/164 (83%)	135 (99%)	1 (1%)	81	87
4	E	138/165 (84%)	138 (100%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	J	116/116 (100%)	115 (99%)	1 (1%)	75	83
6	K	103/103 (100%)	103 (100%)	0	100	100
7	L	76/102 (74%)	76 (100%)	0	100	100
8	N	100/100 (100%)	100 (100%)	0	100	100
9	P	99/99 (100%)	99 (100%)	0	100	100
10	Q	89/89 (100%)	89 (100%)	0	100	100
11	R	84/84 (100%)	84 (100%)	0	100	100
12	T	80/80 (100%)	80 (100%)	0	100	100
13	U	83/83 (100%)	83 (100%)	0	100	100
14	Y	55/55 (100%)	55 (100%)	0	100	100
15	4	34/34 (100%)	34 (100%)	0	100	100
16	G	137/137 (100%)	123 (90%)	14 (10%)	6	21
17	Z	47/48 (98%)	38 (81%)	9 (19%)	1	7
All	All	1408/1497 (94%)	1382 (98%)	26 (2%)	54	71

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
16	G	151	ARG
17	Z	9	THR
17	Z	52	PHE
16	G	162	ARG
17	Z	28	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	D	164	GLN
5	J	136	GLN
8	N	107	ASN
11	R	43	ASN
15	4	35	GLN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	A	2158/2904 (74%)	385 (17%)	11 (0%)

5 of 385 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	A	10	A
2	A	12	U
2	A	25	U
2	A	27	G
2	A	46	G

5 of 11 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	A	1736	U
2	A	1737	G
2	A	2763	G
2	A	1738	G
2	A	828	U

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-51832. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


### 6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 164




Y Index: 142




Z Index: 148


### 6.3.2 Raw map



X Index: 164



Y Index: 142




Z Index: 149

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.268. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_51832\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1209 nm<sup>3</sup>; this corresponds to an approximate mass of 1092 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.220  $\text{\AA}^{-1}$

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.220 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.54	-	-
Author-provided FSC curve	4.49	6.70	4.64
Unmasked-calculated*	9.30	15.95	9.88

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 9.30 differs from the reported value 4.54 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-51832 and PDB model 9H3O. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.268 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.268).







































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 91% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.268) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9060	 0.1970
2	 0.9180	 0.1900
4	 0.8150	 0.1380
A	 0.9250	 0.2000
D	 0.8740	 0.2180
E	 0.8910	 0.2200
G	 0.8420	 0.1490
J	 0.9310	 0.2410
K	 0.6030	 0.0990
L	 0.6980	 0.1030
N	 0.9310	 0.1670
P	 0.6080	 0.1360
Q	 0.9470	 0.2320
R	 0.8900	 0.2160
T	 0.8910	 0.2350
U	 0.9220	 0.2420
Y	 0.8630	 0.1990
Z	 0.8060	 0.1890

