



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 26, 2025 – 12:29 am GMT

PDB ID : 9HEC
Title : Unspecific peroxygenase from *Psathyrella aberdarensis* (PabUPO-II) in complex with alpha-ionone
Authors : Fernandez-Garcia, A.; Sanz-Aparicio, J.
Deposited on : 2024-11-13
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div>2%</div> <div>94%</div> <div>5%</div> </div>
1	B	335	<div> <div>4%</div> <div>96%</div> <div>.</div> </div>
1	C	335	<div> <div>4%</div> <div>94%</div> <div>6%</div> </div>
2	G	5	<div> <div>100%</div> </div>
3	I	7	<div> <div>29%</div> <div>71%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	W	3	 100%
4	l	3	 67% 33%
5	X	6	 17% 50% 33%
6	j	4	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
12	ACT	B	403	-	-	X	-
7	A1IUB	C	401	-	-	X	-
7	A1IUB	C	403	-	-	X	-

2 Entry composition [i](#)


There are 13 unique types of molecules in this entry. The entry contains 9032 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heme-thiolate peroxidase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	10	0
			2646	1682	456	501	7			
1	B	335	Total	C	N	O	S	0	5	0
			2620	1665	452	496	7			
1	C	335	Total	C	N	O	S	0	6	0
			2621	1666	451	497	7			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	W	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	1	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	X	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	j	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 7 is alpha-ionone (three-letter code: A1IUB) (formula: C₁₃H₂₀O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			14	13	1		
7	A	1	Total	C	O	0	0
			14	13	1		
7	B	1	Total	C	O	0	0
			14	13	1		
7	C	1	Total	C	O	0	0
			14	13	1		
7	C	1	Total	C	O	0	0
			14	13	1		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total 6	C 3	O 3	0	0
8	B	1	Total 6	C 3	O 3	0	0
8	C	1	Total 6	C 3	O 3	0	0

- Molecule 9 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
9	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
9	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total Mg 1 1	0	0
10	B	1	Total Mg 1 1	0	0
10	C	1	Total Mg 1 1	0	0

- Molecule 11 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	O	S	0	0
			5	4	1		
11	B	1	Total	O	S	0	0
			5	4	1		
11	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 12 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	253	Total	O	0	0
			253	253		
13	B	154	Total	O	0	0
			154	154		
13	C	155	Total	O	0	0
			155	155		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heme-thiolate peroxidase



- Molecule 1: Heme-thiolate peroxidase



- Molecule 1: Heme-thiolate peroxidase



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain W: 100%




- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain l: 67% 33%



- Molecule 5: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain X: 17% 50% 33%



- Molecule 6: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain j: 100%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	270.97Å 75.05Å 106.06Å 90.00° 112.06° 90.00°	Depositor
Resolution (Å)	49.19 – 2.20 49.19 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.2 (49.19-2.20) 98.2 (49.19-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.182 , 0.213 0.188 , 0.218	Depositor DCC
R_{free} test set	4898 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	35.9	Xtriage
Anisotropy	0.782	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.014 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9032	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, ACT, NAG, HEM, BMA, MG, SO4, A1IUB, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2757	0.74	3/3767 (0.1%)
1	B	0.40	0/2716	0.72	0/3712
1	C	0.41	0/2720	0.70	0/3718
All	All	0.41	0/8193	0.72	3/11197 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	ARG	NE-CZ-NH1	-5.73	117.44	120.30
1	A	318[A]	ARG	NE-CZ-NH1	-5.08	117.76	120.30
1	A	318[B]	ARG	NE-CZ-NH1	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2646	0	2506	22	0
1	B	2620	0	2469	7	0
1	C	2621	0	2473	23	0
2	G	61	0	52	0	0
3	I	83	0	70	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	W	39	0	34	0	0
4	l	39	0	34	0	0
5	X	72	0	61	2	0
6	j	50	0	43	0	0
7	A	28	0	0	7	0
7	B	14	0	0	2	0
7	C	28	0	0	15	0
8	A	6	0	8	0	0
8	B	6	0	8	0	0
8	C	6	0	8	0	0
9	A	43	0	30	1	0
9	B	43	0	30	4	0
9	C	43	0	30	6	0
10	A	1	0	0	0	0
10	B	1	0	0	0	0
10	C	1	0	0	0	0
11	A	5	0	0	0	0
11	B	5	0	0	0	0
11	C	5	0	0	0	0
12	B	4	0	3	2	0
13	A	253	0	0	10	0
13	B	154	0	0	3	0
13	C	155	0	0	5	0
All	All	9032	0	7859	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 68 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212[A]:GLU:HG3	7:A:401:A1IUB:C1	1.83	1.09
7:C:401:A1IUB:C14	7:C:403:A1IUB:C4	2.35	1.03
1:C:92:VAL:HG11	7:C:403:A1IUB:C13	2.06	0.85
1:C:208:THR:HB	7:C:401:A1IUB:C1	2.08	0.83
1:A:136[A]:THR:HG23	13:A:607:HOH:O	1.79	0.83

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/335 (102%)	335 (98%)	7 (2%)	1 (0%)	37	42
1	B	338/335 (101%)	330 (98%)	7 (2%)	1 (0%)	37	42
1	C	339/335 (101%)	331 (98%)	7 (2%)	1 (0%)	37	42
All	All	1020/1005 (102%)	996 (98%)	21 (2%)	3 (0%)	37	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	257	SER
1	B	257	SER
1	C	257	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/274 (104%)	284 (100%)	0	100	100
1	B	279/274 (102%)	278 (100%)	1 (0%)	89	95
1	C	280/274 (102%)	280 (100%)	0	100	100
All	All	843/822 (103%)	842 (100%)	1 (0%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	137	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

28 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	G	1	1,2	14,14,15	0.43	0	17,19,21	1.37	2 (11%)
2	NAG	G	2	2	14,14,15	0.38	0	17,19,21	1.01	1 (5%)
2	BMA	G	3	2	11,11,12	0.69	0	15,15,17	1.10	1 (6%)
2	MAN	G	4	2	11,11,12	0.79	0	15,15,17	1.03	2 (13%)
2	MAN	G	5	2	11,11,12	0.74	0	15,15,17	1.66	3 (20%)
3	NAG	I	1	3,1	14,14,15	0.34	0	17,19,21	0.73	0
3	NAG	I	2	3	14,14,15	0.35	0	17,19,21	0.91	0
3	BMA	I	3	3	11,11,12	0.46	0	15,15,17	1.03	1 (6%)
3	MAN	I	4	3	11,11,12	0.77	0	15,15,17	1.07	1 (6%)
3	MAN	I	5	3	11,11,12	0.81	0	15,15,17	0.94	1 (6%)
3	MAN	I	6	3	11,11,12	0.62	0	15,15,17	1.01	1 (6%)
3	MAN	I	7	3	11,11,12	0.86	0	15,15,17	1.65	3 (20%)
4	NAG	W	1	4,1	14,14,15	0.39	0	17,19,21	2.03	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	W	2	4	14,14,15	0.32	0	17,19,21	0.96	1 (5%)
4	BMA	W	3	4	11,11,12	0.60	0	15,15,17	0.88	1 (6%)
5	NAG	X	1	5,1	14,14,15	0.34	0	17,19,21	0.77	0
5	NAG	X	2	5	14,14,15	0.39	0	17,19,21	0.93	1 (5%)
5	BMA	X	3	5	11,11,12	0.47	0	15,15,17	1.02	1 (6%)
5	MAN	X	4	5	11,11,12	0.71	0	15,15,17	1.50	3 (20%)
5	MAN	X	5	5	11,11,12	0.84	1 (9%)	15,15,17	1.11	1 (6%)
5	MAN	X	6	5	11,11,12	0.61	0	15,15,17	1.47	2 (13%)
6	NAG	j	1	1,6	14,14,15	0.30	0	17,19,21	0.98	1 (5%)
6	NAG	j	2	6	14,14,15	0.38	0	17,19,21	1.18	1 (5%)
6	BMA	j	3	6	11,11,12	0.44	0	15,15,17	1.04	1 (6%)
6	MAN	j	4	6	11,11,12	0.76	0	15,15,17	1.07	2 (13%)
4	NAG	l	1	4,1	14,14,15	0.32	0	17,19,21	0.47	0
4	NAG	l	2	4	14,14,15	0.42	0	17,19,21	1.60	4 (23%)
4	BMA	l	3	4	11,11,12	0.44	0	15,15,17	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	BMA	G	3	2	-	2/2/19/22	0/1/1/1
2	MAN	G	4	2	-	2/2/19/22	0/1/1/1
2	MAN	G	5	2	-	0/2/19/22	0/1/1/1
3	NAG	I	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	0/1/1/1
3	MAN	I	4	3	-	0/2/19/22	0/1/1/1
3	MAN	I	5	3	-	2/2/19/22	0/1/1/1
3	MAN	I	6	3	-	1/2/19/22	0/1/1/1
3	MAN	I	7	3	-	0/2/19/22	0/1/1/1
4	NAG	W	1	4,1	-	3/6/23/26	0/1/1/1
4	NAG	W	2	4	-	0/6/23/26	0/1/1/1
4	BMA	W	3	4	-	0/2/19/22	0/1/1/1
5	NAG	X	1	5,1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	X	2	5	-	0/6/23/26	0/1/1/1
5	BMA	X	3	5	-	2/2/19/22	0/1/1/1
5	MAN	X	4	5	-	1/2/19/22	0/1/1/1
5	MAN	X	5	5	-	2/2/19/22	0/1/1/1
5	MAN	X	6	5	-	2/2/19/22	0/1/1/1
6	NAG	j	1	1,6	-	3/6/23/26	0/1/1/1
6	NAG	j	2	6	-	4/6/23/26	0/1/1/1
6	BMA	j	3	6	-	0/2/19/22	0/1/1/1
6	MAN	j	4	6	-	2/2/19/22	0/1/1/1
4	NAG	l	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	l	2	4	-	0/6/23/26	0/1/1/1
4	BMA	l	3	4	-	1/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	X	5	MAN	O5-C5	2.06	1.47	1.43

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	W	1	NAG	C1-C2-N2	6.63	121.81	110.49
5	X	6	MAN	C1-C2-C3	4.84	115.62	109.67
3	I	7	MAN	C1-C2-C3	4.68	115.42	109.67
2	G	5	MAN	C1-O5-C5	4.31	118.03	112.19
2	G	5	MAN	C1-C2-C3	4.07	114.67	109.67

There are no chirality outliers.

5 of 35 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	W	1	NAG	C1-C2-N2-C7
6	j	1	NAG	O7-C7-N2-C2
6	j	1	NAG	C8-C7-N2-C2
2	G	3	BMA	O5-C5-C6-O6
2	G	4	MAN	C4-C5-C6-O6



There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	X	6	MAN	2	0
5	X	3	BMA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 3 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	GOL	B	402	-	5,5,5	0.20	0	5,5,5	0.50	0
8	GOL	A	403	-	5,5,5	0.20	0	5,5,5	0.59	0
11	SO4	C	406	-	4,4,4	0.18	0	6,6,6	0.16	0
8	GOL	C	402	-	5,5,5	0.22	0	5,5,5	0.48	0
7	A1IUB	C	403	-	13,14,14	0.68	1 (7%)	14,20,20	0.56	0
11	SO4	A	406	-	4,4,4	0.21	0	6,6,6	0.12	0
7	A1IUB	A	402	-	13,14,14	0.52	0	14,20,20	0.68	0
7	A1IUB	B	401	-	13,14,14	0.44	0	14,20,20	0.75	1 (7%)
12	ACT	B	403	-	3,3,3	1.28	0	3,3,3	0.34	0
9	HEM	C	404	10,1	41,50,50	1.37	3 (7%)	45,82,82	2.10	17 (37%)
7	A1IUB	A	401	-	13,14,14	1.06	1 (7%)	14,20,20	1.08	2 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	SO4	B	406	-	4,4,4	0.21	0	6,6,6	0.07	0
9	HEM	B	404	10,1	41,50,50	1.47	7 (17%)	45,82,82	2.04	17 (37%)
7	A1IUB	C	401	-	13,14,14	1.14	1 (7%)	14,20,20	0.72	0
9	HEM	A	404	10,1	41,50,50	1.30	5 (12%)	45,82,82	2.02	14 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	B	402	-	-	0/4/4/4	-
8	GOL	A	403	-	-	3/4/4/4	-
8	GOL	C	402	-	-	0/4/4/4	-
7	A1IUB	C	403	-	-	0/5/22/22	0/1/1/1
7	A1IUB	A	402	-	-	1/5/22/22	0/1/1/1
7	A1IUB	B	401	-	-	1/5/22/22	0/1/1/1
9	HEM	C	404	10,1	-	3/12/54/54	-
7	A1IUB	A	401	-	-	2/5/22/22	0/1/1/1
9	HEM	B	404	10,1	-	1/12/54/54	-
7	A1IUB	C	401	-	-	0/5/22/22	0/1/1/1
9	HEM	A	404	10,1	-	1/12/54/54	-

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	C	404	HEM	C4D-ND	-4.36	1.32	1.40
9	A	404	HEM	C1B-NB	-4.24	1.33	1.40
9	B	404	HEM	C4D-ND	-4.17	1.33	1.40
7	C	401	A1IUB	C6-C7	-3.85	1.45	1.56
7	A	401	A1IUB	C6-C7	-3.46	1.46	1.56

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	B	404	HEM	CHC-C4B-NB	5.40	130.29	124.43
9	A	404	HEM	C1B-NB-C4B	4.81	110.04	105.07
9	B	404	HEM	C1B-NB-C4B	4.80	110.03	105.07
9	A	404	HEM	CHC-C4B-NB	4.37	129.18	124.43
9	C	404	HEM	CBA-CAA-C2A	-4.32	105.25	112.62

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	B	401	A1IUB	C7-C8-C9-C10
8	A	403	GOL	C1-C2-C3-O3
8	A	403	GOL	O1-C1-C2-C3
8	A	403	GOL	O2-C2-C3-O3
7	A	401	A1IUB	O11-C10-C9-C8

There are no ring outliers.


9 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	403	A1IUB	7	0
7	A	402	A1IUB	2	0
7	B	401	A1IUB	2	0
12	B	403	ACT	2	0
9	C	404	HEM	6	0
7	A	401	A1IUB	5	0
9	B	404	HEM	4	0
7	C	401	A1IUB	11	0
9	A	404	HEM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand A1IUB A 402

















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/335 (100%)	-0.25	6 (1%) 67 64	17, 35, 58, 136	10 (2%)
1	B	335/335 (100%)	0.18	14 (4%) 41 37	19, 43, 74, 145	5 (1%)
1	C	335/335 (100%)	0.20	12 (3%) 46 43	18, 45, 73, 143	6 (1%)
All	All	1005/1005 (100%)	0.04	32 (3%) 50 47	17, 41, 71, 145	21 (2%)

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	348	LEU	4.6
1	A	14	GLN	4.5
1	B	15	GLY	4.1
1	B	316	ILE	4.0
1	B	14	GLN	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
11	SO4	B	406	5/5	0.73	0.12	89,93,97,98	0
8	GOL	C	402	6/6	0.74	0.20	69,80,83,83	0
11	SO4	C	406	5/5	0.76	0.11	81,84,97,104	0
7	A1IUB	C	403	14/14	0.77	0.35	54,69,80,81	14
11	SO4	A	406	5/5	0.77	0.12	81,82,94,94	0
7	A1IUB	B	401	14/14	0.79	0.32	58,75,82,88	0
7	A1IUB	A	401	14/14	0.80	0.28	57,71,84,86	14
7	A1IUB	A	402	14/14	0.80	0.34	65,70,79,84	14
8	GOL	A	403	6/6	0.81	0.21	54,59,66,73	0
7	A1IUB	C	401	14/14	0.82	0.34	73,78,88,99	14
12	ACT	B	403	4/4	0.83	0.24	45,52,53,54	0
8	GOL	B	402	6/6	0.84	0.18	54,57,61,67	0
10	MG	A	405	1/1	0.86	0.12	49,49,49,49	0
10	MG	C	405	1/1	0.89	0.11	52,52,52,52	0
10	MG	B	405	1/1	0.90	0.08	49,49,49,49	0
9	HEM	C	404	43/43	0.97	0.08	27,32,41,42	0
9	HEM	A	404	43/43	0.98	0.07	24,27,34,38	0
9	HEM	B	404	43/43	0.98	0.07	25,29,36,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1IUB C 403:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1IUB B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1IUB A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1IUB A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1IUB C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 404:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.