



wwPDB EM Validation Summary Report ⓘ

Nov 17, 2024 – 11:44 AM EST

PDB ID : 3J5M
EMDB ID : EMD-5779
Title : Cryo-EM structure of the BG505 SOSIP.664 HIV-1 Env trimer with 3 PGV04 Fabs
Authors : Lyumkis, D.; Julien, J.-P.; Wilson, I.A.; Ward, A.B.
Deposited on : 2013-10-26
Resolution : 5.80 Å (reported)
Based on initial models : 1ENV, 3SE9, 2B4C, 3U2S

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 5.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	
1	E	475	
1	I	475	
2	B	64	
2	F	64	
2	J	64	
3	C	208	
3	G	208	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	K	208	 <p>25% 96% 11% .</p>
4	D	228	 <p>24% 86% 11% ..</p>
4	H	228	 <p>24% 87% 11% ..</p>
4	L	228	 <p>25% 87% 11% ..</p>

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 20394 atoms, of which 15 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BG505 SOSIP gp120.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	415	3140	1964	554	595	27	0	0
1	E	415	3140	1964	554	595	27	0	0
1	I	415	3140	1964	554	595	27	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	332	ASN	THR	engineered mutation	UNP Q2N0S6
A	501	CYS	ALA	engineered mutation	UNP Q2N0S6
E	332	ASN	THR	engineered mutation	UNP Q2N0S6
E	501	CYS	ALA	engineered mutation	UNP Q2N0S6
I	332	ASN	THR	engineered mutation	UNP Q2N0S6
I	501	CYS	ALA	engineered mutation	UNP Q2N0S6

- Molecule 2 is a protein called BG505 SOSIP gp41.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	B	63	315	189	63	63	0	0
2	F	63	315	189	63	63	0	0
2	J	63	315	189	63	63	0	0

- Molecule 3 is a protein called PGV04 light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	208	1621	1018	275	321	7	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	208	Total	C	N	O	S	0	0
			1621	1018	275	321	7		
3	K	208	Total	C	N	O	S	0	0
			1621	1018	275	321	7		

- Molecule 4 is a protein called PGV04 heavy chain.


Mol	Chain	Residues	Atoms					AltConf	Trace	
4	D	225	Total	C	H	N	O	S	0	0
			1722	1090	5	298	324	5		
4	H	225	Total	C	H	N	O	S	0	0
			1722	1090	5	298	324	5		
4	L	225	Total	C	H	N	O	S	0	0
			1722	1090	5	298	324	5		




• Molecule 1: BG505 SOSIP gp120



• Molecule 2: BG505 SOSIP gp41




• Molecule 2: BG505 SOSIP gp41




• Molecule 2: BG505 SOSIP gp41







• Molecule 3: PGV04 light chain





• Molecule 3: PGV04 light chain



• Molecule 3: PGV04 light chain





• Molecule 4: PGV04 heavy chain







• Molecule 4: PGV04 heavy chain



• Molecule 4: PGV04 heavy chain



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C3	Depositor
Number of particles used	49572	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Frealign	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	32	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	29000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.067	Depositor
Minimum map value	-0.028	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.027	Depositor
Map size (\AA)	309.76, 309.76, 309.76	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.21, 1.21, 1.21	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/3206	0.57	1/4358 (0.0%)
1	E	0.60	0/3206	0.57	1/4358 (0.0%)
1	I	0.60	1/3206 (0.0%)	0.57	1/4358 (0.0%)
3	C	0.67	0/1658	0.56	1/2244 (0.0%)
3	G	0.67	0/1658	0.56	1/2244 (0.0%)
3	K	0.67	0/1658	0.56	1/2244 (0.0%)
4	D	0.58	2/1761 (0.1%)	0.57	2/2396 (0.1%)
4	H	0.58	2/1761 (0.1%)	0.57	2/2396 (0.1%)
4	L	0.58	2/1761 (0.1%)	0.57	2/2396 (0.1%)
All	All	0.61	7/19875 (0.0%)	0.57	12/26994 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	E	0	2
1	I	0	2
3	C	0	1
3	G	0	1
3	K	0	1
4	D	0	1
4	H	0	1
4	L	0	1
All	All	0	12

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	82(B)	GLY	C-N	5.29	1.46	1.34
4	D	82(B)	GLY	C-N	5.27	1.46	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	82(B)	GLY	C-N	5.27	1.46	1.34
4	D	149	PRO	N-CD	5.11	1.55	1.47
1	I	422	GLN	C-N	5.09	1.45	1.34

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	145	GLU	O-C-N	-12.75	102.31	122.70
1	A	145	GLU	O-C-N	-12.73	102.32	122.70
1	I	145	GLU	O-C-N	-12.73	102.33	122.70
3	G	91	LEU	O-C-N	-8.36	109.33	122.70
3	C	91	LEU	O-C-N	-8.34	109.36	122.70

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	145	GLU	Mainchain
1	A	47	ASP	Peptide
3	C	91	LEU	Mainchain
4	D	35(A)	HIS	Mainchain
1	E	47	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3140	0	2943	90	0
1	E	3140	0	2943	93	0
1	I	3140	0	2943	90	0
2	B	315	0	67	0	0
2	F	315	0	67	0	0
2	J	315	0	67	0	0
3	C	1621	0	1579	6	0
3	G	1621	0	1579	5	0
3	K	1621	0	1579	5	0
4	D	1717	5	1690	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	1717	5	1690	25	0
4	L	1717	5	1690	23	0
All	All	20379	15	18837	361	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 361 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:TYR:CE1	1:A:421:LYS:HG2	1.30	1.66
1:E:384:TYR:CE1	1:E:421:LYS:HG2	1.30	1.63
1:I:384:TYR:CE1	1:I:421:LYS:HG2	1.30	1.60
1:E:384:TYR:CE1	1:E:421:LYS:CG	1.89	1.55
1:I:384:TYR:CE1	1:I:421:LYS:CG	1.89	1.54

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/475 (86%)	359 (88%)	33 (8%)	17 (4%)	2	17
1	E	409/475 (86%)	359 (88%)	33 (8%)	17 (4%)	2	17
1	I	409/475 (86%)	359 (88%)	33 (8%)	17 (4%)	2	17
3	C	206/208 (99%)	198 (96%)	7 (3%)	1 (0%)	25	65
3	G	206/208 (99%)	198 (96%)	7 (3%)	1 (0%)	25	65
3	K	206/208 (99%)	198 (96%)	7 (3%)	1 (0%)	25	65
4	D	221/228 (97%)	212 (96%)	9 (4%)	0	100	100
4	H	221/228 (97%)	212 (96%)	9 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	L	221/228 (97%)	212 (96%)	9 (4%)	0	100	100
All	All	2508/2733 (92%)	2307 (92%)	147 (6%)	54 (2%)	8	29

5 of 54 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	LEU
1	A	302	ASN
1	A	318	ALA
1	A	321	ASP
1	A	323	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/422 (79%)	330 (99%)	3 (1%)	75	83
1	E	333/422 (79%)	330 (99%)	3 (1%)	75	83
1	I	333/422 (79%)	330 (99%)	3 (1%)	75	83
3	C	182/182 (100%)	181 (100%)	1 (0%)	86	89
3	G	182/182 (100%)	181 (100%)	1 (0%)	86	89
3	K	182/182 (100%)	181 (100%)	1 (0%)	86	89
4	D	190/193 (98%)	190 (100%)	0	100	100
4	H	190/193 (98%)	190 (100%)	0	100	100
4	L	190/193 (98%)	190 (100%)	0	100	100
All	All	2115/2391 (88%)	2103 (99%)	12 (1%)	82	88

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	91	LEU
1	I	111	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	K	91	LEU
1	I	412	ASP
3	C	91	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
4	H	171	GLN
1	I	67	ASN
4	L	171	GLN
4	D	171	GLN
1	A	67	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-5779. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 136



Y Index: 135




Z Index: 121


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.027. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 207 nm³; this corresponds to an approximate mass of 187 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.172 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-5779 and PDB model 3J5M. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.027 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.027).

9.4 Atom inclusion [i](#)



At the recommended contour level, 64% of all backbone atoms, 60% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.027) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6010	 0.1560
A	 0.6150	 0.1590
B	 0.7810	 0.2100
C	 0.5790	 0.1340
D	 0.5780	 0.1590
E	 0.6150	 0.1590
F	 0.7870	 0.2110
G	 0.5770	 0.1340
H	 0.5830	 0.1620
I	 0.6170	 0.1610
J	 0.7870	 0.2080
K	 0.5740	 0.1350
L	 0.5800	 0.1590

