



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 19, 2025 – 12:42 PM EDT

PDB ID : 4JE5
Title : Crystal structure of the aromatic aminotransferase Aro8, a putative alpha-amino acid aminotransferase in *Saccharomyces cerevisiae*
Authors : Bulfer, S.L.; Brunzelle, J.S.; Trievel, R.C.
Deposited on : 2013-02-26
Resolution : 1.91 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>.</div> </div> </div>
1	B	503	<div> <div>12%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>..</div> </div> </div>
1	C	503	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>
2	D	503	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>..</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 16855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aromatic/aminoadipate aminotransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	0	26	0
			3978	2568	644	758	8			
1	B	497	Total	C	N	O	S	0	16	0
			3882	2511	630	732	9			
1	C	492	Total	C	N	O	S	0	18	0
			3899	2508	636	746	9			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P53090
A	-1	HIS	-	expression tag	UNP P53090
A	0	MET	-	expression tag	UNP P53090
B	-2	GLY	-	expression tag	UNP P53090
B	-1	HIS	-	expression tag	UNP P53090
B	0	MET	-	expression tag	UNP P53090
C	-2	GLY	-	expression tag	UNP P53090
C	-1	HIS	-	expression tag	UNP P53090
C	0	MET	-	expression tag	UNP P53090


- Molecule 2 is a protein called Aromatic/aminoadipate aminotransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	496	Total	C	N	O	P S	0	15	0
			3961	2545	645	761	1 9			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	GLY	-	expression tag	UNP P53090
D	-1	HIS	-	expression tag	UNP P53090
D	0	MET	-	expression tag	UNP P53090

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is 4'-DEOXY-4'-AMINOPYRIDOXAL-5'-PHOSPHATE (three-letter code: PMP) (formula: C₈H₁₃N₂O₅P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			16	8	2	5	1		
5	C	1	Total	C	N	O	P	0	0
			16	8	2	5	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	280	Total	O	0	0
			280	280		
6	B	213	Total	O	0	0
			213	213		
6	C	264	Total	O	0	0
			264	264		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	256	Total 256	O 256	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aromatic/aminoadipate aminotransferase 1





- Molecule 1: Aromatic/aminoadipate aminotransferase 1



- Molecule 1: Aromatic/aminoadipate aminotransferase 1



- Molecule 2: Aromatic/aminoadipate aminotransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	227.44Å 159.83Å 70.64Å 90.00° 106.03° 90.00°	Depositor
Resolution (Å)	34.25 – 1.91 34.25 – 1.91	Depositor EDS
% Data completeness (in resolution range)	96.6 (34.25-1.91) 96.5 (34.25-1.91)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.182 , 0.217 0.186 , 0.221	Depositor DCC
R_{free} test set	9078 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16855	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PMP, PLP, LLP, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/4149	0.65	0/5656
1	B	0.64	0/4041	0.65	1/5525 (0.0%)
1	C	0.69	0/4049	0.67	0/5527
2	D	0.69	0/4074	0.68	2/5566 (0.0%)
All	All	0.68	0/16313	0.66	3/22274 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	312[A]	ARG	NE-CZ-NH2	-5.25	117.67	120.30
2	D	312[B]	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	B	365	ARG	NE-CZ-NH2	-5.22	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3978	0	3889	27	0
1	B	3882	0	3720	35	0
1	C	3899	0	3744	27	0
2	D	3961	0	3784	35	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	15	0	6	1	0
4	A	30	0	34	2	0
4	B	15	0	17	2	0
4	C	30	0	34	3	0
5	B	16	0	11	5	0
5	C	16	0	11	3	0
6	A	280	0	0	1	0
6	B	213	0	0	1	0
6	C	264	0	0	3	0
6	D	256	0	0	1	0
All	All	16855	0	15250	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 123 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:321:LYS:HA	1:A:324[A]:LYS:HE3	1.47	0.94
1:B:251:TYR:HH	5:B:1000:PMP:HO3	1.29	0.79
1:C:267:GLU:HA	1:C:270[B]:GLN:HG2	1.65	0.77
1:C:305:LYS:NZ	5:C:1000:PMP:H4A1	2.00	0.76
1:C:311:THR:O	1:C:341[B]:THR:HG21	1.85	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/503 (103%)	502 (97%)	14 (3%)	0	100	100
1	B	511/503 (102%)	491 (96%)	19 (4%)	1 (0%)	44	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	506/503 (101%)	492 (97%)	14 (3%)	0	100	100
2	D	508/503 (101%)	494 (97%)	12 (2%)	2 (0%)	30	22
All	All	2041/2012 (101%)	1979 (97%)	59 (3%)	3 (0%)	44	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	414	THR
2	D	335	GLN
1	B	464	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	436/439 (99%)	431 (99%)	5 (1%)	70	71
1	B	411/439 (94%)	407 (99%)	4 (1%)	73	74
1	C	420/439 (96%)	413 (98%)	7 (2%)	56	54
2	D	425/438 (97%)	423 (100%)	2 (0%)	86	88
All	All	1692/1755 (96%)	1674 (99%)	18 (1%)	75	71

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	451[A]	GLU
2	D	90	SER
2	D	48	ASP
1	B	488	LEU
1	C	341[B]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LLP	D	305	2	23,24,25	1.95	5 (21%)	25,32,34	1.66	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LLP	D	305	2	-	5/16/17/19	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	305	LLP	O3-C3	-5.52	1.24	1.36
2	D	305	LLP	C2-N1	3.79	1.40	1.33
2	D	305	LLP	C4-C4'	3.12	1.53	1.46
2	D	305	LLP	C4'-NZ	2.61	1.35	1.27
2	D	305	LLP	C6-N1	2.31	1.39	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	305	LLP	OP4-C5'-C5	3.90	116.66	109.36
2	D	305	LLP	C4-C4'-NZ	-3.45	108.10	124.04
2	D	305	LLP	OP2-P-OP4	-3.29	98.09	106.67
2	D	305	LLP	OP3-P-OP2	2.90	118.68	107.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	305	LLP	C5-C6-N1	-2.03	120.53	123.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	305	LLP	C5'-OP4-P-OP3
2	D	305	LLP	CA-CB-CG-CD
2	D	305	LLP	C3-C4-C4'-NZ
2	D	305	LLP	CD-CE-NZ-C4'
2	D	305	LLP	C5-C4-C4'-NZ

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	305	LLP	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PMP	C	1000	-	16,16,16	1.14	1 (6%)	22,23,23	1.30	2 (9%)
3	PLP	A	601	-	15,15,16	1.14	1 (6%)	21,22,23	1.23	2 (9%)
5	PMP	B	1000	-	16,16,16	1.05	1 (6%)	22,23,23	1.24	2 (9%)
4	EPE	C	1002	-	15,15,15	0.85	1 (6%)	19,20,20	2.04	5 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	A	603	-	15,15,15	1.00	1 (6%)	19,20,20	2.09	6 (31%)
4	EPE	C	1001	-	15,15,15	0.87	1 (6%)	19,20,20	1.86	4 (21%)
4	EPE	B	604	-	15,15,15	0.85	1 (6%)	19,20,20	2.00	5 (26%)
4	EPE	A	602	-	15,15,15	0.92	1 (6%)	19,20,20	1.98	5 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PMP	C	1000	-	-	2/8/8/8	0/1/1/1
3	PLP	A	601	-	-	0/6/6/8	0/1/1/1
5	PMP	B	1000	-	-	4/8/8/8	0/1/1/1
4	EPE	C	1002	-	-	4/9/19/19	0/1/1/1
4	EPE	A	603	-	-	2/9/19/19	0/1/1/1
4	EPE	C	1001	-	-	6/9/19/19	0/1/1/1
4	EPE	B	604	-	-	3/9/19/19	0/1/1/1
4	EPE	A	602	-	-	1/9/19/19	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	603	EPE	C10-S	3.27	1.82	1.77
4	B	604	EPE	C10-S	3.02	1.81	1.77
4	A	602	EPE	C10-S	2.93	1.81	1.77
5	B	1000	PMP	C2-N1	2.92	1.39	1.33
4	C	1002	EPE	C10-S	2.85	1.81	1.77

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	604	EPE	C5-N4-C3	5.90	121.55	108.84
4	C	1001	EPE	C5-N4-C3	5.79	121.31	108.84
4	A	602	EPE	C5-N4-C3	5.16	119.95	108.84
4	C	1002	EPE	C5-N4-C3	4.94	119.48	108.84
4	A	603	EPE	C5-N4-C3	4.72	119.00	108.84

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	602	EPE	C8-C7-N4-C3
4	B	604	EPE	C10-C9-N1-C6
4	B	604	EPE	C8-C7-N4-C5
4	C	1001	EPE	C10-C9-N1-C2
5	B	1000	PMP	C3-C4-C4A-N4A

There are no ring outliers.

8 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1000	PMP	3	0
3	A	601	PLP	1	0
5	B	1000	PMP	5	0
4	C	1002	EPE	2	0
4	A	603	EPE	1	0
4	C	1001	EPE	1	0
4	B	604	EPE	2	0
4	A	602	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	494/503 (98%)	-0.24	18 (3%)	46	48	8, 23, 45, 69	28 (5%)
1	B	497/503 (98%)	0.28	61 (12%)	9	10	10, 29, 66, 80	16 (3%)
1	C	492/503 (97%)	-0.16	17 (3%)	47	49	9, 24, 49, 67	18 (3%)
2	D	495/503 (98%)	-0.11	20 (4%)	43	44	8, 24, 54, 65	15 (3%)
All	All	1978/2012 (98%)	-0.06	116 (5%)	29	30	8, 25, 57, 80	77 (3%)

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	PHE	6.0
1	B	498	ILE	5.9
1	B	32	PHE	5.6
1	C	28	CYS	5.6
1	B	88	ASP	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LLP	D	305	24/25	0.92	0.12	16,32,38,39	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EPE	C	1002	15/15	0.80	0.15	44,58,75,75	0
4	EPE	C	1001	15/15	0.87	0.17	42,58,67,67	0
4	EPE	A	603	15/15	0.87	0.13	47,56,69,70	0
3	PLP	A	601	15/16	0.89	0.12	31,38,41,41	0
4	EPE	B	604	15/15	0.90	0.19	32,34,39,40	11
5	PMP	C	1000	16/16	0.91	0.11	24,39,44,47	0
4	EPE	A	602	15/15	0.94	0.11	35,40,50,51	0
5	PMP	B	1000	16/16	0.95	0.09	31,40,43,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.