



# wwPDB NMR Structure Validation Summary Report ⓘ

Jun 24, 2024 – 01:24 PM EDT

PDB ID : 7JXG  
BMRB ID : 50269  
Title : Structural model for Fe-containing human acireductone dioxygenase  
Authors : Pochapsky, T.C.; Liu, X.; Deshpande, A.; Ringe, D.; Garber, A.; Ryan, J.  
Deposited on : 2020-08-27

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*


The overall completeness of chemical shifts assignment is 62%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	179	

## 2 Ensemble composition and analysis

This entry contains 6 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:7, A:18-A:159 (148)	1.07	6
2	A:171-A:178 (8)	0.14	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	3, 4, 5, 6
2	1, 2

### 3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2972 atoms, of which 1457 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called 1,2-dihydroxy-3-keto-5-methylthiopentene dioxygenase.

Mol	Chain	Residues	Atoms						Trace
1	A	178	Total	C	H	N	O	S	0
			2969	968	1457	261	275	8	

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms	
2	A	1	Total	Fe
			1	1

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	
3	A	2	Total	O
			2	2

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: 1,2-dihydroxy-3-keto-5-methylthiopentene dioxygenase



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 6. Colouring as in section 4.1 above.

- Molecule 1: 1,2-dihydroxy-3-keto-5-methylthiopentene dioxygenase



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 6 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure calculation	2.51

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1567
Number of shifts mapped to atoms	1559
Number of unparsed shifts	0
Number of shifts with mapping errors	8
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	62%

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1340	1303	1302	263±27
All	All	8058	7818	7812	1579

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 99.

5 of 1144 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:99:LEU:HD11	1:A:146:MET:SD	1.09	1.86	1	2
1:A:4:ALA:HB3	1:A:26:VAL:HG11	1.09	1.15	4	5
1:A:89:LEU:HD22	1:A:132:TYR:CE2	1.07	1.84	4	2
1:A:4:ALA:HB3	1:A:26:VAL:CG1	1.06	1.80	3	5
1:A:89:LEU:HD11	1:A:130:GLY:C	1.04	1.72	1	2

## 6.3 Torsion angles

### 6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/179 (87%)	145±2 (94±1%)	8±2 (5±2%)	2±1 (2±0%)	14	59
All	All	930/1074 (87%)	871 (94%)	45 (5%)	14 (2%)	14	59

5 of 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	50	PRO	5
1	A	137	VAL	3
1	A	153	PRO	2
1	A	75	PRO	2
1	A	141	ASN	1

### 6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/159 (89%)	123±3 (87±2%)	19±3 (13±2%)	7	48
All	All	852/954 (89%)	739 (87%)	113 (13%)	7	48

5 of 54 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	46	TYR	6
1	A	19	ARG	5
1	A	74	LEU	5
1	A	5	TRP	4
1	A	24	ARG	4



### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 62% for the well-defined parts and 60% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: *assigned\_chemical\_shifts\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1567
Number of shifts mapped to atoms	1559
Number of unparsed shifts	0
Number of shifts with mapping errors	8
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	33

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. First 5 (of 8) occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	179	ALA	H	7.66	0.01	1
1	A	179	ALA	HA	3.87	0.01	1
1	A	179	ALA	HB1	1.16	0.01	1
1	A	179	ALA	HB2	1.16	0.01	1
1	A	179	ALA	HB3	1.16	0.01	1
1	A	179	ALA	CA	52.77	0.01	1
1	A	179	ALA	CB	19.6	0.01	1
1	A	179	ALA	N	131.3	0.1	1

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	163	$-0.08 \pm 0.14$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	145	$0.30 \pm 0.12$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	149	$-0.04 \pm 0.48$	None needed ( $< 0.5$ ppm)

### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 62%, i.e. 1413 atoms were assigned a chemical shift out of a possible 2294. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	558/774 (72%)	284/313 (91%)	141/312 (45%)	133/149 (89%)
Sidechain	809/1263 (64%)	570/812 (70%)	239/394 (61%)	0/57 (0%)
Aromatic	46/257 (18%)	42/124 (34%)	0/123 (0%)	4/10 (40%)
Overall	1413/2294 (62%)	896/1249 (72%)	380/829 (46%)	137/216 (63%)

### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	126	THR	HG21	-0.78	0.08 – 2.19	-9.1
1	A	126	THR	HG22	-0.78	0.08 – 2.19	-9.1
1	A	126	THR	HG23	-0.78	0.08 – 2.19	-9.1
1	A	19	ARG	HD2	1.08	1.97 – 4.26	-8.9
1	A	83	MET	HB2	-0.82	0.42 – 3.63	-8.9
1	A	153	PRO	HA	1.74	2.78 – 6.00	-8.2
1	A	83	MET	HG2	-0.43	0.65 – 4.19	-8.1
1	A	110	LYS	CG	33.62	19.35 – 30.45	7.9
1	A	80	LYS	HB2	-0.02	0.58 – 2.97	-7.5
1	A	59	ARG	HG2	-0.35	0.26 – 2.87	-7.3
1	A	83	MET	HG3	-0.23	0.54 – 4.26	-7.1
1	A	19	ARG	HG3	-0.33	0.15 – 2.94	-6.7
1	A	19	ARG	CD	49.34	38.57 – 47.75	6.7
1	A	19	ARG	HD3	1.38	1.81 – 4.39	-6.7
1	A	20	PRO	CD	57.10	45.11 – 55.58	6.5
1	A	125	VAL	HG11	-0.82	-0.48 – 2.12	-6.3
1	A	125	VAL	HG12	-0.82	-0.48 – 2.12	-6.3

*Continued on next page...*

Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	125	VAL	HG13	-0.82	-0.48 – 2.12	-6.3
1	A	19	ARG	HG2	-0.06	0.26 – 2.87	-6.2
1	A	83	MET	HA	1.65	2.11 – 6.67	-6.0
1	A	83	MET	HE1	-0.34	-0.03 – 3.80	-5.8
1	A	83	MET	HE2	-0.34	-0.03 – 3.80	-5.8
1	A	83	MET	HE3	-0.34	-0.03 – 3.80	-5.8
1	A	83	MET	HB3	0.10	0.33 – 3.66	-5.7
1	A	153	PRO	HB2	0.15	0.37 – 3.78	-5.7
1	A	126	THR	HB	2.40	2.57 – 5.77	-5.5
1	A	19	ARG	HB2	0.41	0.52 – 3.08	-5.4
1	A	6	TYR	HE1	5.50	5.59 – 7.82	-5.4
1	A	37	LEU	HB2	-0.18	-0.07 – 3.30	-5.3
1	A	26	VAL	HG21	-0.65	-0.58 – 2.19	-5.3
1	A	26	VAL	HG22	-0.65	-0.58 – 2.19	-5.3
1	A	26	VAL	HG23	-0.65	-0.58 – 2.19	-5.3
1	A	153	PRO	HB3	0.19	0.25 – 3.76	-5.2

### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

