



# wwPDB X-ray Structure Validation Summary Report i

Nov 17, 2024 – 04:03 PM EST

PDB ID : 4KYI  
Title : Crystal structure of the phospholipase VipD from Legionella pneumophila in complex with the human GTPase Rab5  
Authors : Lucas, M.; Gaspar, A.H.; Pallara, C.; Rojas, A.L.; Fernandez-Recio, J.; Machner, M.P.; Hierro, A.  
Deposited on : 2013-05-29  
Resolution : 3.08 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriaage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1842 (3.10-3.06)
Clashscore	180529	1965 (3.10-3.06)
Ramachandran outliers	177936	1859 (3.10-3.06)
Sidechain outliers	177891	1858 (3.10-3.06)
RSRZ outliers	164620	1842 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain		
2	B	170		80%	16% •
2	D	170		79%	16% ••
2	F	170		85%	11% •
2	H	170		84%	12% ••

## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 21833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VipD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	530	Total	C 4140	N 2608	O 708	S 807	17	0	0
1	C	527	Total	C 4120	N 2596	O 705	S 802	17	0	0
1	E	531	Total	C 4145	N 2609	O 709	S 810	17	0	0
1	G	528	Total	C 4122	N 2594	O 705	S 806	17	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	GLY	-	expression tag	UNP Q5ZRP9
A	15	ALA	-	expression tag	UNP Q5ZRP9
A	16	MET	-	expression tag	UNP Q5ZRP9
A	17	GLY	-	expression tag	UNP Q5ZRP9
A	18	SER	-	expression tag	UNP Q5ZRP9
C	14	GLY	-	expression tag	UNP Q5ZRP9
C	15	ALA	-	expression tag	UNP Q5ZRP9
C	16	MET	-	expression tag	UNP Q5ZRP9
C	17	GLY	-	expression tag	UNP Q5ZRP9
C	18	SER	-	expression tag	UNP Q5ZRP9
E	14	GLY	-	expression tag	UNP Q5ZRP9
E	15	ALA	-	expression tag	UNP Q5ZRP9
E	16	MET	-	expression tag	UNP Q5ZRP9
E	17	GLY	-	expression tag	UNP Q5ZRP9
E	18	SER	-	expression tag	UNP Q5ZRP9
G	14	GLY	-	expression tag	UNP Q5ZRP9
G	15	ALA	-	expression tag	UNP Q5ZRP9
G	16	MET	-	expression tag	UNP Q5ZRP9
G	17	GLY	-	expression tag	UNP Q5ZRP9
G	18	SER	-	expression tag	UNP Q5ZRP9


- Molecule 2 is a protein called Ras-related protein Rab-5C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	164	Total	C	N	O	S	0	0	0
			1283	820	215	242	6			
2	D	164	Total	C	N	O	S	0	0	0
			1283	820	215	242	6			
2	F	164	Total	C	N	O	S	0	0	0
			1283	820	215	242	6			
2	H	164	Total	C	N	O	S	0	0	0
			1283	820	215	242	6			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	13	GLY	-	expression tag	UNP P51148
B	14	ALA	-	expression tag	UNP P51148
B	15	MET	-	expression tag	UNP P51148
B	16	GLY	-	expression tag	UNP P51148
B	17	SER	-	expression tag	UNP P51148
B	80	LEU	GLN	engineered mutation	UNP P51148
D	13	GLY	-	expression tag	UNP P51148
D	14	ALA	-	expression tag	UNP P51148
D	15	MET	-	expression tag	UNP P51148
D	16	GLY	-	expression tag	UNP P51148
D	17	SER	-	expression tag	UNP P51148
D	80	LEU	GLN	engineered mutation	UNP P51148
F	13	GLY	-	expression tag	UNP P51148
F	14	ALA	-	expression tag	UNP P51148
F	15	MET	-	expression tag	UNP P51148
F	16	GLY	-	expression tag	UNP P51148
F	17	SER	-	expression tag	UNP P51148
F	80	LEU	GLN	engineered mutation	UNP P51148
H	13	GLY	-	expression tag	UNP P51148
H	14	ALA	-	expression tag	UNP P51148
H	15	MET	-	expression tag	UNP P51148
H	16	GLY	-	expression tag	UNP P51148
H	17	SER	-	expression tag	UNP P51148
H	80	LEU	GLN	engineered mutation	UNP P51148

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	E	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	E	1	Total C O 6 3 3	0	0

- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
5	D	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
5	F	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
5	H	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		
6	F	1	Total	Mg	0	0
			1	1		
6	H	1	Total	Mg	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: VipD

Chain A:  78% 17% . .

 E209 T214 Y28 S29 V30 V39 I48 L51 Q55 Q60 R60 I88 M106 F111 K245 D254 S260 I261 V264 H271 P272 F116 D116 R119 D123 M127 K31 I141 P142


 E378 V379 T380 T381 N383 S385 P386 T387 Q390 Q401 T402 M403 K396 D400 Q411 T412 H415 P416 N280 A281 V309 E312 R320 I332 R338 L342 V343 I156 A157 L158 Y159 M171 I177 ILE GLY LYS TRP LEU HIS ALA PRO GLN PRO GLU PRO

• Molecule 1: VipD

Chain C:  74% 20% . .


 L179 K184 K188 A192 L193 H194 S195 A196 T204 V210 P211 L215 G216 D217 L218 R222 D90 I191 K92 K93 L94 T230 V238 D239 L248 Y251 S252 D254 D116 N120 D112 N125 D123 Y124 Y125 D126 K131 I141 P142

 L179 K184 K188 A192 L193 H194 S195 A196 T204 V210 P211 L215 G216 D217 L218 R222 D90 I191 K92 K93 L94 T230 V238 D239 L248 Y251 S252 D254 D116 N120 D112 N125 D123 Y124 Y125 D126 K131 I141 P142

 S437 E438 K439 E440 H447 A448 Q449 L454 Q457 I458 E461 D464 S469 D479 A485 K486 M487 D488 D489 T490 Q491 F496 Q498 Q499 R493 D496 Y499 D499 K500 G502 L498 F505 K507 P502 L496 R506 H415 P416 R503 T497 D288 G278 E304 D305 P416 L417 K310 A311 E312 A313 Y421 Y422 I423 G424 R320 L428 R338 D430 E435 K436


- Molecule 1: VipD

Chain E:  75% 19% . .




- Molecule 1: VipD

Chain G: 75% 19% • •




- Molecule 2: Ras-related protein Rab-5C

Chain B: 80% 16% •




- Molecule 2: Ras-related protein Rab-5C



Chain D: 79% 16% • •



- Molecule 2: Ras-related protein Rab-5C



- Molecule 2: Ras-related protein Rab-5C



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.28Å 97.98Å 109.85Å 76.57° 80.71° 78.91°	Depositor
Resolution (Å)	29.83 – 3.08 29.83 – 3.08	Depositor EDS
% Data completeness (in resolution range)	97.6 (29.83-3.08) 97.5 (29.83-3.08)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.02 (at 3.06Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
$R$ , $R_{free}$	0.233 , 0.280 0.235 , 0.282	Depositor DCC
$R_{free}$ test set	3408 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	85.7	Xtriage
Anisotropy	0.370	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 23.6	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	21833	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, GNP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/4202	0.39	0/5668
1	C	0.22	0/4182	0.38	0/5640
1	E	0.22	0/4205	0.39	0/5671
1	G	0.21	0/4182	0.38	0/5640
2	B	0.23	0/1305	0.39	0/1765
2	D	0.22	0/1305	0.38	0/1765
2	F	0.22	0/1305	0.37	0/1765
2	H	0.23	0/1305	0.39	0/1765
All	All	0.22	0/21991	0.38	0/29679

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4140	0	4187	45	0
1	C	4120	0	4166	65	0
1	E	4145	0	4192	61	0
1	G	4122	0	4163	58	0
2	B	1283	0	1282	18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1283	0	1282	21	0
2	F	1283	0	1282	15	0
2	H	1283	0	1282	15	0
3	A	8	0	12	1	0
3	B	8	0	12	1	0
3	E	4	0	6	0	0
3	G	4	0	6	2	0
4	A	12	0	16	0	0
4	E	6	0	8	1	0
5	B	32	0	13	0	0
5	D	32	0	13	4	0
5	F	32	0	13	1	0
5	H	32	0	13	1	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
6	H	1	0	0	0	0
All	All	21833	0	21948	279	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 279 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:112:ARG:HD3	1:E:276:VAL:HG12	1.68	0.75
1:C:312:GLU:OE2	1:C:320:ARG:NH1	2.22	0.72
1:C:58:GLN:HG2	1:C:88:ILE:HD12	1.72	0.72
1:E:60:ARG:HG3	1:E:406:LEU:HD23	1.74	0.68
1:E:112:ARG:NH2	1:E:272:PRO:O	2.27	0.68

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	526/551 (96%)	508 (97%)	16 (3%)	2 (0%)	30 60
1	C	523/551 (95%)	505 (97%)	17 (3%)	1 (0%)	44 72
1	E	525/551 (95%)	503 (96%)	20 (4%)	2 (0%)	30 60
1	G	522/551 (95%)	499 (96%)	22 (4%)	1 (0%)	44 72
2	B	162/170 (95%)	152 (94%)	10 (6%)	0	100 100
2	D	162/170 (95%)	152 (94%)	8 (5%)	2 (1%)	11 35
2	F	162/170 (95%)	151 (93%)	11 (7%)	0	100 100
2	H	162/170 (95%)	149 (92%)	13 (8%)	0	100 100
All	All	2744/2884 (95%)	2619 (95%)	117 (4%)	8 (0%)	37 66

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	382	THR
1	G	106	ASN
1	A	553	ALA
2	D	92	ARG
1	E	552	PRO

### 5.3.2 Protein sidechains [\(1\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	451/466 (97%)	425 (94%)	26 (6%)	17 43
1	C	448/466 (96%)	413 (92%)	35 (8%)	10 33
1	E	452/466 (97%)	417 (92%)	35 (8%)	10 33
1	G	449/466 (96%)	422 (94%)	27 (6%)	16 42
2	B	134/137 (98%)	132 (98%)	2 (2%)	60 78
2	D	134/137 (98%)	132 (98%)	2 (2%)	60 78

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	F	134/137 (98%)	132 (98%)	2 (2%)	60 78
2	H	134/137 (98%)	130 (97%)	4 (3%)	36 62
All	All	2336/2412 (97%)	2203 (94%)	133 (6%)	17 43

5 of 133 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	376	THR
1	G	405	LEU
2	H	54	ILE
1	C	403	MET
1	C	393	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	139	GLN
1	G	411	GLN
2	D	95	GLN
1	E	401	GLN
1	E	411	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 17 ligands modelled in this entry, 4 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	GNP	D	301	6	29,34,34	2.19	8 (27%)	33,54,54	2.37	7 (21%)
3	EDO	E	601	-	3,3,3	0.42	0	2,2,2	0.38	0
4	GOL	E	602	-	5,5,5	0.38	0	5,5,5	0.32	0
4	GOL	A	604	-	5,5,5	0.38	0	5,5,5	0.33	0
3	EDO	A	602	-	3,3,3	0.42	0	2,2,2	0.37	0
3	EDO	G	601	-	3,3,3	0.41	0	2,2,2	0.39	0
3	EDO	B	303	-	3,3,3	0.42	0	2,2,2	0.43	0
4	GOL	A	603	-	5,5,5	0.37	0	5,5,5	0.27	0
5	GNP	H	301	6	29,34,34	2.22	8 (27%)	33,54,54	2.40	7 (21%)
5	GNP	F	301	6	29,34,34	2.18	8 (27%)	33,54,54	2.38	7 (21%)
3	EDO	A	601	-	3,3,3	0.43	0	2,2,2	0.41	0
5	GNP	B	301	6	29,34,34	2.18	8 (27%)	33,54,54	2.38	7 (21%)
3	EDO	B	302	-	3,3,3	0.41	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GNP	D	301	6	-	4/14/38/38	0/3/3/3
3	EDO	E	601	-	-	0/1/1/1	-
4	GOL	E	602	-	-	2/4/4/4	-
4	GOL	A	604	-	-	2/4/4/4	-
3	EDO	A	602	-	-	0/1/1/1	-
3	EDO	G	601	-	-	0/1/1/1	-
3	EDO	B	303	-	-	0/1/1/1	-
4	GOL	A	603	-	-	2/4/4/4	-
5	GNP	H	301	6	-	2/14/38/38	0/3/3/3
5	GNP	F	301	6	-	7/14/38/38	0/3/3/3
3	EDO	A	601	-	-	0/1/1/1	-
5	GNP	B	301	6	-	4/14/38/38	0/3/3/3
3	EDO	B	302	-	-	0/1/1/1	-

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	301	GNP	PA-O3A	-6.68	1.52	1.59
5	D	301	GNP	PA-O3A	-6.54	1.52	1.59
5	B	301	GNP	PA-O3A	-6.53	1.52	1.59
5	F	301	GNP	PA-O3A	-6.50	1.52	1.59
5	H	301	GNP	PB-O3A	-5.48	1.52	1.59

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	301	GNP	C5-C6-N1	-8.79	111.67	123.42
5	D	301	GNP	C5-C6-N1	-8.76	111.71	123.42
5	B	301	GNP	C5-C6-N1	-8.72	111.76	123.42
5	F	301	GNP	C5-C6-N1	-8.61	111.91	123.42
5	H	301	GNP	C2-N1-C6	6.35	124.79	115.96

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	603	GOL	O1-C1-C2-O2
4	A	603	GOL	O1-C1-C2-C3
4	A	604	GOL	O1-C1-C2-C3
5	B	301	GNP	PG-N3B-PB-O1B
5	B	301	GNP	PA-O3A-PB-O2B


There are no ring outliers.



7 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	301	GNP	4	0
4	E	602	GOL	1	0
3	A	602	EDO	1	0
3	G	601	EDO	2	0
3	B	303	EDO	1	0
5	H	301	GNP	1	0
5	F	301	GNP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	530/551 (96%)	-0.68	0   100   100	1, 36, 77, 137	0
1	C	527/551 (95%)	-0.53	0   100   100	5, 57, 109, 169	0
1	E	531/551 (96%)	-0.67	0   100   100	0, 32, 79, 118	0
1	G	528/551 (95%)	-0.42	2 (0%)   89   77	14, 59, 112, 142	0
2	B	164/170 (96%)	-0.47	0   100   100	16, 58, 89, 151	0
2	D	164/170 (96%)	-0.45	0   100   100	25, 73, 108, 133	0
2	F	164/170 (96%)	-0.27	1 (0%)   85   72	22, 72, 108, 149	0
2	H	164/170 (96%)	-0.54	0   100   100	26, 60, 99, 130	0
All	All	2772/2884 (96%)	-0.54	3 (0%)   92   87	0, 52, 102, 169	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	536	ASN	3.1
1	G	358	VAL	2.5
2	F	101	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	602	4/4	0.48	0.11	55,55,55,55	0
3	EDO	G	601	4/4	0.64	0.20	84,84,84,84	0
4	GOL	A	604	6/6	0.64	0.17	78,78,78,78	0
3	EDO	A	601	4/4	0.67	0.23	58,58,58,58	0
4	GOL	E	602	6/6	0.68	0.15	66,66,66,66	0
3	EDO	E	601	4/4	0.75	0.08	56,56,56,56	0
4	GOL	A	603	6/6	0.79	0.13	59,59,59,59	0
3	EDO	B	302	4/4	0.83	0.18	45,45,45,45	0
3	EDO	B	303	4/4	0.88	0.17	41,41,41,41	0
5	GNP	F	301	32/32	0.89	0.08	81,81,81,81	0
5	GNP	B	301	32/32	0.90	0.08	72,72,72,72	0
6	MG	F	302	1/1	0.90	0.12	54,54,54,54	0
5	GNP	D	301	32/32	0.93	0.05	80,80,80,80	0
5	GNP	H	301	32/32	0.94	0.06	63,63,63,63	0
6	MG	B	304	1/1	0.95	0.07	36,36,36,36	0
6	MG	H	302	1/1	0.98	0.06	35,35,35,35	0
6	MG	D	302	1/1	1.00	0.02	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.