



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 08:30 PM EDT

PDB ID : 5LHI  
Title : Structure of the KDM1A/CoREST complex with the inhibitor N-[3-(ethoxymethyl)-2-[[4-[(3R)-pyrrolidin-3-yl]methoxy]phenoxy]methyl]phenyl]-4-methylthieno[3,2-b]pyrrole-5-carboxamide  
Authors : Cecatiello, V.; Pasqualato, S.  
Deposited on : 2016-07-12  
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	852	<div> <div>%</div> <div> <div></div> <div>47%</div> <div>29%</div> <div>•</div> <div>22%</div> </div> </div>
2	B	482	<div> <div>15%</div> <div>13%</div> <div>72%</div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LYSINE-SPECIFIC HISTONE DEMETHYLASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	666	Total	C	N	O	S	0	0	0
			5213	3321	905	967	20			

- Molecule 2 is a protein called REST corepressor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	0	0
			1076	676	194	203	3			


- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is {N}-[3-(ethoxymethyl)-2-[[4-[[{(3 {R})-pyrrolidin-3-yl]methoxy]phenoxy]met

hyl]phenyl]-4-methyl-thieno[3,2-b]pyrrole-5-carboxamide (three-letter code: 6X5) (formula:  $C_{29}H_{33}N_3O_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			37	29	3	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O		
			6	3	3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	17	Total	O	0	0
			17	17		
6	B	2	Total	O	0	0
			2	2		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: LYSINE-SPECIFIC HISTONE DEMETHYLASE 1



#### • Molecule 2: REST corepressor 1

LYS	E387	GLN	SER	ASP	ALA	PRO	ASP	ALA	ALA	PRO	MET
SER	Q388	ALA	VAL	LEU	ARG	PRO	LEU	ARG	ARG	PRO	VAL
PRO	L389	LYS	MET	PRO	ARG	ARG	ASN	SER	SER	ASN	GLU
ASN	L390	ARG	ASP	PHE	ARG	PHE	GLN	GLN	GLY	ASN	GLY
SER	A391	ALA	HIS	THR	HIS	THR	THR	GLU	GLN	PRO	GLY
ILE	V392	LYS	ALA	PRO	ARG	PHE	PRO	ARG	ASN	GLU	VAL
MET	A394	GLN	LYS	PRO	ASN	ASP	ASP	ASN	SER	SER	VAL
PRO	I395	P311	GLN	ASP	LEU	LEU	LEU	LEU	GLY	GLY	GLY
GLU		K312	LYS	GLU	GLY	GLU	GLY	GLY	ALA	LYS	LYS
GLU	V396	G313	ARG	TRP	MET	TRP	MET	MET	ALA	ARG	ARG
GLU		K314	GLU	THR	LEU	THR	LEU	LEU	ALA	ALA	GLY
ASP	I405	F315	ARG	GLU	ARG	GLU	TRP	TRP	ALA	GLY	GLY
GLU	S406	L316	GLU	GLU	GLU	GLU	VAL	VAL	PRO	PRO	ARG
ALA	D407	S317	SER	ASP	GLU	ASP	SER	SER	ASN	ASN	ASN
PRO	V408	Q318	GLU	VAL	LYS	LYS	PRO	PRO	GLY	GLY	ASN
VAL	I409	E319	GLU	VAL	VAL	VAL	ASN	ASN	ASN	ASN	ALA
LEU		D320	ASP	LEU	GLN	LEU	GLN	GLN	SER	SER	ALA
ASP	V410	V321	GLU	GLU	GLU	GLU	LEU	LEU	SER	SER	ALA
VAL	V411	S325	GLU	ALA	ALA	ALA	ALA	GLU	ASN	ASN	SER
ARG	V415		GLU	PHE	PHE	PHE	PHE	GLU	SER	SER	ALA
TYR	D416	A331	ALA	ALA	ALA	ALA	ALA	ALA	TRP	TRP	ALA
ALA	V417		ASN	SER	SER	SER	SER	LYS	GLU	GLU	ALA
SER	K418	V334	GLY	PHE	PHE	PHE	PHE	LEU	GLU	GLU	ALA
ALA	N419	L335	ASN	HIS	HIS	HIS	HIS	ASP	GLY	GLY	ALA
SER	F420	R336	ASN	GLY	GLY	GLY	GLY	TYR	SER	SER	ALA
	F421	Q337	PRO	LYS	LYS	LYS	LYS	TYR	SER	SER	ALA
	V422		ILE	THR	THR	THR	THR	ILE	GLY	GLY	ALA
	M423	K340	ASP	PHE	PHE	PHE	PHE	ALA	SER	SER	ALA
	Y424		ILE	HIS	HIS	HIS	HIS	ILE	SER	SER	ALA
	R425	S344	GLU	ARG	ARG	ARG	ARG	LYS	SER	SER	ALA
	R426	V345	VAL	ILE	ILE	ILE	ILE	GLU	ASP	ASP	ALA
	D427	K346	ASP	GLN	GLN	GLN	GLN	LYS	GLU	GLU	CYS
	F428	R347	GLN	GLN	GLN	GLN	GLN	LYS	GLU	GLU	ALA
	M429	Q348	ASN	MET	MET	MET	MET	HIS	HIS	HIS	SER
	D430	L349	GLY	LEU	LEU	LEU	LEU	GLY	GLY	GLY	PRO
	D431	Q350	GLU	PRO	PRO	PRO	PRO	TYR	GLY	GLY	ALA
	E432	N351	SER	ASP	ASN	ASN	ASN	GLY	GLY	GLY	ALA
	V433	L352	LYS	LYS	LYS	LYS	LYS	MET	GLY	GLY	THR
	L434	K353	LYS	SER	SER	SER	SER	GLU	MET	MET	ALA
	Q435	Q354	GLU	ILE	ILE	ILE	ILE	GLN	ARG	ARG	ALA
		T355	VAL	ALA	ALA	ALA	ALA	VAL	VAL	VAL	SER
	A439	N356	PRO	LEU	LEU	LEU	LEU	GLY	GLY	GLY	ALA
	E440		THR	VAL	VAL	VAL	VAL	MET	GLN	GLN	ALA
	HIS	L359	GLU	LYS	LYS	LYS	LYS	LEU	TYR	TYR	SER
	GLY		THR	THR	THR	THR	THR	PHE	GLN	GLN	ALA
	LYS	L363	VAL	PHE	PHE	PHE	PHE	TRP	ALA	ALA	SER
	GLU		VAL	TYR	TYR	TYR	TYR	HIS	VAL	VAL	ALA
	THR	I367	PRO	SER	SER	SER	SER	LYS	VAL	VAL	SER
	ASN		GLN	ASN	ASN	ASN	ASN	HIS	VAL	VAL	ALA
	GLY	Y370	VAL	VAL	VAL	VAL	VAL	ASN	PRO	PRO	ALA
	PRO	R371	LYS	LYS	LY						

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.03Å 180.37Å 235.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.04 – 3.40 72.04 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.04-3.40) 99.7 (72.04-3.40)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 3.41Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.195 , 0.225 0.196 , 0.226	Depositor DCC
$R_{free}$ test set	1772 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	97.2	Xtriage
Anisotropy	0.432	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 82.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6410	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6X5, GOL, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/5327	0.55	2/7228 (0.0%)
2	B	0.29	0/1091	0.57	0/1471
All	All	0.30	0/6418	0.55	2/8699 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	731	LEU	CA-CB-CG	5.35	127.61	115.30
1	A	445	LEU	CA-CB-CG	5.09	127.01	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	488	THR	Peptide
1	A	522	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5213	0	5241	215	0
2	B	1076	0	1091	68	0
3	A	53	0	31	6	0
4	A	37	0	0	0	0
5	A	12	0	16	0	0
6	A	17	0	0	0	0
6	B	2	0	0	0	0
All	All	6410	0	6379	263	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 263 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:384:THR:O	2:B:388:GLN:NE2	1.93	0.99
1:A:393:SER:HB3	1:A:549:LEU:HD21	1.58	0.86
1:A:632:GLN:HE21	1:A:636:ALA:HB2	1.42	0.85
1:A:355:LYS:NZ	1:A:356:ILE:O	2.12	0.82
2:B:379:CYS:SG	2:B:416:GLN:NE2	2.53	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/852 (78%)	645 (97%)	19 (3%)	0	100	100
2	B	131/482 (27%)	121 (92%)	10 (8%)	0	100	100
All	All	795/1334 (60%)	766 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	565/699 (81%)	544 (96%)	21 (4%)	34	62
2	B	117/395 (30%)	114 (97%)	3 (3%)	46	72
All	All	682/1094 (62%)	658 (96%)	24 (4%)	36	65

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	571	TYR
1	A	668	ARG
1	A	627	LEU
1	A	710	GLU
1	A	447	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	399	ASN
1	A	527	GLN
1	A	535	ASN
1	A	802	HIS
2	B	416	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	903	-	5,5,5	0.37	0	5,5,5	0.27	0
3	FAD	A	901	-	53,58,58	0.92	2 (3%)	68,89,89	1.29	7 (10%)
5	GOL	A	904	-	5,5,5	0.39	0	5,5,5	0.22	0
4	6X5	A	902	-	38,41,41	2.09	10 (26%)	39,56,56	1.19	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	903	-	-	2/4/4/4	-
3	FAD	A	901	-	-	6/30/50/50	0/6/6/6
5	GOL	A	904	-	-	4/4/4/4	-
4	6X5	A	902	-	-	10/19/29/29	0/5/5/5

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	6X5	C25-N24	5.64	1.64	1.46
4	A	902	6X5	C09-N26	4.48	1.50	1.41
4	A	902	6X5	C27-N26	4.28	1.47	1.35
4	A	902	6X5	C22-C23	4.03	1.59	1.53
4	A	902	6X5	C11-C10	3.36	1.57	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	FAD	N3A-C2A-N1A	-4.65	121.41	128.68
4	A	902	6X5	C36-S37-C31	4.29	96.23	91.55
4	A	902	6X5	C22-C21-C25	3.63	106.60	102.14
3	A	901	FAD	C4-N3-C2	-2.74	120.58	125.64
3	A	901	FAD	O4-C4-C4X	-2.54	119.86	126.60

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	FAD	C5B-O5B-PA-O1A
4	A	902	6X5	O19-C20-C21-C22
4	A	902	6X5	O19-C20-C21-C25
4	A	902	6X5	O28-C27-C29-C30
5	A	904	GOL	O1-C1-C2-C3


There are no ring outliers.


1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	FAD	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	666/852 (78%)	0.36	7 (1%) 80 79	44, 91, 142, 181	0
2	B	133/482 (27%)	0.41	2 (1%) 73 72	80, 130, 177, 223	0
All	All	799/1334 (59%)	0.36	9 (1%) 80 79	44, 98, 155, 223	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	311	PRO	3.2
2	B	316	LEU	3.1
1	A	437	THR	2.6
1	A	667	ASP	2.6
1	A	174	VAL	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.







Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	A	903	6/6	0.82	0.47	91,95,98,102	0
5	GOL	A	904	6/6	0.91	0.50	120,122,123,127	0
4	6X5	A	902	37/37	0.93	0.37	55,89,132,136	0
3	FAD	A	901	53/53	0.98	0.27	34,65,76,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 6X5 A 902:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
 and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.