



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 16, 2025 – 08:46 PM EDT

PDB ID : 3LSQ / pdb_00003lsq
Title : Trypanosoma brucei seryl-tRNA synthetase
Authors : Larson, E.T.; Merritt, E.A.; Medical Structural Genomics of Pathogenic Protozoa (MSGPP)
Deposited on : 2010-02-12
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	
1	B	484	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MLI	A	602	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Seryl-tRNA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	431	Total	C	N	O	S	0	5	0
			3447	2170	600	648	29			
1	B	458	Total	C	N	O	S	0	6	0
			3649	2293	629	698	29			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q384V4
A	-3	PRO	-	expression tag	UNP Q384V4
A	-2	GLY	-	expression tag	UNP Q384V4
A	-1	SER	-	expression tag	UNP Q384V4
A	0	MET	-	expression tag	UNP Q384V4
B	-4	GLY	-	expression tag	UNP Q384V4
B	-3	PRO	-	expression tag	UNP Q384V4
B	-2	GLY	-	expression tag	UNP Q384V4
B	-1	SER	-	expression tag	UNP Q384V4
B	0	MET	-	expression tag	UNP Q384V4

- Molecule 2 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	142	Total	O	0	0
			142	142		
4	B	141	Total	O	0	0
			141	141		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Seryl-tRNA synthetase



• Molecule 1: Seryl-tRNA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	176.02Å 176.02Å 248.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.45 – 2.10 37.45 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.45-2.10) 99.5 (37.45-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.98 (at 2.10Å)	Xtriage
Refinement program	REFMAC refmac_5.5.0106	Depositor
R, R_{free}	0.171 , 0.203 0.179 , 0.208	Depositor DCC
R_{free} test set	4328 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.286	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7387	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/3521	0.71	2/4737 (0.0%)
1	B	0.68	1/3729 (0.0%)	0.70	2/5024 (0.0%)
All	All	0.69	1/7250 (0.0%)	0.71	4/9761 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	443	GLU	CB-CG	5.04	1.61	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	230	PHE	CB-CA-C	-6.68	97.04	110.40
1	A	230	PHE	CB-CA-C	-6.17	98.06	110.40
1	B	190	MET	CG-SD-CE	5.45	108.92	100.20
1	A	295	ARG	NE-CZ-NH2	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3447	0	3454	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3649	0	3638	26	0
2	A	7	0	2	2	0
3	A	1	0	0	0	0
4	A	142	0	0	1	0
4	B	141	0	0	0	0
All	All	7387	0	7094	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:ILE:HG22	1:B:34:ILE:CD1	1.97	0.95
1:B:30:ILE:HG22	1:B:34:ILE:HD11	1.50	0.91
1:B:214:LEU:HD13	1:B:290[B]:MET:SD	2.14	0.87
1:B:134:LEU:HD11	1:B:242:LEU:HD12	1.60	0.82
1:B:33:ALA:HB1	1:B:132:LEU:HD23	1.62	0.81
1:B:30:ILE:O	1:B:34:ILE:HD12	1.83	0.79
1:B:290[B]:MET:SD	1:B:317:ILE:CD1	2.74	0.76
1:B:30:ILE:CG2	1:B:34:ILE:HD11	2.14	0.75
1:A:214:LEU:HD13	1:A:290[B]:MET:SD	2.30	0.72
1:B:290[B]:MET:SD	1:B:317:ILE:HD12	2.39	0.61
1:A:127[A]:GLU:HA	1:A:127[A]:GLU:OE1	2.01	0.60
1:A:276:ARG:HD3	2:A:602:MLI:H11	1.85	0.58
1:B:290[B]:MET:SD	1:B:317:ILE:HD13	2.44	0.58
1:B:33:ALA:HB1	1:B:132:LEU:CD2	2.34	0.57
1:A:317:ILE:HD12	1:A:317:ILE:N	2.20	0.56
1:B:128:GLU:O	1:B:132:LEU:HD13	2.04	0.56
1:A:231:LEU:HD22	1:B:190:MET:HG2	1.89	0.53
1:B:241:GLN:HE22	1:B:364:ASN:ND2	2.07	0.52
1:A:306:LEU:HD23	4:A:942:HOH:O	2.07	0.52
1:B:372:LEU:HB3	1:B:386:VAL:HB	1.93	0.51
1:B:81:GLN:OE1	1:B:95:GLN:NE2	2.42	0.51
1:B:30:ILE:CG2	1:B:34:ILE:CD1	2.78	0.51
1:B:33:ALA:CB	1:B:132:LEU:HD23	2.35	0.51
1:A:372:LEU:HB3	1:A:386:VAL:HB	1.94	0.50
1:A:187:VAL:HA	1:A:190:MET:HE2	1.95	0.48
1:A:371:ASP:OD1	1:A:387:SER:HB3	2.14	0.48
1:A:157:VAL:HG12	1:A:158:ARG:HG3	1.96	0.47
1:B:295:ARG:NH1	1:B:297:GLU:OE2	2.45	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:LEU:HD23	1:A:17:ILE:CG1	2.47	0.45
1:A:150:GLU:HG3	1:A:361:ALA:CB	2.48	0.44
1:A:231:LEU:CD2	1:B:190:MET:HG2	2.48	0.43
1:B:294:PHE:HA	1:B:312:HIS:O	2.18	0.43
1:B:316:LYS:HD2	1:B:425:THR:HG23	2.00	0.43
1:B:30:ILE:HG22	1:B:34:ILE:HD12	1.91	0.43
1:A:187:VAL:HA	1:A:190:MET:CE	2.49	0.43
1:A:294:PHE:HA	1:A:312:HIS:O	2.19	0.42
1:B:134:LEU:CD1	1:B:242:LEU:HD12	2.41	0.42
1:A:187:VAL:O	1:A:190:MET:HB2	2.20	0.41
1:A:276:ARG:HH11	2:A:602:MLI:H11	1.85	0.41
1:B:453:ARG:HB2	1:B:454:PRO:HD3	2.02	0.41
1:A:453:ARG:N	1:A:454:PRO:CD	2.84	0.40
1:B:279:THR:HA	1:B:404:GLY:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/484 (89%)	425 (99%)	5 (1%)	0	100	100
1	B	460/484 (95%)	451 (98%)	8 (2%)	1 (0%)	44	45
All	All	890/968 (92%)	876 (98%)	13 (2%)	1 (0%)	48	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	412	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/412 (92%)	370 (98%)	7 (2%)	52	59
1	B	400/412 (97%)	395 (99%)	5 (1%)	65	72
All	All	777/824 (94%)	765 (98%)	12 (2%)	62	67

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	ILE
1	A	108	SER
1	A	127[A]	GLU
1	A	127[B]	GLU
1	A	131	LYS
1	A	264	THR
1	A	306	LEU
1	B	12	THR
1	B	69	LYS
1	B	241	GLN
1	B	264	THR
1	B	274	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	241	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLI	A	602	-	6,6,6	1.89	2 (33%)	7,7,7	1.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	A	602	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	602	MLI	C1-C2	2.84	1.55	1.51
2	A	602	MLI	C1-C3	2.52	1.55	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	602	MLI	C2-C1-C3-O8
2	A	602	MLI	C2-C1-C3-O9

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	602	MLI	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/484 (89%)	0.23	53 (12%) 9 10	16, 33, 93, 129	5 (1%)
1	B	458/484 (94%)	0.06	41 (8%) 17 18	16, 35, 83, 111	6 (1%)
All	All	889/968 (91%)	0.14	94 (10%) 13 13	16, 35, 86, 129	11 (1%)

All (94) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	132	LEU	9.6
1	A	71	ALA	7.3
1	A	410	THR	7.2
1	B	131	LYS	6.5
1	B	2	VAL	5.4
1	A	255	ASP	5.4
1	A	0	MET	5.4
1	B	134	LEU	5.3
1	A	298	ALA	5.1
1	A	411	ALA	5.1
1	A	412	ALA	5.1
1	A	57	ILE	5.0
1	A	407	LEU	4.9
1	B	1	MET	4.6
1	B	242	LEU	4.6
1	A	409	GLY	4.5
1	A	102	LEU	4.3
1	B	3	LEU	4.2
1	B	306	LEU	4.1
1	B	255	ASP	4.1
1	A	242	LEU	4.1
1	B	30	ILE	4.0
1	A	101	VAL	3.9
1	A	408	ARG	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	4	ASP	3.9
1	B	133[A]	MET	3.8
1	A	306	LEU	3.8
1	B	412	ALA	3.7
1	A	246	ASP	3.4
1	A	68	ALA	3.4
1	A	120	LYS	3.3
1	B	241	GLN	3.3
1	B	297	GLU	3.3
1	A	54	LEU	3.3
1	A	241	GLN	3.2
1	A	254	GLY	3.2
1	A	466	ASN	3.2
1	A	69	LYS	3.1
1	A	125	LEU	3.1
1	A	234	ASP	3.1
1	B	414	ASN	3.0
1	A	405	PRO	2.9
1	A	46	PHE	2.9
1	A	103	GLN	2.9
1	A	67	LYS	2.9
1	A	413	GLN	2.9
1	A	53	LYS	2.8
1	B	29	ASP	2.8
1	B	151	GLU	2.8
1	B	415	VAL	2.8
1	A	383[A]	ARG	2.7
1	B	246	ASP	2.7
1	A	406	ASN	2.7
1	B	136	VAL	2.6
1	B	411	ALA	2.6
1	A	297	GLU	2.6
1	A	66	LYS	2.6
1	B	135	ASN	2.6
1	B	6	GLN	2.6
1	B	257	ASP	2.5
1	A	47	LEU	2.5
1	A	245	PHE	2.5
1	A	59	SER	2.5
1	A	256	GLY	2.5
1	A	1	MET	2.4
1	B	298	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	245	PHE	2.4
1	B	123	GLN	2.3
1	B	130	ASP	2.3
1	A	127[A]	GLU	2.3
1	A	104	LEU	2.3
1	A	151	GLU	2.2
1	B	125	LEU	2.2
1	B	410	THR	2.2
1	A	61	ALA	2.2
1	B	243	SER	2.2
1	A	4	ASP	2.2
1	B	254	GLY	2.2
1	A	248	GLU	2.2
1	A	64	ALA	2.2
1	B	258	LYS	2.2
1	B	413	GLN	2.2
1	B	126	GLU	2.2
1	A	105	LYS	2.2
1	B	244	GLN	2.2
1	B	129	ARG	2.1
1	B	7	LEU	2.1
1	A	247	GLU	2.1
1	A	123	GLN	2.1
1	B	94	GLU	2.1
1	A	414	ASN	2.0
1	A	415	VAL	2.0
1	B	34	ILE	2.0
1	A	58	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MLI	A	602	7/7	0.82	0.13	40,51,62,63	0
3	NA	A	600	1/1	0.99	0.11	25,25,25,25	1

6.5 Other polymers [i](#)

There are no such residues in this entry.