



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 28, 2024 – 03:07 am GMT

PDB ID : 5MTZ
Title : Crystal structure of a long form RNase Z from yeast
Authors : Li de la Sierra-Gallay, I.; Miao, M.; van Tilbeurgh, H.
Deposited on : 2017-01-11
Resolution : 2.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	874	 73% 13% • 12%
1	B	874	 73% 12% • 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease Z.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	768	Total	C	N	O	S	Se	0	0	0
			6284	4020	1046	1192	8	18			
1	B	755	Total	C	N	O	S	Se	0	0	0
			6174	3954	1031	1163	8	18			

There are 74 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-35	MSE	-	initiating methionine	UNP P36159
A	-34	ALA	-	expression tag	UNP P36159
A	-33	HIS	-	expression tag	UNP P36159
A	-32	HIS	-	expression tag	UNP P36159
A	-31	HIS	-	expression tag	UNP P36159
A	-30	HIS	-	expression tag	UNP P36159
A	-29	HIS	-	expression tag	UNP P36159
A	-28	HIS	-	expression tag	UNP P36159
A	-27	VAL	-	expression tag	UNP P36159
A	-26	GLY	-	expression tag	UNP P36159
A	-25	THR	-	expression tag	UNP P36159
A	-24	GLY	-	expression tag	UNP P36159
A	-23	SER	-	expression tag	UNP P36159
A	-22	ASN	-	expression tag	UNP P36159
A	-21	ASP	-	expression tag	UNP P36159
A	-20	ASP	-	expression tag	UNP P36159
A	-19	ASP	-	expression tag	UNP P36159
A	-18	ASP	-	expression tag	UNP P36159
A	-17	LYS	-	expression tag	UNP P36159
A	-16	SER	-	expression tag	UNP P36159
A	-15	PRO	-	expression tag	UNP P36159
A	-14	ASP	-	expression tag	UNP P36159
A	-13	PRO	-	expression tag	UNP P36159
A	-12	ASN	-	expression tag	UNP P36159
A	-11	TRP	-	expression tag	UNP P36159

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	GLU	-	expression tag	UNP P36159
A	-9	LEU	-	expression tag	UNP P36159
A	-8	VAL	-	expression tag	UNP P36159
A	-7	TYR	-	expression tag	UNP P36159
A	-6	THR	-	expression tag	UNP P36159
A	-5	ALA	-	expression tag	UNP P36159
A	-4	ARG	-	expression tag	UNP P36159
A	-3	LEU	-	expression tag	UNP P36159
A	-2	GLN	-	expression tag	UNP P36159
A	-1	GLU	-	expression tag	UNP P36159
A	0	PHE	-	expression tag	UNP P36159
A	1	MSE	-	expression tag	UNP P36159
B	-35	MSE	-	initiating methionine	UNP P36159
B	-34	ALA	-	expression tag	UNP P36159
B	-33	HIS	-	expression tag	UNP P36159
B	-32	HIS	-	expression tag	UNP P36159
B	-31	HIS	-	expression tag	UNP P36159
B	-30	HIS	-	expression tag	UNP P36159
B	-29	HIS	-	expression tag	UNP P36159
B	-28	HIS	-	expression tag	UNP P36159
B	-27	VAL	-	expression tag	UNP P36159
B	-26	GLY	-	expression tag	UNP P36159
B	-25	THR	-	expression tag	UNP P36159
B	-24	GLY	-	expression tag	UNP P36159
B	-23	SER	-	expression tag	UNP P36159
B	-22	ASN	-	expression tag	UNP P36159
B	-21	ASP	-	expression tag	UNP P36159
B	-20	ASP	-	expression tag	UNP P36159
B	-19	ASP	-	expression tag	UNP P36159
B	-18	ASP	-	expression tag	UNP P36159
B	-17	LYS	-	expression tag	UNP P36159
B	-16	SER	-	expression tag	UNP P36159
B	-15	PRO	-	expression tag	UNP P36159
B	-14	ASP	-	expression tag	UNP P36159
B	-13	PRO	-	expression tag	UNP P36159
B	-12	ASN	-	expression tag	UNP P36159
B	-11	TRP	-	expression tag	UNP P36159
B	-10	GLU	-	expression tag	UNP P36159
B	-9	LEU	-	expression tag	UNP P36159
B	-8	VAL	-	expression tag	UNP P36159
B	-7	TYR	-	expression tag	UNP P36159
B	-6	THR	-	expression tag	UNP P36159

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	ALA	-	expression tag	UNP P36159
B	-4	ARG	-	expression tag	UNP P36159
B	-3	LEU	-	expression tag	UNP P36159
B	-2	GLN	-	expression tag	UNP P36159
B	-1	GLU	-	expression tag	UNP P36159
B	0	PHE	-	expression tag	UNP P36159
B	1	MSE	-	expression tag	UNP P36159

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ribonuclease Z



• Molecule 1: Ribonuclease Z



F394	S395	Q396	V400	T401	F402	E410	P411	M412	N415	I416	N417	D422	Q426	V446	I447	N448	N449	Q450	V453	D454	V466	LYS	R482	L488	V491	R503	T516	Q523	V526	M535	I536	Y537	H546	I549	D562	T565	V571	W574	
E582	W583	L584	I591	Y597	S607	F608	V609	R610	M611	F621	N622	E623	I637	L637	R641	Y662	F681	R682	M683	F692	R701	L709	S715	L718	L729	H737	N741	M750	N751	A752									
L770	D771	A778	R800	I801	L804	F809	VAL	GLU	GLU	LYS	GLU	GLU	GLU	GLU	ASP	VAL	ASP	ASP	VAL	GLU	SER	VAL	GLN	ASP	LEU	GLU	VAL	LYS	LYS	HIS	LYS	LYS	ASN						

4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	136.00Å 136.00Å 115.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.61 – 2.99 49.61 – 2.99	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.61-2.99) 99.8 (49.61-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.181 , 0.262 0.191 , 0.269	Depositor DCC
R_{free} test set	2135 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	85.6	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.038 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12472	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/6410	0.74	1/8638 (0.0%)
1	B	0.53	1/6300 (0.0%)	0.73	0/8491
All	All	0.53	1/12710 (0.0%)	0.73	1/17129 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	247	GLU	CD-OE1	5.59	1.31	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	LEU	CA-CB-CG	5.02	126.84	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6284	0	6147	49	0
1	B	6174	0	6048	42	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	0	0
3	B	5	0	0	0	0
All	All	12472	0	12195	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:365:ASP:O	1:A:366:THR:HG22	1.78	0.83
1:A:157:ILE:HG22	1:A:161:MSE:HE2	1.65	0.79
1:B:681:PHE:HB2	1:B:683:MSE:CE	2.20	0.72
1:B:681:PHE:HB2	1:B:683:MSE:HE1	1.73	0.71
1:A:343:GLN:HE22	1:A:798:GLN:HE21	1.38	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	762/874 (87%)	686 (90%)	66 (9%)	10 (1%)	10	39
1	B	749/874 (86%)	669 (89%)	69 (9%)	11 (2%)	8	36
All	All	1511/1748 (86%)	1355 (90%)	135 (9%)	21 (1%)	9	37

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	141	ASP
1	B	370	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	145	VAL
1	A	167	PRO
1	A	407	MSE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	707/783 (90%)	655 (93%)	52 (7%)	11	38
1	B	694/783 (89%)	646 (93%)	48 (7%)	13	42
All	All	1401/1566 (90%)	1301 (93%)	100 (7%)	12	40

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	175	SER
1	B	385	ILE
1	B	809	PHE
1	B	199	SER
1	B	353	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	653	GLN
1	B	664	GLN
1	B	762	GLN
1	B	751	ASN
1	B	415	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	903	2	4,4,4	0.97	0	6,6,6	0.56	0
3	PO4	B	903	2	4,4,4	0.78	0	6,6,6	0.97	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	750/874 (85%)	-0.53	3 (0%) 89 77	50, 80, 121, 166	0
1	B	737/874 (84%)	-0.49	1 (0%) 92 88	51, 80, 134, 171	0
All	All	1487/1748 (85%)	-0.51	4 (0%) 90 81	50, 80, 128, 171	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	PHE	4.0
1	A	246	LEU	3.0
1	B	0	PHE	2.5
1	A	143	LEU	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	903	5/5	0.95	0.06	77,89,95,102	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PO4	B	903	5/5	0.96	0.05	75,78,85,86	0
2	ZN	B	901	1/1	0.99	0.04	73,73,73,73	0
2	ZN	B	902	1/1	1.00	0.02	68,68,68,68	0
2	ZN	A	902	1/1	1.00	0.02	73,73,73,73	0
2	ZN	A	901	1/1	1.00	0.03	83,83,83,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.