



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 9, 2024 – 01:39 PM EST

PDB ID : 6N1F
Title : Crystal structure of Oxidoreductase, 2OG-Fe(II) oxygenase family, from Burkholderia pseudomallei
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2018-11-08
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>8%</div> <div>13%</div> </div> </div>
1	B	237	<div> <div>5%</div> <div> <div></div> <div>79%</div> <div>8%</div> <div>12%</div> </div> </div>
1	C	237	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>8%</div> <div>14%</div> </div> </div>
1	D	237	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>5%</div> <div>10%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxidoreductase, 2OG-Fe(II) oxygenase family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	206	Total	C	N	O	S	Se	0	1	0
			1616	1017	299	291	3	6			
1	B	208	Total	C	N	O	S	Se	0	1	0
			1623	1023	301	290	3	6			
1	C	205	Total	C	N	O	S	Se	0	0	0
			1599	1008	296	286	3	6			
1	D	213	Total	C	N	O	S	Se	0	0	0
			1637	1029	299	300	3	6			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MSE	-	expression tag	UNP Q3JFV4
A	-19	ALA	-	expression tag	UNP Q3JFV4
A	-18	HIS	-	expression tag	UNP Q3JFV4
A	-17	HIS	-	expression tag	UNP Q3JFV4
A	-16	HIS	-	expression tag	UNP Q3JFV4
A	-15	HIS	-	expression tag	UNP Q3JFV4
A	-14	HIS	-	expression tag	UNP Q3JFV4
A	-13	HIS	-	expression tag	UNP Q3JFV4
A	-12	MSE	-	expression tag	UNP Q3JFV4
A	-11	GLY	-	expression tag	UNP Q3JFV4
A	-10	THR	-	expression tag	UNP Q3JFV4
A	-9	LEU	-	expression tag	UNP Q3JFV4
A	-8	GLU	-	expression tag	UNP Q3JFV4
A	-7	ALA	-	expression tag	UNP Q3JFV4
A	-6	GLN	-	expression tag	UNP Q3JFV4
A	-5	THR	-	expression tag	UNP Q3JFV4
A	-4	GLN	-	expression tag	UNP Q3JFV4
A	-3	GLY	-	expression tag	UNP Q3JFV4
A	-2	PRO	-	expression tag	UNP Q3JFV4
A	-1	GLY	-	expression tag	UNP Q3JFV4
A	0	SER	-	expression tag	UNP Q3JFV4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MSE	-	expression tag	UNP Q3JFV4
B	-19	ALA	-	expression tag	UNP Q3JFV4
B	-18	HIS	-	expression tag	UNP Q3JFV4
B	-17	HIS	-	expression tag	UNP Q3JFV4
B	-16	HIS	-	expression tag	UNP Q3JFV4
B	-15	HIS	-	expression tag	UNP Q3JFV4
B	-14	HIS	-	expression tag	UNP Q3JFV4
B	-13	HIS	-	expression tag	UNP Q3JFV4
B	-12	MSE	-	expression tag	UNP Q3JFV4
B	-11	GLY	-	expression tag	UNP Q3JFV4
B	-10	THR	-	expression tag	UNP Q3JFV4
B	-9	LEU	-	expression tag	UNP Q3JFV4
B	-8	GLU	-	expression tag	UNP Q3JFV4
B	-7	ALA	-	expression tag	UNP Q3JFV4
B	-6	GLN	-	expression tag	UNP Q3JFV4
B	-5	THR	-	expression tag	UNP Q3JFV4
B	-4	GLN	-	expression tag	UNP Q3JFV4
B	-3	GLY	-	expression tag	UNP Q3JFV4
B	-2	PRO	-	expression tag	UNP Q3JFV4
B	-1	GLY	-	expression tag	UNP Q3JFV4
B	0	SER	-	expression tag	UNP Q3JFV4
C	-20	MSE	-	expression tag	UNP Q3JFV4
C	-19	ALA	-	expression tag	UNP Q3JFV4
C	-18	HIS	-	expression tag	UNP Q3JFV4
C	-17	HIS	-	expression tag	UNP Q3JFV4
C	-16	HIS	-	expression tag	UNP Q3JFV4
C	-15	HIS	-	expression tag	UNP Q3JFV4
C	-14	HIS	-	expression tag	UNP Q3JFV4
C	-13	HIS	-	expression tag	UNP Q3JFV4
C	-12	MSE	-	expression tag	UNP Q3JFV4
C	-11	GLY	-	expression tag	UNP Q3JFV4
C	-10	THR	-	expression tag	UNP Q3JFV4
C	-9	LEU	-	expression tag	UNP Q3JFV4
C	-8	GLU	-	expression tag	UNP Q3JFV4
C	-7	ALA	-	expression tag	UNP Q3JFV4
C	-6	GLN	-	expression tag	UNP Q3JFV4
C	-5	THR	-	expression tag	UNP Q3JFV4
C	-4	GLN	-	expression tag	UNP Q3JFV4
C	-3	GLY	-	expression tag	UNP Q3JFV4
C	-2	PRO	-	expression tag	UNP Q3JFV4
C	-1	GLY	-	expression tag	UNP Q3JFV4
C	0	SER	-	expression tag	UNP Q3JFV4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MSE	-	expression tag	UNP Q3JFV4
D	-19	ALA	-	expression tag	UNP Q3JFV4
D	-18	HIS	-	expression tag	UNP Q3JFV4
D	-17	HIS	-	expression tag	UNP Q3JFV4
D	-16	HIS	-	expression tag	UNP Q3JFV4
D	-15	HIS	-	expression tag	UNP Q3JFV4
D	-14	HIS	-	expression tag	UNP Q3JFV4
D	-13	HIS	-	expression tag	UNP Q3JFV4
D	-12	MSE	-	expression tag	UNP Q3JFV4
D	-11	GLY	-	expression tag	UNP Q3JFV4
D	-10	THR	-	expression tag	UNP Q3JFV4
D	-9	LEU	-	expression tag	UNP Q3JFV4
D	-8	GLU	-	expression tag	UNP Q3JFV4
D	-7	ALA	-	expression tag	UNP Q3JFV4
D	-6	GLN	-	expression tag	UNP Q3JFV4
D	-5	THR	-	expression tag	UNP Q3JFV4
D	-4	GLN	-	expression tag	UNP Q3JFV4
D	-3	GLY	-	expression tag	UNP Q3JFV4
D	-2	PRO	-	expression tag	UNP Q3JFV4
D	-1	GLY	-	expression tag	UNP Q3JFV4
D	0	SER	-	expression tag	UNP Q3JFV4

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	C	1	Total Cl 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	118	Total 119	O 119	0	1
4	B	112	Total 112	O 112	0	0
4	C	109	Total 109	O 109	0	0
4	D	103	Total 103	O 103	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Oxidoreductase, 2OG-Fe(II) oxygenase family




- Molecule 1: Oxidoreductase, 2OG-Fe(II) oxygenase family




- Molecule 1: Oxidoreductase, 2OG-Fe(II) oxygenase family



- Molecule 1: Oxidoreductase, 2OG-Fe(II) oxygenase family





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.03Å 54.99Å 112.10Å 77.66° 87.03° 64.73°	Depositor
Resolution (Å)	27.35 – 2.05 27.35 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.3 (27.35-2.05) 96.3 (27.35-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.26 (at 2.05Å)	Xtriage
Refinement program	PHENIX (dev_3304)	Depositor
R, R_{free}	0.208 , 0.242 0.210 , 0.241	Depositor DCC
R_{free} test set	4294 reflections (3.38%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.477	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6924	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1650	0.54	0/2227
1	B	0.38	0/1657	0.56	0/2237
1	C	0.35	0/1630	0.54	0/2201
1	D	0.36	0/1669	0.55	0/2258
All	All	0.36	0/6606	0.55	0/8923

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	111	THR	Peptide
1	D	111	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1616	0	1552	13	0
1	B	1623	0	1556	13	0
1	C	1599	0	1531	12	0
1	D	1637	0	1551	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	119	0	0	3	0
4	B	112	0	0	5	0
4	C	109	0	0	3	0
4	D	103	0	0	3	0
All	All	6924	0	6190	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:10:CYS:SG	4:D:499:HOH:O	2.38	0.80
1:C:124:ARG:NH2	1:C:197:GLU:OE1	2.21	0.74
1:B:136:ASP:OD2	4:B:401:HOH:O	2.11	0.68
1:B:111:THR:HG21	1:B:115:TYR:HE2	1.61	0.64
1:B:23:ARG:NH1	4:B:405:HOH:O	2.31	0.63

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/237 (86%)	198 (98%)	5 (2%)	0	100	100
1	B	205/237 (86%)	200 (98%)	5 (2%)	0	100	100
1	C	201/237 (85%)	195 (97%)	6 (3%)	0	100	100
1	D	211/237 (89%)	204 (97%)	7 (3%)	0	100	100
All	All	820/948 (86%)	797 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/182 (90%)	164 (100%)	0	100	100
1	B	163/182 (90%)	160 (98%)	3 (2%)	54	52
1	C	161/182 (88%)	161 (100%)	0	100	100
1	D	163/182 (90%)	160 (98%)	3 (2%)	54	52
All	All	651/728 (89%)	645 (99%)	6 (1%)	75	77

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	111	THR
1	D	183	SER
1	D	196	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	111	THR
1	B	108	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	196	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/237 (84%)	0.41	13 (6%) 26 28	18, 36, 63, 86	1 (0%)
1	B	202/237 (85%)	0.24	11 (5%) 32 34	18, 32, 59, 85	1 (0%)
1	C	199/237 (83%)	0.41	13 (6%) 26 28	19, 35, 60, 89	0
1	D	207/237 (87%)	0.24	12 (5%) 30 32	17, 32, 61, 85	0
All	All	808/948 (85%)	0.32	49 (6%) 28 29	17, 34, 61, 89	2 (0%)

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	153	ARG	5.3
1	B	112	ARG	4.8
1	C	153	ARG	4.8
1	D	137	ALA	4.5
1	D	112	ARG	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FE	C	300	1/1	0.97	0.06	52,52,52,52	1
2	FE	A	300	1/1	0.98	0.05	52,52,52,52	0
2	FE	D	300	1/1	0.98	0.07	50,50,50,50	0
2	FE	B	300	1/1	0.99	0.06	44,44,44,44	1
3	CL	A	301	1/1	0.99	0.02	21,21,21,21	0
3	CL	C	301	1/1	0.99	0.10	5,5,5,5	1

6.5 Other polymers [i](#)

There are no such residues in this entry.