



wwPDB EM Validation Summary Report ⓘ

Jul 27, 2026 – 10:16 AM EDT

PDB ID : 9N3Y / pdb_00009n3y
EMDB ID : EMD-48860
Title : Type IV-A1 CRISPR effector complex bound to dsDNA and CasDinG - loaded state
Authors : Kiernan, K.A.; Taylor, D.W.
Deposited on : 2025-02-01
Resolution : 3.00 Å(reported)
Based on initial model : 7XG4

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	220	
2	H	232	
3	I	61	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	J	54	
5	K	54	
6	A	241	
7	C	359	
7	D	359	
7	E	359	
7	F	359	
7	G	359	
8	L	748	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 25206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCRISPR type AFERR-associated protein Csf3 (Cas5).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	219	Total	C	N	O	S	0	0
			1710	1080	316	306	8		

- Molecule 2 is a protein called CRISPR type AFERR-associated protein Csf5 (Cas6).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	226	Total	C	N	O	S	0	0
			1737	1112	309	309	7		

- Molecule 3 is a RNA chain called crRNA (61-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
3	I	61	Total	C	N	O	P	0	0
			1314	586	247	421	60		

- Molecule 4 is a DNA chain called DNA, non-target strand (48-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
4	J	48	Total	C	N	O	P	0	0
			996	470	196	282	48		

- Molecule 5 is a DNA chain called DNA, target strand (48-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	K	48	Total	C	N	O	P	0	0
			972	464	166	294	48		

- Molecule 6 is a protein called CRISPR type AFERR-associated protein Csf1 (Cas8).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	A	241	Total	C	N	O	S	0	0
			1889	1196	334	347	12		

- Molecule 7 is a protein called CRISPR type AFERR-associated protein Csf2 (Cas7).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	C	329	Total	C	N	O	S	0	0
			2482	1567	436	467	12		
7	D	331	Total	C	N	O	S	0	0
			2499	1577	440	470	12		
7	E	327	Total	C	N	O	S	0	0
			2465	1555	434	464	12		
7	F	326	Total	C	N	O	S	0	0
			2460	1552	433	463	12		
7	G	276	Total	C	N	O	S	0	0
			2090	1324	364	391	11		

There are 55 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-10	MET	-	initiating methionine	UNP A0AAB0B1V3
C	-9	TRP	-	expression tag	UNP A0AAB0B1V3
C	-8	SER	-	expression tag	UNP A0AAB0B1V3
C	-7	HIS	-	expression tag	UNP A0AAB0B1V3
C	-6	PRO	-	expression tag	UNP A0AAB0B1V3
C	-5	GLN	-	expression tag	UNP A0AAB0B1V3
C	-4	PHE	-	expression tag	UNP A0AAB0B1V3
C	-3	GLU	-	expression tag	UNP A0AAB0B1V3
C	-2	LYS	-	expression tag	UNP A0AAB0B1V3
C	-1	GLY	-	expression tag	UNP A0AAB0B1V3
C	0	ALA	-	expression tag	UNP A0AAB0B1V3
D	-10	MET	-	initiating methionine	UNP A0AAB0B1V3
D	-9	TRP	-	expression tag	UNP A0AAB0B1V3
D	-8	SER	-	expression tag	UNP A0AAB0B1V3
D	-7	HIS	-	expression tag	UNP A0AAB0B1V3
D	-6	PRO	-	expression tag	UNP A0AAB0B1V3
D	-5	GLN	-	expression tag	UNP A0AAB0B1V3
D	-4	PHE	-	expression tag	UNP A0AAB0B1V3
D	-3	GLU	-	expression tag	UNP A0AAB0B1V3
D	-2	LYS	-	expression tag	UNP A0AAB0B1V3
D	-1	GLY	-	expression tag	UNP A0AAB0B1V3
D	0	ALA	-	expression tag	UNP A0AAB0B1V3
E	-10	MET	-	initiating methionine	UNP A0AAB0B1V3
E	-9	TRP	-	expression tag	UNP A0AAB0B1V3
E	-8	SER	-	expression tag	UNP A0AAB0B1V3
E	-7	HIS	-	expression tag	UNP A0AAB0B1V3
E	-6	PRO	-	expression tag	UNP A0AAB0B1V3
E	-5	GLN	-	expression tag	UNP A0AAB0B1V3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-4	PHE	-	expression tag	UNP A0AAB0B1V3
E	-3	GLU	-	expression tag	UNP A0AAB0B1V3
E	-2	LYS	-	expression tag	UNP A0AAB0B1V3
E	-1	GLY	-	expression tag	UNP A0AAB0B1V3
E	0	ALA	-	expression tag	UNP A0AAB0B1V3
F	-10	MET	-	initiating methionine	UNP A0AAB0B1V3
F	-9	TRP	-	expression tag	UNP A0AAB0B1V3
F	-8	SER	-	expression tag	UNP A0AAB0B1V3
F	-7	HIS	-	expression tag	UNP A0AAB0B1V3
F	-6	PRO	-	expression tag	UNP A0AAB0B1V3
F	-5	GLN	-	expression tag	UNP A0AAB0B1V3
F	-4	PHE	-	expression tag	UNP A0AAB0B1V3
F	-3	GLU	-	expression tag	UNP A0AAB0B1V3
F	-2	LYS	-	expression tag	UNP A0AAB0B1V3
F	-1	GLY	-	expression tag	UNP A0AAB0B1V3
F	0	ALA	-	expression tag	UNP A0AAB0B1V3
G	-10	MET	-	initiating methionine	UNP A0AAB0B1V3
G	-9	TRP	-	expression tag	UNP A0AAB0B1V3
G	-8	SER	-	expression tag	UNP A0AAB0B1V3
G	-7	HIS	-	expression tag	UNP A0AAB0B1V3
G	-6	PRO	-	expression tag	UNP A0AAB0B1V3
G	-5	GLN	-	expression tag	UNP A0AAB0B1V3
G	-4	PHE	-	expression tag	UNP A0AAB0B1V3
G	-3	GLU	-	expression tag	UNP A0AAB0B1V3
G	-2	LYS	-	expression tag	UNP A0AAB0B1V3
G	-1	GLY	-	expression tag	UNP A0AAB0B1V3
G	0	ALA	-	expression tag	UNP A0AAB0B1V3

- Molecule 8 is a protein called CRISPR-associated DEAD/DEAH-box helicase CasDinG.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	L	613	Total	C	N	O	S	0	0
			4591	2897	826	850	18		


- Molecule 9 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
9	A	1	Total	Zn	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCRISPR type AFERR-associated protein Csf3 (Cas5)




- Molecule 2: CRISPR type AFERR-associated protein Csf5 (Cas6)




- Molecule 3: crRNA (61-MER)




- Molecule 4: DNA, non-target strand (48-MER)




- Molecule 5: DNA, target strand (48-MER)

Chain K: 




- Molecule 6: CRISPR type AFERR-associated protein Csf1 (Cas8)

Chain A: 




- Molecule 7: CRISPR type AFERR-associated protein Csf2 (Cas7)

Chain C: 




- Molecule 7: CRISPR type AFERR-associated protein Csf2 (Cas7)

Chain D: 




- Molecule 7: CRISPR type AFERR-associated protein Csf2 (Cas7)

Chain E: 




- Molecule 7: CRISPR type AFERR-associated protein Csf2 (Cas7)

Chain F: 71% 20% 9%




- Molecule 7: CRISPR type AFERR-associated protein Csf2 (Cas7)


Chain G: 65% 12% 23%



- Molecule 8: CRISPR-associated DEAD/DEAH-box helicase CasDinG

Chain L: 12% 60% 21% 18%





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	83553	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	80	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOCONTINUUM (6k x 4k)	Depositor
Maximum map value	1.420	Depositor
Minimum map value	-0.233	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.033	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	426.5984, 426.5984, 426.5984	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.8332, 0.8332, 0.8332	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.42	0/1755	0.59	1/2384 (0.0%)
2	H	0.21	0/1772	0.49	0/2391
3	I	0.69	0/1472	0.91	0/2297
4	J	0.60	0/1121	0.85	0/1730
5	K	0.61	0/1085	0.87	0/1670
6	A	0.18	0/1937	0.47	0/2635
7	C	0.18	0/2531	0.45	0/3432
7	D	0.15	0/2548	0.40	0/3454
7	E	0.20	0/2514	0.45	0/3412
7	F	0.17	0/2509	0.40	0/3405
7	G	0.16	0/2133	0.41	0/2894
8	L	0.35	0/4683	0.60	2/6381 (0.0%)
All	All	0.33	0/26060	0.57	3/36085 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	L	483	PRO	N-CA-C	-8.26	105.09	114.92
8	L	300	TYR	N-CA-CB	-5.58	101.63	109.94
1	B	50	TYR	N-CA-C	-5.46	106.99	113.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1710	0	1694	46	0
2	H	1737	0	1772	32	0
3	I	1314	0	663	51	0
4	J	996	0	537	51	0
5	K	972	0	543	44	0
6	A	1889	0	1869	32	0
7	C	2482	0	2484	52	0
7	D	2499	0	2502	50	0
7	E	2465	0	2458	46	0
7	F	2460	0	2456	46	0
7	G	2090	0	2096	27	0
8	L	4591	0	4611	132	0
9	A	1	0	0	0	0
All	All	25206	0	23685	517	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 517 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:L:381:MET:SD	8:L:395:ALA:HB1	1.70	1.32
3:I:35:A:O2'	3:I:36:U:H5'	1.36	1.25
8:L:194:THR:HG23	8:L:314:LEU:CD2	1.65	1.24
8:L:194:THR:HG23	8:L:314:LEU:HD21	1.19	1.14
4:J:22:DA:H2''	4:J:23:DC:H5''	1.30	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	217/220 (99%)	207 (95%)	10 (5%)	0	100	100
2	H	224/232 (97%)	216 (96%)	7 (3%)	1 (0%)	30	65
6	A	239/241 (99%)	229 (96%)	10 (4%)	0	100	100
7	C	325/359 (90%)	317 (98%)	8 (2%)	0	100	100
7	D	327/359 (91%)	314 (96%)	13 (4%)	0	100	100
7	E	323/359 (90%)	309 (96%)	14 (4%)	0	100	100
7	F	322/359 (90%)	307 (95%)	15 (5%)	0	100	100
7	G	272/359 (76%)	262 (96%)	10 (4%)	0	100	100
8	L	609/748 (81%)	571 (94%)	38 (6%)	0	100	100
All	All	2858/3236 (88%)	2732 (96%)	125 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	220	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	176/177 (99%)	173 (98%)	3 (2%)	53	78
2	H	184/190 (97%)	181 (98%)	3 (2%)	55	79
6	A	208/210 (99%)	208 (100%)	0	100	100
7	C	262/283 (93%)	262 (100%)	0	100	100
7	D	263/283 (93%)	260 (99%)	3 (1%)	65	83
7	E	259/283 (92%)	257 (99%)	2 (1%)	73	86
7	F	259/283 (92%)	257 (99%)	2 (1%)	73	86
7	G	223/283 (79%)	223 (100%)	0	100	100
8	L	473/602 (79%)	468 (99%)	5 (1%)	65	83
All	All	2307/2594 (89%)	2289 (99%)	18 (1%)	70	86

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	L	380	LEU
8	L	595	THR
8	L	476	VAL
7	D	256	LEU
8	L	250	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
7	D	249	HIS
7	E	209	GLN
7	E	68	ASN
7	F	30	ASN
2	H	59	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	I	60/61 (98%)	34 (56%)	4 (6%)

5 of 34 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	I	2	U
3	I	4	A
3	I	9	U
3	I	10	G
3	I	12	A

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	I	12	A
3	I	33	U
3	I	34	G
3	I	38	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-48860. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256



Y Index: 256




Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 253



Y Index: 238




Z Index: 305


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 215 nm^3 ; this corresponds to an approximate mass of 194 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-48860 and PDB model 9N3Y. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8150	0.3920
A	0.8220	0.4100
B	0.8130	0.4000
C	0.8350	0.4260
D	0.8660	0.4410
E	0.8690	0.4450
F	0.8520	0.4370
G	0.8540	0.4610
H	0.7950	0.3950
I	0.9450	0.4110
J	0.6930	0.2520
K	0.9290	0.3850
L	0.6790	0.2770

1.0

0.0

<0.0