



Full wwPDB X-ray Structure Validation Report i

Feb 14, 2024 – 10:57 PM EST

PDB ID : 3N67

Title : Structure of endothelial nitric oxide synthase N368D/V106M double mutant heme domain complexed with 6,6'-(2,2'-(5-amino-1,3-phenylene)bis(ethane-2,1-diyl))bis(4-methylpyridin-2-amine)

Authors : Delker, S.L.; Li, H.; Poulos, T.L.

Deposited on : 2010-05-25

Resolution : 2.09 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 7038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	404	Total	C 3247	N 2061	O 571	S 598	17	0	4	0
1	B	403	Total	C 3238	N 2055	O 572	S 594	17	0	3	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	SEE REMARK 999	UNP P29476
A	106	MET	VAL	engineered mutation	UNP P29476
A	368	ASP	ASN	engineered mutation	UNP P29476
B	100	ARG	CYS	SEE REMARK 999	UNP P29476
B	106	MET	VAL	engineered mutation	UNP P29476
B	368	ASP	ASN	engineered mutation	UNP P29476


- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂⁻).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 6,6'-[{(5-aminobenzene-1,3-diyl)diethane-2,1-diyl]bis(4-methylpyridin-2-amine) (three-letter code: XFN) (formula: C₂₂H₂₇N₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			27	22	5		
4	B	1	Total	C	N	0	0
			27	22	5		

- Molecule 5 is CACODYLIC ACID (three-letter code: CAD) (formula: $C_2H_7AsO_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	As	C	0	0
			3	1	2		
5	B	1	Total	As	C	0	0
			3	1	2		

- Molecule 6 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C₉H₁₅N₅O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O 17 9 5 3	0	0
6	B	1	Total C N O 17 9 5 3	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Zn 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	182	Total O 182 182	0	0
8	B	182	Total O 182 182	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase



- Molecule 1: Nitric oxide synthase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.97 Å 106.61 Å 156.77 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.19 – 2.09 42.70 – 2.79	Depositor EDS
% Data completeness (in resolution range)	94.4 (39.19-2.09) 97.7 (42.70-2.79)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) >$ ¹	1.84 (at 2.81 Å)	Xtriage
Refinement program	REFMAC 5.5.0109, CNS	Depositor
R , R_{free}	0.173 , 0.218 0.189 , 0.225	Depositor DCC
R_{free} test set	1196 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	53.6	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.1	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7038	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, H4B, XFN, HEM, ACT, CAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/3337	0.76	3/4543 (0.1%)
1	B	0.73	1/3327 (0.0%)	0.73	2/4528 (0.0%)
All	All	0.74	1/6664 (0.0%)	0.74	5/9071 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	203	CYS	CB-SG	-5.33	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	ARG	NE-CZ-NH2	-6.67	116.96	120.30
1	B	240	ARG	NE-CZ-NH1	6.37	123.49	120.30
1	A	328	LEU	CA-CB-CG	5.34	127.59	115.30
1	A	458	LEU	CB-CG-CD1	5.28	119.97	111.00
1	A	328	LEU	CB-CG-CD1	5.13	119.72	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3247	0	3139	38	0
1	B	3238	0	3137	38	0
2	A	43	0	30	3	0
2	B	43	0	30	5	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	A	27	0	27	4	0
4	B	27	0	27	2	0
5	A	3	0	0	2	0
5	B	3	0	0	0	0
6	A	17	0	15	2	0
6	B	17	0	15	0	0
7	A	1	0	0	0	0
8	A	182	0	0	4	0
8	B	182	0	0	4	0
All	All	7038	0	6426	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:CYS:SG	5:A:950:CAD:AS	2.46	1.34
1:A:257:ARG:HG3	1:A:257:ARG:HH11	1.12	1.09
1:B:236:ARG:HG3	1:B:349:GLU:HB2	1.62	0.82
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.63	0.79
1:A:257:ARG:HG3	1:A:257:ARG:NH1	1.91	0.77
1:A:126:LEU:HD11	1:A:156:GLU:HB3	1.70	0.71
1:A:72:ARG:HD3	8:A:1070:HOH:O	1.92	0.69
1:A:68:PRO:O	1:B:109:ARG:NH2	2.26	0.67
1:B:330:LEU:HB2	8:B:1189:HOH:O	1.95	0.67
1:A:99:ARG:HG2	1:A:100:ARG:HD2	1.78	0.65
1:B:178:GLN:HE22	1:B:181:ARG:HH11	1.46	0.64
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.81	0.62
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.30	0.62
1:B:378:LEU:HB2	8:B:1191:HOH:O	2.00	0.61
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.84	0.59
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.84	0.59
1:A:246:TRP:CZ2	1:A:302:PRO:HG3	2.38	0.58
1:B:126:LEU:O	1:B:130:ARG:HG3	2.04	0.58
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.87	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.35	0.56
1:A:384:CYS:CB	5:A:950:CAD:AS	3.17	0.53
1:B:99:ARG:HG3	1:B:100:ARG:HD2	1.91	0.53
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.48	0.52
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	1.92	0.52
2:B:500:HEM:HBC2	2:B:500:HEM:HMC1	1.92	0.52
1:A:340:ASN:H	1:A:340:ASN:HD22	1.57	0.51
1:A:126:LEU:HD12	1:A:130:ARG:CZ	2.41	0.51
1:B:281:TRP:CD1	1:B:292:PRO:HG3	2.45	0.51
1:A:104:SER:O	6:A:600:H4B:O10	2.27	0.50
1:B:379:GLU:HB2	8:B:1342:HOH:O	2.12	0.50
1:A:378:LEU:HB2	8:A:1010:HOH:O	2.11	0.50
1:B:343:LEU:HD21	1:B:345:ILE:HD11	1.92	0.49
1:B:249:GLN:NE2	4:B:800:XFN:H03	2.27	0.49
1:B:240:ARG:HD2	1:B:241:GLY:O	2.12	0.48
1:A:472:SER:HA	1:A:473:PRO:C	2.35	0.47
1:A:476:ARG:HD2	8:A:1139:HOH:O	2.12	0.47
1:B:236:ARG:HG2	1:B:349:GLU:O	2.15	0.47
1:A:92:GLN:NE2	8:A:1097:HOH:O	2.46	0.47
1:A:449:TRP:HA	6:A:600:H4B:N1	2.30	0.47
1:A:205:SER:O	1:A:209:MET:HG3	2.14	0.47
1:B:167:LEU:HG	1:B:348:LEU:HD12	1.97	0.47
1:A:374:ARG:HH11	1:A:374:ARG:HG3	1.80	0.46
1:A:455:SER:O	1:A:458:LEU:HB2	2.15	0.46
1:B:231:THR:O	1:B:353:ALA:HA	2.16	0.46
2:A:500:HEM:O2D	4:A:800:XFN:H08	2.16	0.46
1:B:434:ASP:OD2	1:B:438:LYS:HE3	2.15	0.46
1:A:344:GLU:OE1	1:A:476:ARG:NH2	2.49	0.45
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.56	0.45
2:B:500:HEM:HMC1	2:B:500:HEM:CBC	2.46	0.45
1:B:72:ARG:NH1	8:B:1275:HOH:O	2.50	0.44
1:B:249:GLN:NE2	4:B:800:XFN:C03	2.81	0.44
1:A:363:GLU:OE1	4:A:800:XFN:N40	2.51	0.44
2:A:500:HEM:O2D	4:A:800:XFN:C08	2.65	0.44
1:B:185:ARG:NH1	1:B:477:TYR:OH	2.49	0.44
1:A:149:GLU:O	1:A:153:GLN:HG3	2.18	0.43
1:A:374:ARG:HG3	1:A:374:ARG:NH1	2.34	0.43
1:B:372:PRO:HA	1:B:376:ASN:ND2	2.33	0.43
1:A:308:PRO:HA	1:A:309:PRO:HD3	1.92	0.43
1:B:186:CYS:HB2	2:B:500:HEM:ND	2.34	0.42
1:A:423:HIS:HB2	1:B:392:THR:HB	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:GLU:OE1	1:A:166:HIS:HD2	2.02	0.42
1:A:480:ASP:HA	1:A:481:PRO:HD3	1.93	0.42
1:A:451:VAL:HA	1:A:452:PRO:HD3	1.87	0.42
1:B:308:PRO:HA	1:B:309:PRO:HD3	1.90	0.42
1:B:322:LEU:HD13	1:B:324:TRP:CZ2	2.55	0.42
1:B:362:THR:HA	1:B:405:ASN:ND2	2.36	0.41
1:B:452:PRO:HG2	1:B:459:THR:HG21	2.03	0.41
1:B:246:TRP:CH2	1:B:302:PRO:HG3	2.56	0.41
1:B:364:ILE:HA	1:B:368:ASP:HB2	2.02	0.41
2:A:500:HEM:HBA1	4:A:800:XFN:H34	2.02	0.41
1:B:178:GLN:HE22	1:B:181:ARG:NH1	2.15	0.41
1:A:100:ARG:NH1	1:A:102:LEU:HD22	2.36	0.41
1:A:392:THR:HB	1:B:423:HIS:HB2	2.03	0.41
1:B:236:ARG:HG2	1:B:236:ARG:H	1.50	0.41
1:A:366:THR:O	1:A:370:CYS:HB2	2.21	0.41
1:A:431:LYS:O	1:A:434[B]:ASP:HB2	2.20	0.40
1:B:180:TRP:CE3	1:B:192:TRP:HA	2.56	0.40
1:B:273:THR:O	1:B:277:ILE:HG13	2.21	0.40
1:A:156:GLU:H	1:A:156:GLU:HG2	1.77	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	404/444 (91%)	392 (97%)	11 (3%)	1 (0%)	47 49
1	B	402/444 (90%)	392 (98%)	10 (2%)	0	100 100
All	All	806/888 (91%)	784 (97%)	21 (3%)	1 (0%)	51 54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	347/377 (92%)	325 (94%)	22 (6%)	18 15
1	B	346/377 (92%)	327 (94%)	19 (6%)	21 19
All	All	693/754 (92%)	652 (94%)	41 (6%)	20 17

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	126	LEU
1	A	161	SER
1	A	169[A]	GLU
1	A	169[B]	GLU
1	A	257	ARG
1	A	260	ASP
1	A	278	GLN
1	A	287[A]	ARG
1	A	287[B]	ARG
1	A	293	LEU
1	A	294	LEU
1	A	298	PRO
1	A	328	LEU
1	A	340	ASN
1	A	378	LEU
1	A	379	GLU
1	A	416	LYS
1	A	458	LEU
1	A	468	ASN
1	B	99	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	100	ARG
1	B	102	LEU
1	B	106	MET
1	B	109	ARG
1	B	139	SER
1	B	142	ARG
1	B	194	LYS
1	B	236	ARG
1	B	240	ARG
1	B	259	GLN
1	B	262	SER
1	B	264	ARG
1	B	278	GLN
1	B	330	LEU
1	B	340	ASN
1	B	378	LEU
1	B	379	GLU
1	B	389	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	GLN
1	A	166	HIS
1	A	191	GLN
1	A	340	ASN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	259	GLN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ACT	B	860	-	3,3,3	0.79	0	3,3,3	0.73	0
2	HEM	B	500	1	41,50,50	2.15	10 (24%)	45,82,82	1.89	11 (24%)
3	ACT	A	860	-	3,3,3	0.84	0	3,3,3	0.69	0
6	H4B	A	600	-	16,18,18	1.04	1 (6%)	11,26,26	3.01	7 (63%)
5	CAD	B	950	-	0,2,4	-	-	0,1,6	-	-
5	CAD	A	950	-	0,2,4	-	-	0,1,6	-	-
6	H4B	B	600	-	16,18,18	1.55	3 (18%)	11,26,26	2.58	6 (54%)
4	XFN	B	800	-	29,29,29	0.69	0	38,40,40	1.78	6 (15%)
2	HEM	A	500	1	41,50,50	1.85	9 (21%)	45,82,82	1.38	6 (13%)
4	XFN	A	800	-	29,29,29	0.73	0	38,40,40	1.42	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	500	1	-	0/12/54/54	-
6	H4B	A	600	-	-	0/8/17/17	0/2/2/2
6	H4B	B	600	-	-	0/8/17/17	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XFN	B	800	-	-	3/10/10/10	0/3/3/3
2	HEM	A	500	1	-	0/12/54/54	-
4	XFN	A	800	-	-	3/10/10/10	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C3D-C2D	7.73	1.53	1.36
2	A	500	HEM	C3D-C2D	6.34	1.50	1.36
2	B	500	HEM	C3C-C2C	-4.22	1.34	1.40
2	A	500	HEM	C3C-C2C	-3.99	1.34	1.40
2	B	500	HEM	FE-ND	3.80	2.15	1.96
2	B	500	HEM	C3C-CAC	3.64	1.55	1.47
6	B	600	H4B	C4A-C4	-3.57	1.36	1.41
2	A	500	HEM	C3C-CAC	3.49	1.55	1.47
6	B	600	H4B	C7-C6	3.42	1.55	1.52
2	B	500	HEM	CAB-C3B	3.20	1.56	1.47
2	B	500	HEM	CMB-C2B	2.88	1.56	1.50
2	B	500	HEM	FE-NB	2.78	2.10	1.96
2	B	500	HEM	C1B-NB	-2.69	1.35	1.40
2	B	500	HEM	CAA-C2A	2.68	1.56	1.52
2	A	500	HEM	CAB-C3B	2.63	1.54	1.47
2	A	500	HEM	CMA-C3A	2.59	1.57	1.51
2	A	500	HEM	C3B-C2B	-2.39	1.32	1.37
2	A	500	HEM	CMB-C2B	2.30	1.55	1.50
6	B	600	H4B	C4-N3	2.16	1.36	1.33
2	A	500	HEM	C4A-NA	2.15	1.40	1.36
2	A	500	HEM	CMC-C2C	2.14	1.56	1.51
2	B	500	HEM	CMA-C3A	2.08	1.56	1.51
6	A	600	H4B	O9-C9	2.06	1.47	1.43

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	HEM	C4D-ND-C1D	6.64	111.94	105.07
6	A	600	H4B	C8A-C4A-C4	6.20	120.07	114.57
4	B	800	XFN	C39-N40-C35	5.96	122.62	118.10
4	B	800	XFN	C11-N12-C13	5.33	122.14	118.10
2	B	500	HEM	CBA-CAA-C2A	-4.81	104.41	112.62
6	B	600	H4B	N1-C2-N3	-4.17	118.88	125.42
6	A	600	H4B	C2-N3-C4	4.04	122.35	115.93
2	B	500	HEM	C1B-NB-C4B	3.83	109.03	105.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	XFN	C11-N12-C13	3.69	120.90	118.10
2	B	500	HEM	C3B-C2B-C1B	3.53	109.10	106.49
2	A	500	HEM	C4D-ND-C1D	3.50	108.69	105.07
6	A	600	H4B	C4-C4A-N5	3.45	122.01	119.12
6	B	600	H4B	C8A-C4A-C4	3.40	117.59	114.57
6	B	600	H4B	C2-N1-C8A	3.35	122.05	114.54
6	B	600	H4B	C4-C4A-N5	3.34	121.92	119.12
6	A	600	H4B	N1-C2-N3	-3.28	120.27	125.42
4	B	800	XFN	C36-C35-N40	-3.28	119.43	122.90
6	B	600	H4B	N2-C2-N3	3.14	122.13	117.25
4	B	800	XFN	C14-C13-N12	-3.12	119.59	122.90
4	A	800	XFN	C09-C13-N12	3.03	120.47	115.95
4	A	800	XFN	C14-C13-N12	-2.98	119.74	122.90
2	A	500	HEM	CBA-CAA-C2A	-2.88	107.71	112.62
2	A	500	HEM	CAD-C3D-C4D	2.75	129.47	124.66
2	B	500	HEM	CAD-CBD-CGD	-2.70	107.78	113.60
6	B	600	H4B	C2-N3-C4	2.65	120.14	115.93
6	A	600	H4B	C2-N1-C8A	2.63	120.42	114.54
2	B	500	HEM	C4C-CHD-C1D	2.59	125.97	122.56
2	B	500	HEM	CBD-CAD-C3D	-2.58	105.47	112.63
2	B	500	HEM	C4A-C3A-C2A	2.54	108.77	107.00
4	A	800	XFN	N17-C11-N12	2.51	120.46	116.49
2	B	500	HEM	C1D-C2D-C3D	-2.42	104.41	106.96
2	B	500	HEM	C2B-C1B-NB	-2.41	106.98	109.84
4	A	800	XFN	C39-N40-C35	2.34	119.88	118.10
6	A	600	H4B	C4A-C4-N3	-2.33	117.39	124.01
4	A	800	XFN	C34-C35-N40	2.30	119.38	115.95
4	B	800	XFN	C09-C13-N12	2.29	119.36	115.95
2	A	500	HEM	CAD-CBD-CGD	-2.23	108.80	113.60
2	A	500	HEM	C4D-C3D-C2D	-2.16	103.75	106.90
4	B	800	XFN	N17-C11-N12	2.13	119.86	116.49
2	B	500	HEM	C2C-C3C-C4C	2.12	108.38	106.90
2	A	500	HEM	CMB-C2B-C1B	2.03	128.13	125.04
6	A	600	H4B	N2-C2-N3	2.01	120.38	117.25

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	800	XFN	C06-C08-C09-C13
4	A	800	XFN	C01-C06-C08-C09
4	A	800	XFN	C05-C06-C08-C09

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
4	A	800	XFN	C06-C08-C09-C13
4	B	800	XFN	C01-C06-C08-C09
4	B	800	XFN	C05-C06-C08-C09



There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	500	HEM	5	0
6	A	600	H4B	2	0
5	A	950	CAD	2	0
4	B	800	XFN	2	0
2	A	500	HEM	3	0
4	A	800	XFN	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/444 (90%)	1.32	79 (19%) 1 1	29, 42, 65, 84	0
1	B	403/444 (90%)	1.26	73 (18%) 1 1	30, 45, 66, 90	0
All	All	807/888 (90%)	1.29	152 (18%) 1 1	29, 44, 66, 90	0

All (152) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	69	LYS	6.8
1	A	160	ALA	6.6
1	B	259	GLN	6.1
1	A	123	GLU	5.4
1	B	121	PRO	5.3
1	A	259	GLN	5.1
1	A	239	GLY	5.0
1	A	99	ARG	5.0
1	B	263	VAL	4.9
1	B	122	ALA	4.8
1	A	126	LEU	4.8
1	A	121	PRO	4.8
1	A	153	GLN	4.8
1	A	106	MET	4.7
1	A	91	GLN	4.6
1	B	277	ILE	4.5
1	A	127	SER	4.5
1	B	123	GLU	4.5
1	B	280	GLY	4.4
1	B	270	VAL	4.3
1	A	257	ARG	4.2
1	B	106	MET	4.2
1	A	122	ALA	4.2
1	A	159	VAL	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	142	ARG	4.1
1	A	124	GLN	4.0
1	B	262	SER	4.0
1	A	68	PRO	3.9
1	A	161	SER	3.9
1	B	223	ARG	3.9
1	A	130	ARG	3.8
1	A	156	GLU	3.8
1	B	449	TRP	3.6
1	B	141	LYS	3.6
1	A	70	PHE	3.6
1	A	256	TYR	3.6
1	B	304	LEU	3.6
1	A	281	TRP	3.5
1	A	152	LEU	3.5
1	B	261	GLY	3.5
1	A	125	LEU	3.5
1	A	90	SER	3.5
1	A	93	ASP	3.4
1	B	281	TRP	3.4
1	A	157	ALA	3.4
1	A	260	ASP	3.4
1	A	238	PRO	3.4
1	B	260	ASP	3.4
1	A	67	GLY	3.4
1	A	337	ALA	3.3
1	A	414	LEU	3.3
1	A	451	VAL	3.3
1	A	223	ARG	3.2
1	B	70	PHE	3.2
1	A	163	GLY	3.1
1	B	147	ALA	3.0
1	B	258	GLN	3.0
1	A	476	ARG	3.0
1	A	198	PHE	3.0
1	B	338	VAL	3.0
1	B	80	SER	2.9
1	B	99	ARG	2.9
1	B	257	ARG	2.9
1	B	339	SER	2.9
1	A	246	TRP	2.9
1	B	448	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	140	ILE	2.8
1	A	364	ILE	2.8
1	A	304	LEU	2.8
1	A	311	LEU	2.8
1	B	232	VAL	2.8
1	A	258	GLN	2.8
1	A	278	GLN	2.8
1	A	199	ASP	2.7
1	A	232	VAL	2.7
1	B	218	LYS	2.7
1	B	341	MET	2.7
1	A	274	GLU	2.7
1	B	246	TRP	2.7
1	A	270	VAL	2.7
1	B	98	PRO	2.7
1	B	439	ALA	2.7
1	A	225	ASN	2.7
1	B	358	TRP	2.6
1	A	92	GLN	2.6
1	B	390	ARG	2.6
1	A	231	THR	2.6
1	A	449	TRP	2.6
1	A	250	LEU	2.6
1	B	278	GLN	2.6
1	A	427	VAL	2.6
1	A	162	THR	2.6
1	B	146	GLN	2.6
1	B	306	VAL	2.6
1	B	474	ALA	2.6
1	B	248	SER	2.6
1	A	390	ARG	2.5
1	B	434	ASP	2.5
1	B	337	ALA	2.5
1	A	146	GLN	2.5
1	A	155	VAL	2.5
1	B	362	THR	2.5
1	B	438	LYS	2.5
1	A	210	PHE	2.5
1	A	299	ASP	2.4
1	A	230	ILE	2.4
1	A	149	GLU	2.4
1	A	249	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	137	TYR	2.4
1	B	239	GLY	2.4
1	B	130	ARG	2.4
1	B	264	ARG	2.4
1	A	482	TRP	2.4
1	B	81	ILE	2.4
1	B	69	LYS	2.3
1	A	169[A]	GLU	2.3
1	B	276	CYS	2.3
1	A	295	LEU	2.3
1	B	462	PHE	2.3
1	A	302	PRO	2.3
1	A	72	ARG	2.3
1	B	78	LEU	2.3
1	B	126	LEU	2.3
1	A	168	ARG	2.3
1	A	339	SER	2.2
1	B	294	LEU	2.2
1	A	338	VAL	2.2
1	A	128	GLN	2.2
1	B	102	LEU	2.2
1	B	316	PRO	2.2
1	B	282	THR	2.2
1	B	108	PRO	2.2
1	B	414	LEU	2.2
1	B	476	ARG	2.2
1	A	353	ALA	2.1
1	A	301	ALA	2.1
1	A	309	PRO	2.1
1	A	197	VAL	2.1
1	B	417	VAL	2.1
1	B	83	TYR	2.1
1	B	336	PRO	2.1
1	A	240	ARG	2.1
1	B	109	ARG	2.1
1	B	234	PRO	2.1
1	B	451	VAL	2.1
1	A	137	TYR	2.1
1	B	265	GLY	2.1
1	B	323	GLU	2.0
1	A	206	ALA	2.0
1	B	454	ILE	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	186	CYS	2.0
1	B	197	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	XFN	B	800	27/27	0.81	0.37	38,73,93,93	0
4	XFN	A	800	27/27	0.84	0.34	33,67,93,93	0
5	CAD	A	950	3/5	0.93	0.23	61,61,61,64	0
3	ACT	B	860	4/4	0.94	0.45	54,54,56,56	0
3	ACT	A	860	4/4	0.95	0.21	47,47,49,49	0
6	H4B	A	600	17/17	0.95	0.16	35,38,41,42	0
6	H4B	B	600	17/17	0.95	0.16	33,37,40,41	0
2	HEM	B	500	43/43	0.96	0.21	28,34,43,47	0
2	HEM	A	500	43/43	0.97	0.20	27,32,43,48	0
5	CAD	B	950	3/5	0.97	0.13	67,67,68,68	0
7	ZN	A	900	1/1	0.99	0.07	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.