



## wwPDB EM Validation Summary Report ⓘ

Mar 3, 2025 – 10:18 AM EST

PDB ID : 9NJT  
EMDB ID : EMD-49490  
Title : Structure of native octahedral assembly of D. discoideum Odo2  
Authors : Hoogerbrugge, G.; Keatinge-Clay, A.T.; Marcotte, E.M.  
Deposited on : 2025-02-27  
Resolution : 5.72 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev117  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.41.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 5.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	
1	B	439	
1	C	439	
1	D	439	
1	E	439	
1	F	439	
1	G	439	
1	H	439	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	I	439	
1	J	439	
1	K	439	
1	L	439	
1	M	439	
1	N	439	
1	O	439	
1	P	439	
1	Q	439	
1	R	439	
1	S	439	
1	T	439	
1	U	439	
1	V	439	
1	W	439	
1	X	439	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 43224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial.


Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	B	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	C	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	D	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	E	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	F	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	G	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	H	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	I	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	J	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	K	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	L	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	M	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	N	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	O	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	P	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms					AltConf	Trace
1	Q	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	R	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	S	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	T	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	U	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	V	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	W	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		
1	X	230	Total	C	N	O	S	0	0
			1801	1137	319	335	10		






- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial

Chain C: 37% 16% 48%



- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial

Chain D: 39% 13% 48%





- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial

Chain E: 38% 15% 48%

MET ARG SER ALA THR LYS LEU ILE LYS ASN VAL ASN ILE ASN VAL GLN SER ASN MET VAL GLY VAL ARG ASP ARG SER THR SER ARG LEU THR ILE ASN ASN ALA TRP SER ILE THR LYS THR VAL ARG GLN PHE THR VAL ARG SER SER SER SER PHE THR VAL CYS THR SER SER THR SER LEU PHE THR ASP ASN LYS PRO ASN VAL THR

ASN ILE LYS TYR GLN ARG PHE TYR SER GLU SER ALA ASN ASP VAL VAL ILE CYS VAL VAL ASN VAL PRO THR SER MET GLY VAL THR ASP ARG SER LEU THR ILE SER GLY THR ILE ASN VAL ALA THR LYS THR ASN THR VAL GLY VAL ARG

ALA PRO VAL SER GLY THR ILE VAL GLU SER PHE VAL ALA ASN LYS GLU GLY VAL VAL ASN VAL THR VAL THR GLY VAL THR ASP ARG SER THR LYS ILE THR ILE GLY GLU VAL ALA TRP ALA THR LYS THR ASN THR VAL GLY VAL ARG

THR PRO LYS ALA PRO ALA THR ILE VAL GLU SER PHE VAL ALA ASN LYS GLU GLY VAL VAL ASN VAL THR VAL THR GLY VAL THR ASP ARG SER THR LYS ILE THR ILE GLY GLU VAL ALA TRP ALA THR LYS THR ASN THR VAL GLY VAL ARG

K72 I76 A79 S80 K214 K215 K216 D218 N84 D85 I86 V87 Y88 V92 R102 P107 V108 I109 G112 L115 E123 L127 M150 G155 S156 M157 F158 M173 H174 A175 I176 P180 Y181 V182 V183 V188 V189 R190 P191 Y194 L197 I203 R207

V210 T211 F212 L213 K214 K215 K216 D218 N219 L220 L221 L227 L228 E229 L230

- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial

Chain F: 38% 15% 48%

MET ARG SER ALA THR LYS LEU ILE LYS ASN VAL ASN ILE ASN VAL GLN SER ASN MET VAL GLY VAL ARG ASP ARG SER THR SER ARG LEU THR ILE ASN ASN ALA TRP SER ILE THR LYS THR ASN THR VAL GLY VAL ARG GLN PHE THR VAL ARG SER SER SER SER PHE THR SER LEU PHE THR ASP ASN LYS PRO ASN VAL THR

ASN ILE LYS TYR GLN ARG PHE TYR SER GLU SER ALA ASN ASP VAL VAL ILE CYS VAL VAL ASN VAL THR VAL THR GLY VAL THR ASP ARG SER THR LYS ILE THR ILE GLY THR ILE ASN VAL ALA TRP ALA THR LYS THR ASN THR VAL GLY VAL ARG

ALA PRO VAL SER GLY THR ILE VAL GLU SER PHE VAL ALA ASN LYS GLU GLY VAL VAL ASN VAL THR VAL THR GLY VAL THR ASP ARG SER THR LYS ILE THR ILE GLY GLU VAL ALA TRP ALA THR LYS THR ASN THR VAL GLY VAL ARG


THR PRO LYS ALA PRO ALA THR ILE VAL GLU SER PHE VAL ALA ASN LYS GLU GLY VAL VAL ASN VAL THR VAL THR GLY VAL THR ASP ARG SER THR LYS ILE THR ILE GLY GLU VAL ALA TRP ALA THR LYS THR ASN THR VAL GLY VAL ARG

F63 V64 K65 T68 V189 I69 A70 L71 K72 V77 N78 D79 S80 V81 D85 I86 V87 Y88 H89 N90 N91 V92 N93 I94 A100 P101 R102 G103 L104 V105 V106 P107 V108 I109 E123 L127 S128 R132 E139 I142 T147 F158 I163 I170 H174 P180


V183 V188 V189 R190 L197 H201 R202 I203 I204 E205 G206 V210 L213 K214 K215 K216 K217 D218 V219 N222 R225 L230

- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial

48%




48%




48%







- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial




- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial






- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial



- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial












- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial

- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial







- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial



- Molecule 1: Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	836	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	49.0	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.489	Depositor
Minimum map value	-0.202	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.026	Depositor
Recommended contour level	0.147	Depositor
Map size ( $\text{\AA}$ )	447.8464, 447.8464, 447.8464	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.7494, 1.7494, 1.7494	Depositor

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/1828	0.51	0/2466
1	B	0.25	0/1828	0.52	0/2466
1	C	0.25	0/1828	0.52	0/2466
1	D	0.25	0/1828	0.53	0/2466
1	E	0.25	0/1828	0.53	0/2466
1	F	0.25	0/1828	0.51	0/2466
1	G	0.25	0/1828	0.51	0/2466
1	H	0.25	0/1828	0.53	0/2466
1	I	0.25	0/1828	0.52	0/2466
1	J	0.25	0/1828	0.53	0/2466
1	K	0.26	0/1828	0.53	0/2466
1	L	0.25	0/1828	0.51	0/2466
1	M	0.25	0/1828	0.53	0/2466
1	N	0.25	0/1828	0.53	1/2466 (0.0%)
1	O	0.25	0/1828	0.52	0/2466
1	P	0.25	0/1828	0.53	0/2466
1	Q	0.25	0/1828	0.54	0/2466
1	R	0.25	0/1828	0.51	0/2466
1	S	0.25	0/1828	0.51	0/2466
1	T	0.26	0/1828	0.56	0/2466
1	U	0.25	0/1828	0.52	0/2466
1	V	0.25	0/1828	0.52	0/2466
1	W	0.26	0/1828	0.53	0/2466
1	X	0.25	0/1828	0.51	0/2466
All	All	0.25	0/43872	0.52	1/59184 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	178	ASP	CB-CG-OD1	5.28	123.05	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1801	0	1850	49	0
1	B	1801	0	1850	36	0
1	C	1801	0	1850	49	0
1	D	1801	0	1850	44	0
1	E	1801	0	1850	54	0
1	F	1801	0	1850	49	0
1	G	1801	0	1850	32	0
1	H	1801	0	1850	38	0
1	I	1801	0	1850	40	0
1	J	1801	0	1850	36	0
1	K	1801	0	1850	49	0
1	L	1801	0	1850	47	0
1	M	1801	0	1850	50	0
1	N	1801	0	1850	46	0
1	O	1801	0	1850	50	0
1	P	1801	0	1850	46	0
1	Q	1801	0	1850	42	0
1	R	1801	0	1850	51	0
1	S	1801	0	1850	45	0
1	T	1801	0	1850	55	0
1	U	1801	0	1850	43	0
1	V	1801	0	1850	54	0
1	W	1801	0	1850	51	0
1	X	1801	0	1850	44	0
All	All	43224	0	44400	971	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 971 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Q:33:LEU:HD13	1:Q:217:LYS:HG2	1.51	0.92

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:150:ASN:HA	1:X:173:MET:HB2	1.60	0.84
1:J:150:ASN:HA	1:J:173:MET:HB2	1.64	0.80
1:V:35:MET:HG3	1:V:176:ILE:HG12	1.67	0.76
1:M:150:ASN:HA	1:M:173:MET:HB2	1.68	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/439 (52%)	224 (98%)	4 (2%)	0	100	100
1	B	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	C	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	D	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	E	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	F	228/439 (52%)	227 (100%)	1 (0%)	0	100	100
1	G	228/439 (52%)	224 (98%)	4 (2%)	0	100	100
1	H	228/439 (52%)	222 (97%)	6 (3%)	0	100	100
1	I	228/439 (52%)	224 (98%)	4 (2%)	0	100	100
1	J	228/439 (52%)	222 (97%)	5 (2%)	1 (0%)	30	68
1	K	228/439 (52%)	222 (97%)	6 (3%)	0	100	100
1	L	228/439 (52%)	224 (98%)	4 (2%)	0	100	100
1	M	228/439 (52%)	221 (97%)	7 (3%)	0	100	100
1	N	228/439 (52%)	227 (100%)	1 (0%)	0	100	100
1	O	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	P	228/439 (52%)	222 (97%)	6 (3%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	R	228/439 (52%)	222 (97%)	5 (2%)	1 (0%)	30	68
1	S	228/439 (52%)	223 (98%)	5 (2%)	0	100	100
1	T	228/439 (52%)	222 (97%)	6 (3%)	0	100	100
1	U	228/439 (52%)	224 (98%)	4 (2%)	0	100	100
1	V	228/439 (52%)	221 (97%)	7 (3%)	0	100	100
1	W	228/439 (52%)	224 (98%)	4 (2%)	0	100	100
1	X	228/439 (52%)	225 (99%)	3 (1%)	0	100	100
All	All	5472/10536 (52%)	5358 (98%)	112 (2%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	J	83	GLU
1	R	83	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/375 (53%)	198 (100%)	0	100	100
1	B	198/375 (53%)	198 (100%)	0	100	100
1	C	198/375 (53%)	198 (100%)	0	100	100
1	D	198/375 (53%)	198 (100%)	0	100	100
1	E	198/375 (53%)	198 (100%)	0	100	100
1	F	198/375 (53%)	198 (100%)	0	100	100
1	G	198/375 (53%)	198 (100%)	0	100	100
1	H	198/375 (53%)	198 (100%)	0	100	100
1	I	198/375 (53%)	198 (100%)	0	100	100
1	J	198/375 (53%)	197 (100%)	1 (0%)	86	89

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	198/375 (53%)	196 (99%)	2 (1%)	73	81
1	L	198/375 (53%)	198 (100%)	0	100	100
1	M	198/375 (53%)	198 (100%)	0	100	100
1	N	198/375 (53%)	198 (100%)	0	100	100
1	O	198/375 (53%)	198 (100%)	0	100	100
1	P	198/375 (53%)	198 (100%)	0	100	100
1	Q	198/375 (53%)	198 (100%)	0	100	100
1	R	198/375 (53%)	198 (100%)	0	100	100
1	S	198/375 (53%)	197 (100%)	1 (0%)	86	89
1	T	198/375 (53%)	197 (100%)	1 (0%)	86	89
1	U	198/375 (53%)	198 (100%)	0	100	100
1	V	198/375 (53%)	198 (100%)	0	100	100
1	W	198/375 (53%)	198 (100%)	0	100	100
1	X	198/375 (53%)	198 (100%)	0	100	100
All	All	4752/9000 (53%)	4747 (100%)	5 (0%)	92	95

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	102	ARG
1	K	102	ARG
1	K	207	ARG
1	S	102	ARG
1	T	16	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	Q	201	HIS
1	P	167	GLN
1	O	95	ASN
1	L	201	HIS
1	O	167	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-49490. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


### 6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 106




Y Index: 106




Z Index: 105


### 6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 255


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.147. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_49490\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 455 nm<sup>3</sup>; this corresponds to an approximate mass of 411 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.175 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.175 \text{ \AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.72	-	-
Author-provided FSC curve	5.72	7.63	6.22
Unmasked-calculated*	8.47	16.31	9.03

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.47 differs from the reported value 5.72 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-49490 and PDB model 9NJT. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.147 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.147).



















































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.147) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8210	 0.3250
A	 0.8150	 0.3230
B	 0.8240	 0.3270
C	 0.8230	 0.3270
D	 0.8210	 0.3270
E	 0.8250	 0.3250
F	 0.8190	 0.3240
G	 0.8220	 0.3280
H	 0.8190	 0.3260
I	 0.8250	 0.3290
J	 0.8190	 0.3250
K	 0.8150	 0.3270
L	 0.8180	 0.3240
M	 0.8160	 0.3230
N	 0.8200	 0.3270
O	 0.8220	 0.3250
P	 0.8190	 0.3220
Q	 0.8230	 0.3230
R	 0.8230	 0.3260
S	 0.8210	 0.3240
T	 0.8180	 0.3240
U	 0.8230	 0.3250
V	 0.8210	 0.3240
W	 0.8190	 0.3210
X	 0.8230	 0.3290

