



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 26, 2024 – 06:44 PM EDT

PDB ID : 3NKM
Title : Crystal structure of mouse autotaxin
Authors : Nishimasu, H.; Ishitani, R.; Mihara, E.; Takagi, J.; Aoki, J.; Nureki, O.
Deposited on : 2010-06-20
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	831	<div> <div>85%</div> <div>12%</div> <div>.</div> </div>
2	B	2	<div> <div>50%</div> <div>50%</div> </div>
2	D	2	<div> <div>50%</div> <div>50%</div> </div>
3	C	6	<div> <div>83%</div> <div>17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	EDO	A	1026	-	-	X	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 7391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	805	Total	C	N	O	S	0	4	0
			6461	4104	1104	1203	50			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LYS	SEE REMARK 999	UNP Q9R1E6
A	?	-	VAL	SEE REMARK 999	UNP Q9R1E6
A	?	-	GLU	SEE REMARK 999	UNP Q9R1E6
A	?	-	PRO	SEE REMARK 999	UNP Q9R1E6
A	859	SER	-	expression tag	UNP Q9R1E6
A	860	ARG	-	expression tag	UNP Q9R1E6
A	861	GLU	-	expression tag	UNP Q9R1E6
A	862	ASN	-	expression tag	UNP Q9R1E6
A	863	LEU	-	expression tag	UNP Q9R1E6
A	864	TYR	-	expression tag	UNP Q9R1E6
A	865	PHE	-	expression tag	UNP Q9R1E6
A	866	GLN	-	expression tag	UNP Q9R1E6

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		


- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		

- Molecule 7 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	K	0	0
			1	1		

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is (2R)-2-hydroxy-3-(phosphonoxy)propyl (9E)-octadec-9-enoate (three-letter code: NKP) (formula: C₂₁H₄₁O₇P).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	C	0	0
			6	6		

- Molecule 10 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	S	0	0
			3	1	1	1		
10	A	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 11 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			4	2	2		
11	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0
11	A	1	Total C O 4 2 2	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	712	Total O 712 712	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3
MAN4
MAN5
MAN6

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.59Å 94.73Å 75.65Å 90.00° 94.05° 90.00°	Depositor
Resolution (Å)	34.18 – 2.00 34.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.3 (34.18-2.00) 97.3 (34.18-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.6_289	Depositor
R, R_{free}	0.163 , 0.213 0.159 , 0.208	Depositor DCC
R_{free} test set	2875 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7391	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NA, NAG, MAN, SO4, BMA, ZN, K, SCN, NKP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/6659	0.51	0/9049

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6461	0	6168	81	0
2	B	28	0	25	0	0
2	D	28	0	25	0	0
3	C	72	0	61	0	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	5	0	0	0	0
9	A	6	0	11	0	0
10	A	6	0	0	1	0
11	A	68	0	102	19	0
12	A	712	0	0	6	0
All	All	7391	0	6392	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:675:PHE:H	11:A:1012:EDO:H11	1.10	1.11
1:A:118:HIS:HD2	1:A:128:ASP:HB3	1.40	0.84
1:A:675:PHE:N	11:A:1012:EDO:H11	1.94	0.78
1:A:424:MET:HG3	12:A:1560:HOH:O	1.84	0.77
1:A:368:ARG:HH22	11:A:1026:EDO:C2	2.00	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	807/831 (97%)	779 (96%)	25 (3%)	3 (0%)	30 27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	GLU
1	A	109	GLU
1	A	386	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	721/756 (95%)	717 (99%)	4 (1%)	84 88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	LEU
1	A	128	ASP
1	A	400	LEU
1	A	481	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	827	HIS
1	A	802	ASN
1	A	374	ASN
1	A	348	HIS
1	A	667	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.49	0	17,19,21	0.98	1 (5%)
2	NAG	B	2	2	14,14,15	0.54	0	17,19,21	0.71	0
3	NAG	C	1	3,1	14,14,15	0.69	0	17,19,21	1.08	2 (11%)
3	NAG	C	2	3	14,14,15	0.57	0	17,19,21	0.77	0
3	BMA	C	3	3	11,11,12	0.62	0	15,15,17	0.71	0
3	MAN	C	4	3	11,11,12	0.68	0	15,15,17	0.83	0
3	MAN	C	5	3	11,11,12	0.58	0	15,15,17	0.71	0
3	MAN	C	6	3	11,11,12	0.56	0	15,15,17	0.73	0
2	NAG	D	1	2,1	14,14,15	0.64	0	17,19,21	0.95	1 (5%)
2	NAG	D	2	2	14,14,15	0.53	0	17,19,21	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	MAN	C	6	3	-	0/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	2.86	116.02	112.19
3	C	1	NAG	C3-C4-C5	2.35	114.49	110.23
2	D	1	NAG	O5-C1-C2	-2.29	107.75	111.29
3	C	1	NAG	O5-C1-C2	-2.00	108.19	111.29

There are no chirality outliers.


5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
3	C	3	BMA	O5-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6
2	D	2	NAG	C8-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

Of 26 ligands modelled in this entry, 5 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	EDO	A	1024	-	3,3,3	0.60	0	2,2,2	0.30	0
11	EDO	A	1012	-	3,3,3	0.32	0	2,2,2	0.48	0
11	EDO	A	1014	-	3,3,3	0.53	0	2,2,2	0.06	0
11	EDO	A	1013	-	3,3,3	0.43	0	2,2,2	0.52	0
11	EDO	A	1019	-	3,3,3	0.47	0	2,2,2	0.30	0
9	NKP	A	1007	-	5,5,28	0.39	0	4,4,32	0.52	0
11	EDO	A	1011	-	3,3,3	0.63	0	2,2,2	0.04	0
11	EDO	A	1023	-	3,3,3	0.50	0	2,2,2	0.41	0
8	SO4	A	1006	4	4,4,4	0.23	0	6,6,6	0.30	0
10	SCN	A	1008	-	1,2,2	0.97	0	0,1,1	-	-
11	EDO	A	1015	-	3,3,3	0.45	0	2,2,2	0.43	0
11	EDO	A	1025	-	3,3,3	0.46	0	2,2,2	0.22	0
11	EDO	A	1021	-	3,3,3	0.48	0	2,2,2	0.33	0
11	EDO	A	1022	-	3,3,3	0.50	0	2,2,2	0.35	0
11	EDO	A	1016	-	3,3,3	0.50	0	2,2,2	0.33	0
10	SCN	A	1009	-	1,2,2	0.67	0	0,1,1	-	-
11	EDO	A	1010	-	3,3,3	0.48	0	2,2,2	0.47	0
11	EDO	A	1026	-	3,3,3	0.39	0	2,2,2	0.48	0
11	EDO	A	1020	-	3,3,3	0.43	0	2,2,2	0.43	0
11	EDO	A	1018	-	3,3,3	0.51	0	2,2,2	0.13	0
11	EDO	A	1017	-	3,3,3	0.40	0	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	EDO	A	1010	-	-	1/1/1/1	-
11	EDO	A	1011	-	-	0/1/1/1	-
11	EDO	A	1024	-	-	1/1/1/1	-
11	EDO	A	1023	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	EDO	A	1012	-	-	0/1/1/1	-
11	EDO	A	1015	-	-	0/1/1/1	-
11	EDO	A	1022	-	-	0/1/1/1	-
11	EDO	A	1014	-	-	0/1/1/1	-
11	EDO	A	1025	-	-	1/1/1/1	-
11	EDO	A	1026	-	-	0/1/1/1	-
11	EDO	A	1013	-	-	0/1/1/1	-
11	EDO	A	1020	-	-	1/1/1/1	-
11	EDO	A	1016	-	-	0/1/1/1	-
11	EDO	A	1019	-	-	1/1/1/1	-
9	NKP	A	1007	-	-	0/3/3/28	-
11	EDO	A	1018	-	-	0/1/1/1	-
11	EDO	A	1017	-	-	0/1/1/1	-
11	EDO	A	1021	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	A	1010	EDO	O1-C1-C2-O2
11	A	1019	EDO	O1-C1-C2-O2
11	A	1025	EDO	O1-C1-C2-O2
11	A	1020	EDO	O1-C1-C2-O2
11	A	1024	EDO	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	A	1012	EDO	3	0
11	A	1013	EDO	1	0
11	A	1025	EDO	3	0
10	A	1009	SCN	1	0
11	A	1010	EDO	1	0
11	A	1026	EDO	10	0
11	A	1017	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	805/831 (96%)	-0.50	5 (0%) 85 85	9, 22, 47, 69	4 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	51	TRP	2.5
1	A	71	PRO	2.3
1	A	556	MET	2.2
1	A	54	THR	2.2
1	A	465	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	2	14/15	0.76	0.13	44,53,61,62	0
2	NAG	B	2	14/15	0.78	0.12	60,67,70,71	0
3	BMA	C	3	11/12	0.83	0.10	37,41,45,47	0
2	NAG	B	1	14/15	0.86	0.10	49,54,58,62	0
3	MAN	C	4	11/12	0.87	0.08	36,38,43,51	0
3	MAN	C	5	11/12	0.87	0.12	33,35,40,50	0
2	NAG	D	1	14/15	0.93	0.09	17,26,37,41	0

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MAN	C	6	11/12	0.95	0.08	23,31,36,36	0
3	NAG	C	2	14/15	0.96	0.06	19,24,29,37	0
3	NAG	C	1	14/15	0.98	0.05	9,14,19,20	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	NKP	A	1007	6/29	0.68	0.19	32,34,37,38	0
7	K	A	1005	1/1	0.81	0.13	38,38,38,38	0
10	SCN	A	1009	3/3	0.87	0.15	11,11,34,36	0
11	EDO	A	1011	4/4	0.88	0.14	19,22,26,28	0
11	EDO	A	1010	4/4	0.91	0.14	21,27,30,30	0
11	EDO	A	1013	4/4	0.92	0.11	26,27,27,29	0
11	EDO	A	1022	4/4	0.92	0.09	26,31,35,36	0
11	EDO	A	1025	4/4	0.92	0.12	22,25,29,41	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
11	EDO	A	1018	4/4	0.93	0.10	18,25,28,36	0
11	EDO	A	1019	4/4	0.93	0.07	25,25,32,33	0
11	EDO	A	1012	4/4	0.93	0.08	19,22,23,27	0
11	EDO	A	1016	4/4	0.93	0.09	20,26,26,28	0
11	EDO	A	1026	4/4	0.93	0.13	24,24,25,28	0
11	EDO	A	1020	4/4	0.94	0.10	17,23,28,33	0
11	EDO	A	1017	4/4	0.95	0.09	13,25,25,26	0
11	EDO	A	1024	4/4	0.95	0.10	13,15,15,26	0
11	EDO	A	1014	4/4	0.95	0.08	18,23,24,24	0
11	EDO	A	1021	4/4	0.95	0.09	30,30,33,41	0
11	EDO	A	1015	4/4	0.97	0.10	21,22,23,23	0
11	EDO	A	1023	4/4	0.98	0.04	13,14,15,16	0
10	SCN	A	1008	3/3	0.98	0.08	22,22,27,38	0
8	SO4	A	1006	5/5	0.98	0.06	13,18,24,28	0
6	NA	A	1004	1/1	0.98	0.05	21,21,21,21	0
5	CA	A	1003	1/1	1.00	0.02	19,19,19,19	0
4	ZN	A	1001	1/1	1.00	0.01	21,21,21,21	0
4	ZN	A	1002	1/1	1.00	0.02	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.