



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 01:30 PM EDT

PDB ID : 6O16
Title : Crystal structure of murine DHX37 in complex with RNA
Authors : Boneberg, F.; Brandmann, T.; Kobel, L.; van den Heuvel, J.; Bargsten, K.; Bammert, L.; Kutay, U.; Jinek, M.
Deposited on : 2019-02-18
Resolution : 2.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)
RNA backbone	3102	1121 (3.16-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	975	 6% 53% 27% 18%
1	B	975	 5% 54% 26% 18%
2	C	10	 30% 40% 30%
2	D	10	 30% 50% 20%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DEAH (Asp-Glu-Ala-His) box polypeptide 37.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	798	Total	C	N	O	S	0	0	0
			6310	4053	1090	1129	38			
1	B	797	Total	C	N	O	S	0	0	0
			6295	4050	1087	1119	39			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	176	SER	-	expression tag	UNP Q6NZL1
A	177	ASN	-	expression tag	UNP Q6NZL1
A	178	ALA	-	expression tag	UNP Q6NZL1
B	176	SER	-	expression tag	UNP Q6NZL1
B	177	ASN	-	expression tag	UNP Q6NZL1
B	178	ALA	-	expression tag	UNP Q6NZL1

- Molecule 2 is a RNA chain called RNA (5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*UP*U)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			197	90	20	78	9			
2	D	10	Total	C	N	O	P	0	0	0
			197	90	20	78	9			


- Molecule 1: DEAH (Asp-Glu-Ala-His) box polypeptide 37





• Molecule 1: DEAH (Asp-Glu-Ala-His) box polypeptide 37






● Molecule 2: RNA (5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*UP*U)-3')



● Molecule 2: RNA (5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*UP*U)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.83Å 137.59Å 94.22Å 90.00° 93.34° 90.00°	Depositor
Resolution (Å)	47.03 – 2.88 47.03 – 2.88	Depositor EDS
% Data completeness (in resolution range)	97.7 (47.03-2.88) 84.5 (47.03-2.88)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 2.86Å)	Xtriage
Refinement program	PHENIX (1.14_3260)	Depositor
R, R_{free}	0.270 , 0.295 0.270 , 0.291	Depositor DCC
R_{free} test set	2388 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	64.7	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	12999	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/6451	0.49	0/8747
1	B	0.30	0/6438	0.48	0/8731
2	C	0.30	0/216	0.79	0/332
2	D	0.29	0/216	0.87	0/332
All	All	0.31	0/13321	0.50	0/18142

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6310	0	6453	211	0
1	B	6295	0	6460	201	0
2	C	197	0	102	7	0
2	D	197	0	102	6	0
All	All	12999	0	13117	413	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 413 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:634:VAL:HG23	1:B:653:VAL:HG22	1.48	0.95
1:A:295:SER:HB3	1:A:363:LYS:HB2	1.48	0.94
1:A:634:VAL:HG23	1:A:653:VAL:HG12	1.56	0.87
1:A:249:PRO:HG3	1:A:430:SER:HA	1.57	0.85
1:A:791:PRO:HB2	1:A:907:MET:SD	2.19	0.82

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	780/975 (80%)	758 (97%)	22 (3%)	0	100	100
1	B	779/975 (80%)	751 (96%)	28 (4%)	0	100	100
All	All	1559/1950 (80%)	1509 (97%)	50 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	690/836 (82%)	656 (95%)	34 (5%)	25	55
1	B	690/836 (82%)	650 (94%)	40 (6%)	20	48
All	All	1380/1672 (82%)	1306 (95%)	74 (5%)	22	51

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	784	SER
1	B	1053	ARG
1	B	869	GLN
1	B	1011	LEU
1	A	874	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	438	HIS
1	A	440	ASN
1	A	660	GLN
1	B	280	GLN
1	B	660	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	9/10 (90%)	5 (55%)	0
2	D	9/10 (90%)	3 (33%)	1 (11%)
All	All	18/20 (90%)	8 (44%)	1 (5%)

5 of 8 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	U
2	C	4	U
2	C	5	U
2	C	8	U
2	C	9	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	D	4	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	798/975 (81%)	0.48	58 (7%) 15 11	30, 60, 101, 144	0
1	B	797/975 (81%)	0.46	53 (6%) 18 14	36, 63, 100, 132	0
2	C	10/10 (100%)	0.28	0 100 100	49, 57, 93, 100	0
2	D	10/10 (100%)	0.34	0 100 100	54, 66, 83, 87	0
All	All	1615/1970 (81%)	0.47	111 (6%) 16 13	30, 61, 100, 144	0

The worst 5 of 111 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	VAL	7.0
1	B	483	ARG	6.8
1	B	1073	PHE	5.7
1	A	294	ASP	5.2
1	B	667	ARG	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.