



wwPDB EM Validation Summary Report ⓘ

May 26, 2025 – 12:49 PM EDT

PDB ID : 6OAX / pdb_00006oax
EMDB ID : EMD-20004
Title : Structure of the hyperactive ClpB mutant K476C, bound to casein, pre-state
Authors : Rizo, A.R.; Lin, J.-B.; Gates, S.N.; Tse, E.; Bart, S.M.; Castellano, L.M.;
Dimaio, F.; Shorter, J.; Southworth, D.R.
Deposited on : 2019-03-18
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0rc1
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	867	<div> <div>15%</div> <div>51%</div> <div>15%</div> <div>33%</div> </div>
1	B	867	<div> <div>49%</div> <div>18%</div> <div>33%</div> </div>
1	C	867	<div> <div>49%</div> <div>18%</div> <div>33%</div> </div>
1	D	867	<div> <div>48%</div> <div>19%</div> <div>33%</div> </div>
1	E	867	<div> <div>7%</div> <div>50%</div> <div>16%</div> <div>33%</div> </div>
1	F	867	<div> <div>45%</div> <div>54%</div> <div>11%</div> <div>34%</div> </div>
2	P	26	<div> <div>8%</div> <div>73%</div> <div>27%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AGS	C	901	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 27871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hyperactive disaggregase ClpB.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	C	581	Total	C	N	O	S	0	0
			4577	2872	830	859	16		
1	F	571	Total	C	N	O	S	0	0
			4518	2837	819	847	15		
1	E	579	Total	C	N	O	S	0	0
			4555	2858	822	859	16		
1	D	581	Total	C	N	O	S	0	0
			4577	2872	830	859	16		
1	B	581	Total	C	N	O	S	0	0
			4577	2872	830	859	16		
1	A	581	Total	C	N	O	S	0	0
			4577	2872	830	859	16		

- Molecule 2 is a protein called Alpha-S1-casein.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	P	26	Total	C	N	O	0	0
			130	78	26	26		

- Molecule 3 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: C₁₀H₁₆N₅O₁₂P₃S).




Mol	Chain	Residues	Atoms						AltConf
3	C	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	C	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	E	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	E	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	D	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	D	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	B	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	B	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	A	1	Total 31	C 10	N 5	O 12	P 3	S 1	0

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					AltConf
4	F	1	Total	C	N	O	P	0
			27	10	5	10	2	
4	F	1	Total	C	N	O	P	0
			27	10	5	10	2	
4	A	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 1: Hyperactive disaggregase ClpB










• Molecule 1: Hyperactive disaggregase ClpB

Chain B:






• Molecule 1: Hyperactive disaggregase ClpB



• Molecule 2: Alpha-S1-casein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	712910	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	56	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	7.873	Depositor
Minimum map value	-3.901	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.189	Depositor
Recommended contour level	0.5	Depositor
Map size (Å)	264.192, 264.192, 264.192	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.032, 1.032, 1.032	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, AGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	1/4644 (0.0%)	0.70	4/6264 (0.1%)
1	B	0.55	0/4644	0.68	1/6264 (0.0%)
1	C	0.63	0/4644	0.72	1/6264 (0.0%)
1	D	0.60	0/4644	0.71	1/6264 (0.0%)
1	E	0.42	0/4623	0.67	2/6238 (0.0%)
1	F	0.27	0/4584	0.72	1/6183 (0.0%)
All	All	0.49	1/27783 (0.0%)	0.70	10/37477 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	815	ARG	C-N	6.14	1.40	1.34

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	337	PHE	N-CA-C	8.09	121.97	110.23
1	C	850	GLU	N-CA-C	-6.42	100.71	109.95
1	F	343	VAL	N-CA-C	-6.03	105.80	113.22
1	A	215	ILE	N-CA-C	-6.03	106.63	111.81
1	B	548	VAL	N-CA-C	-5.83	106.81	112.29

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4577	0	4650	106	0
1	B	4577	0	4650	153	0
1	C	4577	0	4650	169	0
1	D	4577	0	4650	156	0
1	E	4555	0	4615	112	0
1	F	4518	0	4590	80	0
2	P	130	0	33	7	0
3	A	31	0	12	1	0
3	B	62	0	24	5	0
3	C	62	0	24	12	0
3	D	62	0	24	7	0
3	E	62	0	24	3	0
4	A	27	0	12	2	0
4	F	54	0	24	2	0
All	All	27871	0	27982	713	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 713 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:282:THR:HG22	1:B:293:MET:CG	1.53	1.36
1:C:732:LEU:CD1	1:C:740:LEU:HD22	1.59	1.31
1:D:204:LEU:CD2	1:D:215:ILE:HB	1.59	1.31
1:D:204:LEU:HD21	1:D:215:ILE:CB	1.60	1.29
1:C:386:LEU:HB3	1:C:387:PRO:CD	1.66	1.22

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	577/867 (67%)	520 (90%)	55 (10%)	2 (0%)	37	66
1	B	577/867 (67%)	544 (94%)	32 (6%)	1 (0%)	44	73
1	C	577/867 (67%)	537 (93%)	37 (6%)	3 (0%)	25	56
1	D	577/867 (67%)	542 (94%)	34 (6%)	1 (0%)	44	73
1	E	575/867 (66%)	529 (92%)	42 (7%)	4 (1%)	19	49
1	F	565/867 (65%)	497 (88%)	67 (12%)	1 (0%)	44	73
All	All	3448/5202 (66%)	3169 (92%)	267 (8%)	12 (0%)	38	66

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	386	LEU
1	E	340	GLU
1	E	292	ALA
1	E	341	PRO
1	E	338	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	487/726 (67%)	483 (99%)	4 (1%)	79	93
1	B	487/726 (67%)	480 (99%)	7 (1%)	62	86
1	C	487/726 (67%)	481 (99%)	6 (1%)	67	89
1	D	487/726 (67%)	485 (100%)	2 (0%)	89	97
1	E	485/726 (67%)	478 (99%)	7 (1%)	62	86
1	F	483/726 (66%)	482 (100%)	1 (0%)	92	98
All	All	2916/4356 (67%)	2889 (99%)	27 (1%)	74	92

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	203	VAL
1	B	283	MET
1	A	246	VAL
1	B	203	VAL
1	B	288	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	822	GLN
1	A	195	GLN
1	B	192	GLN
1	B	726	GLN
1	A	297	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	F	902	-	24,29,29	0.88	0	29,45,45	1.65	4 (13%)
3	AGS	B	902	-	28,33,33	1.28	3 (10%)	31,52,52	1.10	2 (6%)
3	AGS	A	901	-	28,33,33	1.03	3 (10%)	31,52,52	1.07	2 (6%)
3	AGS	B	901	-	28,33,33	1.11	2 (7%)	31,52,52	0.87	2 (6%)
3	AGS	D	901	-	28,33,33	1.35	5 (17%)	31,52,52	0.99	4 (12%)
4	ADP	F	901	-	24,29,29	0.88	0	29,45,45	1.21	2 (6%)
3	AGS	C	901	-	28,33,33	1.74	5 (17%)	31,52,52	1.23	3 (9%)
4	ADP	A	902	-	24,29,29	0.93	1 (4%)	29,45,45	1.26	3 (10%)
3	AGS	C	902	-	28,33,33	1.37	4 (14%)	31,52,52	1.03	3 (9%)
3	AGS	E	902	-	28,33,33	1.19	3 (10%)	31,52,52	0.97	2 (6%)
3	AGS	D	902	-	28,33,33	1.45	4 (14%)	31,52,52	1.11	3 (9%)
3	AGS	E	901	-	28,33,33	1.05	2 (7%)	31,52,52	0.86	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	F	902	-	-	3/12/32/32	0/3/3/3
3	AGS	B	902	-	-	0/17/38/38	0/3/3/3
3	AGS	A	901	-	-	3/17/38/38	0/3/3/3
3	AGS	B	901	-	-	6/17/38/38	0/3/3/3
3	AGS	D	901	-	-	3/17/38/38	0/3/3/3
4	ADP	F	901	-	-	0/12/32/32	0/3/3/3
3	AGS	C	901	-	-	6/17/38/38	0/3/3/3
4	ADP	A	902	-	-	6/12/32/32	0/3/3/3
3	AGS	C	902	-	-	1/17/38/38	0/3/3/3
3	AGS	E	902	-	-	1/17/38/38	0/3/3/3
3	AGS	D	902	-	-	2/17/38/38	0/3/3/3
3	AGS	E	901	-	-	8/17/38/38	0/3/3/3

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	901	AGS	PB-O3A	-5.24	1.53	1.59
3	C	901	AGS	PB-O3B	-4.54	1.54	1.59
3	D	902	AGS	PA-O3A	-4.03	1.55	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	901	AGS	PA-O3A	-3.64	1.55	1.59
3	C	902	AGS	PB-O3A	-3.51	1.55	1.59

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	902	ADP	O4'-C1'-N9	4.64	114.90	108.75
3	C	901	AGS	PB-O3B-PG	-4.50	116.69	133.17
4	F	902	ADP	C4'-O4'-C1'	-4.32	105.97	109.92
3	A	901	AGS	PB-O3B-PG	-4.28	117.52	133.17
3	B	902	AGS	PB-O3B-PG	-3.89	118.94	133.17

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	901	AGS	C5'-O5'-PA-O1A
3	C	901	AGS	C5'-O5'-PA-O3A
3	E	901	AGS	PB-O3B-PG-O3G
3	D	902	AGS	PB-O3B-PG-O2G
3	D	902	AGS	PB-O3B-PG-O3G


There are no ring outliers.

12 monomers are involved in 32 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	902	ADP	1	0
3	B	902	AGS	1	0
3	A	901	AGS	1	0
3	B	901	AGS	4	0
3	D	901	AGS	3	0
4	F	901	ADP	1	0
3	C	901	AGS	10	0
4	A	902	ADP	2	0
3	C	902	AGS	2	0
3	E	902	AGS	1	0
3	D	902	AGS	4	0
3	E	901	AGS	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In


addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand AGS B 902









Ligand AGS D 901




Ligand ADP F 901




Ligand AGS C 901





Ligand ADP A 902






Ligand AGS E 902







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20004. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 120



Y Index: 106




Z Index: 131


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 162 nm^3 ; this corresponds to an approximate mass of 147 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.345 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20004 and PDB model 6OAX. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.5).

9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7180	<div></div> 0.5000
A	<div></div> 0.6370	<div></div> 0.4660
B	<div></div> 0.8440	<div></div> 0.5490
C	<div></div> 0.8860	<div></div> 0.5720
D	<div></div> 0.8820	<div></div> 0.5730
E	<div></div> 0.7580	<div></div> 0.5030
F	<div></div> 0.2880	<div></div> 0.3370
P	<div></div> 0.8540	<div></div> 0.5400

1.0

0.0

<0.0