



wwPDB EM Validation Summary Report ⓘ

Dec 26, 2024 – 03:41 AM EST

PDB ID : 1OED
EMDB ID : EMD-1044
Title : STRUCTURE OF ACETYLCHOLINE RECEPTOR PORE FROM ELECTRON IMAGES
Authors : Miyazawa, A.; Fujiyoshi, Y.; Unwin, N.
Deposited on : 2003-03-24
Resolution : 4.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>37%</div> <div>5% 40% 10% 44%</div> </div>
1	D	227	<div> <div>30%</div> <div>5% 41% 9% 44%</div> </div>
2	B	250	<div> <div>32%</div> <div>• 34% 11% • 49%</div> </div>
3	C	260	<div> <div>34%</div> <div>6% 33% 8% • 51%</div> </div>
4	E	260	<div> <div>32%</div> <div>7% 30% 12% 51%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholine receptor subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	127	Total	C	N	O	S	0	0
			978	659	144	167	8		
1	D	127	Total	C	N	O	S	0	0
			978	659	144	167	8		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	230	GLY	VAL	variant	UNP P02711
A	291	ILE	VAL	variant	UNP P02711
A	318	ASP	ASN	variant	UNP P02711
D	230	GLY	VAL	variant	UNP P02711
D	291	ILE	VAL	variant	UNP P02711
D	318	ASP	ASN	variant	UNP P02711

- Molecule 2 is a protein called Acetylcholine receptor beta subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	127	Total	C	N	O	S	0	0
			995	677	147	165	6		

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	226	ILE	VAL	variant	UNP Q6S3I0
B	259	VAL	LEU	variant	UNP Q6S3I0
B	282	ARG	SER	variant	UNP Q6S3I0
B	442	VAL	ILE	variant	UNP Q6S3I0
B	444	PHE	ILE	variant	UNP Q6S3I0
B	445	VAL	THR	variant	UNP Q6S3I0
B	446	ILE	MET	variant	UNP Q6S3I0

- Molecule 3 is a protein called Acetylcholine receptor delta subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	127	Total	C	N	O	S	0	0
			982	660	151	163	8		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	244	SER	ALA	variant	UNP Q6S3H8
C	262	SER	CYS	variant	UNP Q6S3H8
C	303	ILE	VAL	variant	UNP Q6S3H8
C	343	ALA	VAL	variant	UNP Q6S3H8
C	346	SER	ILE	variant	UNP Q6S3H8
C	480	HIS	ARG	variant	UNP Q6S3H8

- Molecule 4 is a protein called Acetylcholine receptor gamma subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	128	Total	C	N	O	S	0	0
			993	676	149	161	7		


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	295	MET	LEU	variant	UNP Q6S3H9
E	296	LEU	VAL	variant	UNP Q6S3H9
E	299	MET	THR	variant	UNP Q6S3H9
E	327	GLY	GLU	variant	UNP Q6S3H9
E	336	GLN	HIS	variant	UNP Q6S3H9
E	461	ILE	LEU	variant	UNP Q6S3H9
E	472	PHE	LEU	variant	UNP Q6S3H9


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Acetylcholine receptor subunit alpha




• Molecule 1: Acetylcholine receptor subunit alpha





• Molecule 2: Acetylcholine receptor beta subunit







• Molecule 3: Acetylcholine receptor delta subunit



• Molecule 4: Acetylcholine receptor gamma subunit



THR	VAL	ASP	LEU	TYR	LYS	ASP	LEU	ALA	ASN	PHE	ALA	PRO	GLU	ILE	LYS	SER	PHE	GLY	VAL	ILE	MET	ILE	LYS	ALA	GLU	TYR	ILE	LEU	LYS	LYS	PRO	ARG	ASN	GLU	LEU	MET	PHE	GLY	GLU	ASN	GLU	LYS	ASN	TRP	VAL	LEU	ILE	G444	K445	V446	I447	D448	K449	A450	C451	F452	W453	I454	A455	L456	L457	L458
PRO	SER	GLU	THR	PRO	GLU	LYS	PRO	GLN	PRO	ARG	ARG	ARG	ARG	SER	SER	SER	PHE	GLY	ILE	MET	ILE	LYS	ALA	GLU	TYR	ILE	LEU	LYS	LYS	PRO	ARG	ASN	GLU	LEU	MET	PHE	GLY	GLU	ASN	GLU	LYS	ASN	TRP	VAL	LEU	ILE	LEU	LYS	ARG	VAL	ASN	LYS	MET	THR	SER	ASP	ILE	ASP	ILE	GLY	THR	

4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of segments used	Not provided	
Resolution determination method	FSC 0.5 CUT-OFF	Depositor
CTF correction method	Not provided	
Microscope	JEOL 3000SFF	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	Not provided	
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	36800	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	6.320	Depositor
Minimum map value	-2.439	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	2.18	Depositor
Map size (Å)	128, 128, 168	wwPDB
Map dimensions	128, 128, 55	wwPDB
Map angles (°)	90, 90, 90	wwPDB
Pixel spacing (Å)	1, 1, 1	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	1/996 (0.1%)	1.11	8/1357 (0.6%)
1	D	0.71	1/996 (0.1%)	1.11	7/1357 (0.5%)
2	B	0.99	4/1019 (0.4%)	1.16	7/1393 (0.5%)
3	C	0.75	0/1005	1.16	6/1367 (0.4%)
4	E	0.72	0/1014	1.17	5/1381 (0.4%)
All	All	0.78	6/5030 (0.1%)	1.14	33/6855 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	E	0	1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	271	PRO	N-CD	-18.04	1.22	1.47
2	B	445	VAL	CB-CG1	9.40	1.72	1.52
1	A	403	ALA	C-O	8.17	1.38	1.23
1	D	403	ALA	C-O	8.16	1.38	1.23
2	B	448	SER	CB-OG	-6.73	1.33	1.42

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	445	VAL	CG1-CB-CG2	-8.62	97.11	110.90
3	C	315	ARG	NE-CZ-NH2	7.83	124.21	120.30
3	C	455	ARG	NE-CZ-NH2	7.74	124.17	120.30
2	B	307	ARG	NE-CZ-NH2	7.61	124.10	120.30
4	E	310	ARG	NE-CZ-NH2	7.35	123.98	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	E	476	PRO	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	978	0	1059	510	0
1	D	978	0	1059	528	0
2	B	995	0	1064	503	0
3	C	982	0	1040	441	0
4	E	993	0	1077	513	0
All	All	4926	0	5299	2261	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 221.

The worst 5 of 2261 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:286:PHE:CE1	2:B:456:LEU:HB3	1.31	1.64
2:B:463:PRO:CG	2:B:464:PRO:HD3	1.26	1.59
3:C:285:VAL:CG1	3:C:286:PRO:HD2	1.33	1.52
1:A:296:ILE:HD11	4:E:239:LEU:CD1	1.38	1.51
2:B:463:PRO:HG2	2:B:464:PRO:CD	1.43	1.48

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/227 (54%)	118 (96%)	4 (3%)	1 (1%)	16	53
1	D	123/227 (54%)	117 (95%)	5 (4%)	1 (1%)	16	53
2	B	123/250 (49%)	108 (88%)	10 (8%)	5 (4%)	2	22
3	C	123/260 (47%)	114 (93%)	6 (5%)	3 (2%)	5	31
4	E	124/260 (48%)	114 (92%)	7 (6%)	3 (2%)	5	31
All	All	616/1224 (50%)	571 (93%)	32 (5%)	13 (2%)	8	33

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	218	LEU
2	B	282	ARG
2	B	249	MET
3	C	289	GLY
3	C	451	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/209 (56%)	98 (83%)	20 (17%)	1	11
1	D	118/209 (56%)	101 (86%)	17 (14%)	2	15
2	B	116/229 (51%)	89 (77%)	27 (23%)	0	4
3	C	111/233 (48%)	93 (84%)	18 (16%)	2	12
4	E	114/235 (48%)	90 (79%)	24 (21%)	1	5
All	All	577/1115 (52%)	471 (82%)	106 (18%)	3	9

5 of 106 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	457	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	269	SER
4	E	309	LEU
3	C	465	MET
1	D	225	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	300	HIS
1	D	408	HIS
4	E	473	ASN
1	D	435	GLN
3	C	267	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-1044. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.18. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 46 nm³; this corresponds to an approximate mass of 41 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-1044 and PDB model 1OED. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 2.18 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.18).

9.4 Atom inclusion [i](#)



At the recommended contour level, 40% of all backbone atoms, 34% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (2.18) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.3400	<div></div> 0.0660
A	<div></div> 0.3150	<div></div> 0.0560
B	<div></div> 0.3440	<div></div> 0.0520
C	<div></div> 0.2750	<div></div> 0.0660
D	<div></div> 0.4210	<div></div> 0.0960
E	<div></div> 0.3410	<div></div> 0.0570

