



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2025 – 08:23 PM EDT

PDB ID : 2OHQ / pdb_00002ohq
Title : X-ray crystal structure of beta secretase complexed with compound 4
Authors : Patel, S.
Deposited on : 2007-01-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	<div> <div>27%</div> <div>73%</div> <div>19%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	405	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3300 atoms, of which 21 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	377	2987	1913	499	561	14	0	3	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	LYS	ARG	engineered mutation	UNP P56817
A	-4	LYS	ARG	engineered mutation	UNP P56817

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total I 2 2	0	0

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is 6-[2-(3'-METHOXYBIPHENYL-3-YL)ETHYL]PYRIDIN-2-AMINE (CCD ID: 7IP) (formula: C₂₀H₂₀N₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	0	0
			44	20	21	2	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	257	Total	O	0	0
			257	257		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-secretase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.69Å 102.69Å 169.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.60 – 2.10 47.60 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.9 (47.60-2.10) 98.8 (47.60-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.97 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019g	Depositor
R, R_{free}	0.218 , 0.276 0.227 , 0.279	Depositor DCC
R_{free} test set	1568 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3300	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, GOL, 7IP, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/3071	0.86	0/4172

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	233	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2987	0	2915	68	0
2	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	6	0	0
4	A	23	21	20	4	0
5	A	6	0	8	4	0
6	A	257	0	0	5	2
All	All	3279	21	2949	70	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:ILE:HD12	1:A:213:LEU:HD11	1.45	0.96
1:A:118[A]:ILE:HD12	4:A:404:7IP:H142	1.56	0.86
1:A:37:ASN:HD21	1:A:128[B]:ARG:H	1.30	0.78
4:A:404:7IP:H13	5:A:405:GOL:H11	1.66	0.77
1:A:37:ASN:HD21	1:A:128[A]:ARG:H	1.31	0.75
1:A:96:ARG:H	1:A:143:GLN:HE22	1.36	0.71
1:A:215:MET:CE	1:A:243:ALA:HB3	2.21	0.70
1:A:215:MET:HE1	1:A:243:ALA:HB3	1.73	0.69
1:A:73:GLN:OE1	1:A:107:LYS:NZ	2.25	0.67
1:A:304:GLN:HE21	1:A:304:GLN:N	1.92	0.67
1:A:215:MET:HE2	1:A:243:ALA:CB	2.27	0.65
4:A:404:7IP:H13	5:A:405:GOL:C1	2.26	0.64
1:A:259:ASP:O	1:A:263:LEU:HD22	2.01	0.60
1:A:55:GLN:HE21	1:A:55:GLN:H	1.50	0.58
1:A:118[A]:ILE:HD12	4:A:404:7IP:C14	2.32	0.58
1:A:303:GLN:C	1:A:304:GLN:HE21	2.12	0.58
1:A:254:THR:CB	1:A:279:ILE:HD12	2.34	0.58
1:A:20:VAL:HG11	1:A:99:ILE:HD13	1.86	0.58
1:A:215:MET:CE	1:A:243:ALA:CB	2.83	0.57
1:A:67:VAL:HG12	1:A:129:PRO:HD3	1.86	0.56
1:A:170:VAL:HG21	5:A:405:GOL:O3	2.05	0.56
1:A:224:LYS:HE3	1:A:329:THR:O	2.07	0.55
1:A:184:TYR:O	6:A:407:HOH:O	2.18	0.55
1:A:179:ILE:HG23	1:A:342:TYR:HE2	1.72	0.54
1:A:301:LEU:HB3	1:A:302:PRO:HD2	1.89	0.54
1:A:300:ILE:HG21	1:A:337:ILE:HD13	1.88	0.54
1:A:304:GLN:N	1:A:304:GLN:NE2	2.55	0.54
1:A:323:ALA:HB1	1:A:336:VAL:HG21	1.90	0.53
1:A:49:HIS:HD2	6:A:565:HOH:O	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:PRO:O	1:A:306:LEU:HB2	2.10	0.52
1:A:181:HIS:HB3	6:A:549:HOH:O	2.09	0.52
1:A:321:LYS:C	6:A:658:HOH:O	2.53	0.51
1:A:333:MET:HE2	1:A:337:ILE:HG21	1.93	0.51
1:A:30:LEU:HB3	1:A:118[B]:ILE:HG22	1.92	0.51
1:A:320:TYR:HB3	6:A:658:HOH:O	2.10	0.50
1:A:304:GLN:HE21	1:A:304:GLN:CA	2.24	0.50
1:A:67:VAL:CG1	1:A:80:LEU:HD11	2.42	0.50
1:A:215:MET:O	1:A:216:ASP:C	2.54	0.49
1:A:154:LEU:O	1:A:339:GLU:HA	2.13	0.48
1:A:55:GLN:H	1:A:55:GLN:NE2	2.12	0.48
1:A:271:GLN:O	1:A:274:THR:OG1	2.22	0.47
1:A:50:ARG:O	1:A:116:GLU:HG2	2.15	0.47
1:A:254:THR:HG21	1:A:279:ILE:HD12	1.97	0.47
1:A:312:VAL:O	1:A:312:VAL:HG23	2.15	0.46
1:A:8:GLY:C	1:A:170:VAL:HG22	2.41	0.46
1:A:207:GLU:HA	1:A:212:ASP:HA	1.98	0.45
1:A:272:ALA:O	1:A:274:THR:HG23	2.16	0.45
1:A:125:GLU:OE2	1:A:195:ARG:NH1	2.49	0.44
1:A:271:GLN:HB2	1:A:274:THR:HG21	1.97	0.44
1:A:361:VAL:O	1:A:361:VAL:HG23	2.17	0.44
1:A:199:TYR:HB3	1:A:352:ILE:HD11	2.00	0.43
1:A:156:GLY:O	1:A:157:ALA:HB3	2.19	0.43
1:A:252:SER:HA	1:A:279:ILE:HG22	2.01	0.43
1:A:234:LEU:O	1:A:324:ILE:HA	2.19	0.43
1:A:20:VAL:CG1	1:A:99:ILE:CD1	2.97	0.42
1:A:11:GLY:H	5:A:405:GOL:H2	1.83	0.42
1:A:20:VAL:CG1	1:A:99:ILE:HD13	2.50	0.42
1:A:254:THR:HB	1:A:279:ILE:HG21	2.01	0.42
1:A:254:THR:HB	1:A:279:ILE:HD12	2.01	0.42
1:A:50:ARG:HH22	1:A:114:ASN:ND2	2.18	0.42
1:A:232:THR:O	1:A:336:VAL:HG23	2.20	0.41
1:A:302:PRO:O	1:A:306:LEU:N	2.44	0.41
1:A:156:GLY:HA2	1:A:170:VAL:CG1	2.51	0.41
1:A:241:PHE:CZ	1:A:245:VAL:HG21	2.55	0.41
1:A:236:LEU:O	1:A:326:GLN:HA	2.19	0.41
1:A:305:TYR:CD1	1:A:306:LEU:HD13	2.56	0.41
1:A:67:VAL:CG1	1:A:129:PRO:HD3	2.51	0.41
1:A:110:ILE:HB	1:A:113:SER:HB3	2.03	0.40
1:A:174:MET:HE3	1:A:176:ILE:HD11	2.03	0.40
1:A:303:GLN:HG3	1:A:361:VAL:HG23	2.02	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:413:HOH:O	6:A:520:HOH:O[12_555]	1.89	0.31
6:A:443:HOH:O	6:A:513:HOH:O[6_655]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/402 (94%)	360 (96%)	15 (4%)	1 (0%)	37	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	73	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/343 (94%)	310 (96%)	14 (4%)	25	25

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	55	GLN
1	A	72	THR
1	A	75	LYS
1	A	149	LEU
1	A	209	ASN
1	A	212	ASP
1	A	234	LEU
1	A	263	LEU
1	A	304	GLN
1	A	306	LEU
1	A	315	SER
1	A	326	GLN
1	A	328	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	37	ASN
1	A	49	HIS
1	A	55	GLN
1	A	89	HIS
1	A	98	ASN
1	A	114	ASN
1	A	143	GLN
1	A	209	ASN
1	A	233	ASN
1	A	304	GLN
1	A	362	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	405	-	5,5,5	0.69	0	5,5,5	0.80	0
4	7IP	A	404	-	24,25,25	0.70	0	29,33,33	1.31	3 (10%)
3	DMS	A	403	-	3,3,3	2.64	1 (33%)	3,3,3	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	405	-	-	2/4/4/4	-
4	7IP	A	404	-	-	0/7/11/11	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	DMS	O-S	4.43	1.79	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	404	7IP	C20-N22-C16	4.40	121.36	118.07
4	A	404	7IP	O2-C3-C8	-3.02	115.91	124.40
4	A	404	7IP	C17-C16-N22	-2.16	119.67	122.40

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	405	GOL	O1-C1-C2-C3
5	A	405	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	405	GOL	4	0
4	A	404	7IP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/402 (93%)	1.12	109 (28%) 1 1	15, 37, 81, 97	5 (1%)

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	TRP	8.2
1	A	312	VAL	7.5
1	A	319	CYS	7.4
1	A	313	ALA	7.2
1	A	365	PHE	6.9
1	A	254	THR	6.3
1	A	276	PRO	6.1
1	A	314	THR	5.8
1	A	279	ILE	5.7
1	A	244	ALA	5.5
1	A	367	THR	5.3
1	A	170	VAL	4.7
1	A	280	PHE	4.7
1	A	245	VAL	4.6
1	A	256	LYS	4.6
1	A	260	GLY	4.5
1	A	264	GLY	4.5
1	A	315	SER	4.4
1	A	251	ALA	4.4
1	A	282	VAL	4.3
1	A	267	LEU	4.3
1	A	169	SER	4.2
1	A	275	THR	4.2
1	A	263	LEU	4.2
1	A	241	PHE	4.2
1	A	262	TRP	4.1
1	A	168	ALA	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	363	ASP	4.0
1	A	272	ALA	3.9
1	A	308	PRO	3.9
1	A	278	ASN	3.8
1	A	326	GLN	3.8
1	A	274	THR	3.7
1	A	281	PRO	3.7
1	A	384	TYR	3.7
1	A	364	GLU	3.7
1	A	257	PHE	3.6
1	A	311	ASP	3.5
1	A	157	ALA	3.4
1	A	305	TYR	3.4
1	A	238	LYS	3.4
1	A	322	PHE	3.3
1	A	258	PRO	3.3
1	A	222	TYR	3.3
1	A	301	LEU	3.3
1	A	156	GLY	3.2
1	A	215	MET	3.2
1	A	309	VAL	3.2
1	A	242	GLU	3.2
1	A	268	VAL	3.1
1	A	271	GLN	3.1
1	A	307	ARG	3.1
1	A	73	GLN	3.1
1	A	318	ASP	3.1
1	A	361	VAL	3.1
1	A	-1	PHE	3.1
1	A	208	ILE	3.0
1	A	362	HIS	3.0
1	A	302	PRO	3.0
1	A	306	LEU	2.9
1	A	243	ALA	2.9
1	A	235	ARG	2.9
1	A	213	LEU	2.9
1	A	320	TYR	2.9
1	A	250	ALA	2.9
1	A	72	THR	2.9
1	A	261	PHE	2.8
1	A	329	THR	2.8
1	A	253	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	269	CYS	2.8
1	A	310	GLU	2.8
1	A	211	GLN	2.7
1	A	380	GLU	2.7
1	A	323	ALA	2.7
1	A	255	GLU	2.7
1	A	209	ASN	2.7
1	A	128[A]	ARG	2.7
1	A	239	LYS	2.6
1	A	10	SER	2.6
1	A	317	ASP	2.6
1	A	325	SER	2.6
1	A	181	HIS	2.6
1	A	65	LYS	2.6
1	A	212	ASP	2.6
1	A	236	LEU	2.6
1	A	210	GLY	2.5
1	A	71	TYR	2.5
1	A	74	GLY	2.5
1	A	68	TYR	2.5
1	A	303	GLN	2.5
1	A	321	LYS	2.5
1	A	92	ASN	2.5
1	A	383	GLY	2.5
1	A	259	ASP	2.5
1	A	270	TRP	2.5
1	A	328	SER	2.4
1	A	324	ILE	2.4
1	A	9	LYS	2.4
1	A	7	ARG	2.4
1	A	366	ARG	2.3
1	A	0	VAL	2.2
1	A	273	GLY	2.2
1	A	69	VAL	2.2
1	A	283	ILE	2.2
1	A	237	PRO	2.2
1	A	266	GLN	2.1
1	A	214	LYS	2.1
1	A	385	ASN	2.0
1	A	265	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	A	405	6/6	0.81	0.21	50,52,56,60	0
4	7IP	A	404	23/23	0.83	0.15	28,55,60,61	0
3	DMS	A	403	4/4	0.96	0.13	54,55,57,57	0
2	IOD	A	402	1/1	0.98	0.05	47,47,47,47	1
2	IOD	A	401	1/1	1.00	0.03	28,28,28,28	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7IP A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.