



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 28, 2024 – 06:51 pm GMT

PDB ID : 5OJ2
Title : Crystal structure of the chicken MDGA1 ectodomain
Authors : Elegheert, J.; Clayton, A.J.; Aricescu, A.R.
Deposited on : 2017-07-20
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1370 (3.20-3.20)
Clashscore	180529	1497 (3.20-3.20)
Ramachandran outliers	177936	1479 (3.20-3.20)
Sidechain outliers	177891	1478 (3.20-3.20)
RSRZ outliers	164620	1371 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	910	 2% 67% 8% 24%
1	B	910	 % 68% 8% 24%
2	C	3	 100%
3	D	4	 50% 50%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MAM domain-containing glycosylphosphatidylinositol anchor protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	691	Total	C	N	O	S	Se	0	0	0
			5384	3396	930	1031	16	11			
1	B	692	Total	C	N	O	S	Se	0	0	0
			5404	3412	935	1030	16	11			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	120	LYS	ARG	conflict	UNP Q0WYX8
A	920	GLY	-	expression tag	UNP Q0WYX8
A	921	THR	-	expression tag	UNP Q0WYX8
A	922	LYS	-	expression tag	UNP Q0WYX8
A	923	HIS	-	expression tag	UNP Q0WYX8
A	924	HIS	-	expression tag	UNP Q0WYX8
A	925	HIS	-	expression tag	UNP Q0WYX8
A	926	HIS	-	expression tag	UNP Q0WYX8
A	927	HIS	-	expression tag	UNP Q0WYX8
A	928	HIS	-	expression tag	UNP Q0WYX8
B	120	LYS	ARG	engineered mutation	UNP Q0WYX8
B	920	GLY	-	expression tag	UNP Q0WYX8
B	921	THR	-	expression tag	UNP Q0WYX8
B	922	LYS	-	expression tag	UNP Q0WYX8
B	923	HIS	-	expression tag	UNP Q0WYX8
B	924	HIS	-	expression tag	UNP Q0WYX8
B	925	HIS	-	expression tag	UNP Q0WYX8
B	926	HIS	-	expression tag	UNP Q0WYX8
B	927	HIS	-	expression tag	UNP Q0WYX8
B	928	HIS	-	expression tag	UNP Q0WYX8

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

GLU
LYS
ILE
CYS
GLY
PHE
VAL
GLN
ASP
LYS
NSE
ALA
ASN
PHE
ASP
TRP
THR
ARG
GLN
ASN
ALA
ALA
THR
THR
GLN
ASN
PRO
LYS
ARG
THR
VAL
PRO
THR
THR
ASP
ILE
SER
GLY
THR
THR
GLU
TYR
TYR

ALA
ARG
LEU
ILE
SER
PRO
LEU
TYR
ASN
ILE
THR
ALA
LYS
TYR
CYS
TRP
VAL
SER
PHE
GLN
TYR
TYR
HIS
NSE
TYR
GLN
GLY
LYS
HIS
ILE
GLY
SER
VAL
LEU
ASN
GLY
LEU
VAL
ARG
THR
VAL
ARG
ASN
LYS
ARG
ALA
PRO
ILE
ASP
THR
GLN
VAL
TRP
SER
LEU
SER
GLY
ASN
ARG
PRO
GLY
ASN
VAL
NSE
THR
GLN
LYS

ALA
HIS
VAL
PRO
ILE
ASN
PRO
PRO
GLY
PRO
PHE
GLN
ILE
ILE
PHE
GLU
GLY
VAL
ARG
GLY
THR
SER
TYR
GLU
GLY
ASP
ILE
ALA
ILE
ILE
ASP
VAL
THR
LEU
LYS
LYS
GLY
ASP
CYS
PRO
ARG
LYS
PRO
ILE
GLY
PRO
ASN
LYS
GLY
THR
LYS
HIS
HIS
HIS
HIS
HIS

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

100%

MAG1
MAG2
BMA3

- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

50%

50%

MAG1
MAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.18Å 109.04Å 208.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.65 – 3.20 96.65 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (96.65-3.20) 99.9 (96.65-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 3.19Å)	Xtriage
Refinement program	PHENIX (dev_2044)	Depositor
R, R_{free}	0.214 , 0.251 0.228 , 0.267	Depositor DCC
R_{free} test set	2030 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	94.6	Xtriage
Anisotropy	0.196	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 66.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11087	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/5484	0.45	0/7452
1	B	0.25	0/5507	0.45	0/7481
All	All	0.25	0/10991	0.45	0/14933

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5384	0	5351	50	0
1	B	5404	0	5391	42	0
2	C	39	0	34	0	0
3	D	50	0	43	0	0
4	A	112	0	104	0	0
4	B	98	0	91	0	0
All	All	11087	0	11014	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:680:ARG:NH2	1:B:584:PRO:O	2.13	0.80
1:B:572:SER:N	1:B:587:GLU:OE2	2.19	0.73
1:B:19:GLN:OE1	1:B:19:GLN:N	2.31	0.63
1:B:546:ASP:OD1	1:B:548:ARG:NE	2.33	0.61
1:A:174:GLU:N	1:A:174:GLU:OE1	2.35	0.59

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	681/910 (75%)	627 (92%)	53 (8%)	1 (0%)	48	80
1	B	682/910 (75%)	632 (93%)	50 (7%)	0	100	100
All	All	1363/1820 (75%)	1259 (92%)	103 (8%)	1 (0%)	48	80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	391	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	608/795 (76%)	596 (98%)	12 (2%)	50	75
1	B	612/795 (77%)	602 (98%)	10 (2%)	58	79
All	All	1220/1590 (77%)	1198 (98%)	22 (2%)	54	77

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	526	ARG
1	B	648	LYS
1	B	645	LYS
1	B	675	ARG
1	A	553	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	676	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.22	0	17,19,21	0.50	0
2	NAG	C	2	2	14,14,15	0.42	0	17,19,21	0.47	0
2	BMA	C	3	2	11,11,12	0.55	0	15,15,17	0.83	0
3	NAG	D	1	1,3	14,14,15	0.18	0	17,19,21	0.56	0
3	NAG	D	2	3	14,14,15	0.24	0	17,19,21	0.43	0
3	BMA	D	3	3	11,11,12	0.74	0	15,15,17	1.52	4 (26%)
3	MAN	D	4	3	11,11,12	1.04	1 (9%)	15,15,17	0.84	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	4	MAN	O5-C1	-2.68	1.39	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3	BMA	C1-C2-C3	-2.91	106.09	109.67
3	D	3	BMA	C3-C4-C5	-2.54	105.70	110.24
3	D	3	BMA	O3-C3-C2	2.53	114.83	109.99
3	D	4	MAN	O2-C2-C3	-2.27	105.59	110.14
3	D	3	BMA	O3-C3-C4	2.22	115.49	110.35

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	4	MAN	O5-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
3	D	4	MAN	C4-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1011	1	14,14,15	0.20	0	17,19,21	0.48	0
4	NAG	B	1001	1	14,14,15	0.27	0	17,19,21	0.44	0
4	NAG	A	1008	1	14,14,15	0.37	0	17,19,21	0.58	0
4	NAG	B	1009	1	14,14,15	0.27	0	17,19,21	0.41	0
4	NAG	A	1002	1	14,14,15	0.18	0	17,19,21	0.54	0
4	NAG	A	1007	1	14,14,15	0.30	0	17,19,21	0.46	0
4	NAG	A	1010	1	14,14,15	0.18	0	17,19,21	0.59	0
4	NAG	B	1011	1	14,14,15	0.39	0	17,19,21	0.43	0
4	NAG	B	1003	1	14,14,15	0.20	0	17,19,21	0.39	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1003	1	14,14,15	0.20	0	17,19,21	0.38	0
4	NAG	A	1001	1	14,14,15	0.19	0	17,19,21	0.52	0
4	NAG	B	1002	1	14,14,15	0.26	0	17,19,21	0.38	0
4	NAG	B	1010	1	14,14,15	0.21	0	17,19,21	0.62	1 (5%)
4	NAG	A	1009	1	14,14,15	0.20	0	17,19,21	0.55	0
4	NAG	B	1008	1	14,14,15	0.51	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1011	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1001	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1008	1	-	4/6/23/26	0/1/1/1
4	NAG	B	1009	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1002	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1007	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1010	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1011	1	-	1/6/23/26	0/1/1/1
4	NAG	B	1003	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1003	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1001	1	-	1/6/23/26	0/1/1/1
4	NAG	B	1002	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1010	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1009	1	-	2/6/23/26	0/1/1/1
4	NAG	B	1008	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1010	NAG	C1-O5-C5	2.13	115.08	112.19

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1009	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	1009	NAG	O5-C5-C6-O6
4	B	1008	NAG	O5-C5-C6-O6
4	A	1009	NAG	C4-C5-C6-O6
4	B	1008	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	680/910 (74%)	-0.00	16 (2%) 59 44	47, 100, 156, 304	0
1	B	681/910 (74%)	-0.08	8 (1%) 76 61	48, 102, 151, 215	0
All	All	1361/1820 (74%)	-0.04	24 (1%) 67 52	47, 101, 154, 304	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	VAL	4.0
1	A	176	LEU	3.4
1	B	86	THR	3.2
1	A	638	PRO	3.0
1	A	383	ARG	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	D	4	11/12	0.31	0.18	216,223,228,237	0
2	BMA	C	3	11/12	0.58	0.13	163,172,181,186	0
3	BMA	D	3	11/12	0.62	0.13	173,181,193,205	0
3	NAG	D	1	14/15	0.85	0.15	83,101,111,112	0
3	NAG	D	2	14/15	0.87	0.19	128,135,150,160	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	C	2	14/15	0.88	0.14	124,129,138,152	0
2	NAG	C	1	14/15	0.89	0.13	92,109,118,120	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	B	1011	14/15	0.39	0.13	166,192,232,253	0
4	NAG	A	1007	14/15	0.64	0.13	158,178,215,222	0
4	NAG	A	1008	14/15	0.67	0.12	142,145,148,150	0
4	NAG	A	1002	14/15	0.67	0.11	136,148,152,157	0
4	NAG	B	1009	14/15	0.70	0.10	130,142,164,189	0
4	NAG	A	1010	14/15	0.70	0.15	141,165,172,173	0
4	NAG	B	1002	14/15	0.73	0.13	147,158,163,166	0
4	NAG	B	1008	14/15	0.73	0.14	170,175,212,230	0
4	NAG	A	1009	14/15	0.81	0.08	142,147,162,169	0
4	NAG	B	1003	14/15	0.84	0.12	108,116,119,120	0
4	NAG	B	1001	14/15	0.84	0.13	147,166,182,188	0
4	NAG	A	1001	14/15	0.85	0.10	108,114,117,120	0
4	NAG	B	1010	14/15	0.91	0.12	101,108,123,126	0
4	NAG	A	1003	14/15	0.92	0.08	109,114,119,124	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	1011	14/15	0.94	0.08	74,78,84,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.