



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 09:50 AM EDT

PDB ID : 1P1H  
Title : Crystal structure of the 1L-myo-inositol/NAD<sup>+</sup> complex  
Authors : Jin, X.; Geiger, J.H.  
Deposited on : 2003-04-12  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	533	
1	B	533	
1	C	533	
1	D	533	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol-3-phosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	506	Total	C	N	O	S	0	0	0
			3988	2538	669	765	16			
1	B	491	Total	C	N	O	S	0	0	0
			3876	2467	652	741	16			
1	C	517	Total	C	N	O	S	0	0	0
			4081	2596	686	783	16			
1	D	487	Total	C	N	O	S	0	0	0
			3839	2439	643	743	14			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ARG	SEE REMARK 999	UNP P11986
A	14	VAL	LEU	SEE REMARK 999	UNP P11986
A	?	-	PHE	SEE REMARK 999	UNP P11986
A	60	LEU	GLU	SEE REMARK 999	UNP P11986
A	?	-	ALA	SEE REMARK 999	UNP P11986
A	98	GLU	LYS	SEE REMARK 999	UNP P11986
A	140	ASN	LYS	SEE REMARK 999	UNP P11986
A	141	ASP	HIS	SEE REMARK 999	UNP P11986
A	201	ASN	GLN	SEE REMARK 999	UNP P11986
A	444	PRO	ALA	SEE REMARK 999	UNP P11986
B	?	-	ARG	SEE REMARK 999	UNP P11986
B	14	VAL	LEU	SEE REMARK 999	UNP P11986
B	?	-	PHE	SEE REMARK 999	UNP P11986
B	60	LEU	GLU	SEE REMARK 999	UNP P11986
B	?	-	ALA	SEE REMARK 999	UNP P11986
B	98	GLU	LYS	SEE REMARK 999	UNP P11986
B	140	ASN	LYS	SEE REMARK 999	UNP P11986
B	141	ASP	HIS	SEE REMARK 999	UNP P11986
B	201	ASN	GLN	SEE REMARK 999	UNP P11986
B	444	PRO	ALA	SEE REMARK 999	UNP P11986
C	?	-	ARG	SEE REMARK 999	UNP P11986

*Continued on next page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	14	VAL	LEU	SEE REMARK 999	UNP P11986
C	?	-	PHE	SEE REMARK 999	UNP P11986
C	60	LEU	GLU	SEE REMARK 999	UNP P11986
C	?	-	ALA	SEE REMARK 999	UNP P11986
C	98	GLU	LYS	SEE REMARK 999	UNP P11986
C	140	ASN	LYS	SEE REMARK 999	UNP P11986
C	141	ASP	HIS	SEE REMARK 999	UNP P11986
C	201	ASN	GLN	SEE REMARK 999	UNP P11986
C	444	PRO	ALA	SEE REMARK 999	UNP P11986
D	?	-	ARG	SEE REMARK 999	UNP P11986
D	14	VAL	LEU	SEE REMARK 999	UNP P11986
D	?	-	PHE	SEE REMARK 999	UNP P11986
D	60	LEU	GLU	SEE REMARK 999	UNP P11986
D	?	-	ALA	SEE REMARK 999	UNP P11986
D	98	GLU	LYS	SEE REMARK 999	UNP P11986
D	140	ASN	LYS	SEE REMARK 999	UNP P11986
D	141	ASP	HIS	SEE REMARK 999	UNP P11986
D	201	ASN	GLN	SEE REMARK 999	UNP P11986
D	444	PRO	ALA	SEE REMARK 999	UNP P11986

- # NAD

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	425	Total	O	0	0
			425	425		
3	B	354	Total	O	0	0
			354	354		
3	C	383	Total	O	0	0
			383	383		
3	D	367	Total	O	0	0
			367	367		


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Inositol-3-phosphate synthase




#### • Molecule 1: Inositol-3-phosphate synthase







• Molecule 1: Inositol-3-phosphate synthase



• Molecule 1: Inositol-3-phosphate synthase







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.79Å 185.58Å 94.08Å 90.00° 114.77° 90.00°	Depositor
Resolution (Å)	10.00 – 1.95 26.35 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.95) 96.3 (26.35-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.41 (at 1.89Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.208 , 0.249 0.232 , 0.230	Depositor DCC
$R_{free}$ test set	10775 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.1	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 89.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for l,k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	17489	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.35	0/4066	0.61	0/5513
1	B	0.35	0/3951	0.60	0/5354
1	C	0.36	0/4160	0.60	1/5637 (0.0%)
1	D	0.37	0/3913	0.62	1/5308 (0.0%)
All	All	0.36	0/16090	0.61	2/21812 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	318	GLY	N-CA-C	5.40	126.59	113.10
1	C	516	LEU	CA-CB-CG	5.15	127.15	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3988	0	3989	213	0
1	B	3876	0	3879	220	0
1	C	4081	0	4088	258	0
1	D	3839	0	3816	229	0
2	A	44	0	26	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	44	0	26	3	0
2	C	44	0	26	5	0
2	D	44	0	26	1	0
3	A	425	0	0	27	0
3	B	354	0	0	20	0
3	C	383	0	0	25	0
3	D	367	0	0	18	0
All	All	17489	0	15876	856	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 856 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:494:ARG:HB3	1:C:495:PRO:HD2	1.34	1.08
1:C:494:ARG:HH22	1:D:533:LEU:HB3	0.94	1.06
1:A:352:LEU:HD21	1:A:357:GLY:HA3	1.37	1.06
1:C:69:MET:HE1	1:C:227:ILE:HG12	1.42	1.01
1:D:191:ILE:HG22	1:D:192:ALA:H	1.25	1.01

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	502/533 (94%)	465 (93%)	28 (6%)	9 (2%)	<b>8</b> <b>2</b>
1	B	485/533 (91%)	445 (92%)	32 (7%)	8 (2%)	<b>9</b> <b>2</b>
1	C	513/533 (96%)	474 (92%)	27 (5%)	12 (2%)	<b>6</b> <b>1</b>
1	D	481/533 (90%)	445 (92%)	28 (6%)	8 (2%)	<b>9</b> <b>2</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1981/2132 (93%)	1829 (92%)	115 (6%)	37 (2%)	8 2

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	ALA
1	A	211	VAL
1	A	319	ASP
1	A	495	PRO
1	B	192	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	446/471 (95%)	418 (94%)	28 (6%)	18 7
1	B	434/471 (92%)	412 (95%)	22 (5%)	24 11
1	C	458/471 (97%)	435 (95%)	23 (5%)	24 11
1	D	429/471 (91%)	403 (94%)	26 (6%)	18 7
All	All	1767/1884 (94%)	1668 (94%)	99 (6%)	21 9

5 of 99 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	251	VAL
1	C	516	LEU
1	C	298	GLN
1	C	397	ASP
1	D	34	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 94 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	263	ASN
1	D	88	ASN
1	C	298	GLN
1	C	512	ASN
1	D	169	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAD	B	660	-	42,48,48	2.51	10 (23%)	50,73,73	1.26	6 (12%)
2	NAD	C	670	-	42,48,48	2.19	8 (19%)	50,73,73	1.27	6 (12%)
2	NAD	A	650	-	42,48,48	2.35	8 (19%)	50,73,73	1.26	5 (10%)
2	NAD	D	680	-	42,48,48	2.33	10 (23%)	50,73,73	1.27	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	660	-	-	1/26/62/62	0/5/5/5
2	NAD	C	670	-	-	3/26/62/62	0/5/5/5
2	NAD	A	650	-	-	2/26/62/62	0/5/5/5
2	NAD	D	680	-	-	3/26/62/62	0/5/5/5

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	660	NAD	C2N-N1N	8.85	1.44	1.35
2	A	650	NAD	C2N-N1N	7.78	1.43	1.35
2	B	660	NAD	C4N-C3N	6.97	1.50	1.39
2	C	670	NAD	C2N-N1N	6.91	1.42	1.35
2	A	650	NAD	C4N-C3N	6.86	1.49	1.39

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	660	NAD	C3N-C7N-N7N	3.73	122.34	117.74
2	A	650	NAD	C3N-C7N-N7N	3.45	121.99	117.74
2	C	670	NAD	O7N-C7N-N7N	-3.44	117.64	122.62
2	A	650	NAD	O2A-PA-O3	3.32	116.25	107.27
2	D	680	NAD	O2A-PA-O3	3.26	116.07	107.27

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	680	NAD	C5D-O5D-PN-O1N
2	D	680	NAD	PN-O3-PA-O1A
2	C	670	NAD	PN-O3-PA-O1A
2	C	670	NAD	PN-O3-PA-O2A
2	D	680	NAD	PN-O3-PA-O2A


There are no ring outliers.

4 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	660	NAD	3	0
2	C	670	NAD	5	0
2	A	650	NAD	3	0
2	D	680	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	506/533 (94%)	0.65	72 (14%) 2 4	24, 45, 85, 90	0
1	B	491/533 (92%)	0.71	77 (15%) 2 3	23, 45, 88, 90	0
1	C	517/533 (96%)	0.65	68 (13%) 3 5	22, 45, 86, 90	0
1	D	487/533 (91%)	0.61	66 (13%) 3 4	21, 41, 88, 90	0
All	All	2001/2132 (93%)	0.65	283 (14%) 2 4	21, 44, 87, 90	0

The worst 5 of 283 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	192	ALA	8.9
1	B	193	ALA	8.6
1	C	41	ALA	8.1
1	D	353	GLY	8.0
1	B	41	ALA	8.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAD	D	680	44/44	0.95	0.13	26,35,70,75	0
2	NAD	A	650	44/44	0.96	0.14	32,40,65,68	0
2	NAD	C	670	44/44	0.97	0.14	29,34,65,70	0
2	NAD	B	660	44/44	0.97	0.12	29,40,56,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around NAD A 650:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAD C 670:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.