



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 15, 2024 – 04:07 AM EST

PDB ID : 1PJX  
Title : 0.85 ANGSTROM STRUCTURE OF SQUID GANGLION DFPASE  
Authors : Koepke, J.; Rueterjans, H.; Luecke, C.; Fritzsche, G.  
Deposited on : 2003-06-04  
Resolution : 0.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 0.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	1407 (1.00-0.72)
Ramachandran outliers	177936	1328 (1.00-0.72)
Sidechain outliers	177891	1329 (1.00-0.72)
RSRZ outliers	164620	1266 (1.00-0.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	<div> <div>5%</div> <div>74%</div> <div>22%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	PEG	A	461	-	X	X	-
3	ME2	A	471	-	-	X	-
4	MES	A	411	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MES	A	412	-	-	X	-
5	EDO	A	423	-	-	X	-
5	EDO	A	425	-	-	X	-
5	EDO	A	427	-	-	X	-
5	EDO	A	428	-	-	X	-
6	PGE	A	433	-	-	X	-
6	PGE	A	434	-	X	X	-
7	DXE	A	441	-	-	X	-
7	DXE	A	443	-	-	X	-
8	MXE	A	451	-	X	X	-
8	MXE	A	452	-	-	X	-
9	GOL	A	401	-	X	-	-

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 3295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DIISOPROPYLFLUOROPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	46	0
			2657	1688	456	495	18			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is 1-ETHOXY-2-(2-METHOXYETHOXY)ETHANE (three-letter code: ME2) (formula: C<sub>7</sub>H<sub>16</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	7	3		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 10 6 4	0	0
6	A	1	Total C O 10 6 4	0	0

- Molecule 7 is 1,2-DIMETHOXYETHANE (three-letter code: DXE) (formula:  $C_4H_{10}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	4	2		
7	A	1	Total	C	O	0	0
			6	4	2		
7	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 8 is 2-METHOXYETHANOL (three-letter code: MXE) (formula:  $C_3H_8O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			5	3	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 10 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			7	4	3		
10	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	479	Total	O	0	17
			496	496		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: DIISOPROPYLFLUOROPHOSPHATASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.11Å 81.85Å 86.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 0.85 10.00 – 0.85	Depositor EDS
% Data completeness (in resolution range)	90.8 (10.00-0.85) 87.0 (10.00-0.85)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.45 (at 0.85Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.111 , 0.128 0.158 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	6.3	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 76.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3295	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PEG, MES, ME2, EDO, PGE, DXE, MXE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.53	33/2949 (1.1%)	1.67	55/3975 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	264	ARG	NE-CZ	-23.31	1.02	1.33
1	A	278	GLN	CD-OE1	21.50	1.71	1.24
1	A	39	GLU	CD-OE1	20.33	1.48	1.25
1	A	150	GLU	CD-OE2	15.97	1.43	1.25
1	A	112	GLU	CD-OE2	-15.57	1.08	1.25

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	ARG	NE-CZ-NH1	25.43	133.02	120.30
1	A	43	LYS	CG-CD-CE	19.11	169.22	111.90
1	A	186[A]	ARG	NE-CZ-NH2	17.94	129.27	120.30
1	A	186[B]	ARG	NE-CZ-NH2	17.94	129.27	120.30
1	A	115[A]	ARG	CD-NE-CZ	16.26	146.36	123.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	188	TYR	Sidechain
1	A	278	GLN	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2657	0	2566	139	0
2	A	2	0	0	0	0
3	A	10	0	16	11	0
4	A	24	0	26	17	0
5	A	32	0	48	33	0
6	A	20	0	28	16	0
7	A	18	0	30	28	0
8	A	10	0	16	37	0
9	A	12	0	9	0	0
10	A	14	0	20	7	0
11	A	496	0	0	57	0
All	All	3295	0	2759	191	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 191 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:LYS:CD	1:A:109:LYS:CE	1.83	1.57
1:A:109:LYS:CE	1:A:109:LYS:NZ	1.80	1.41
1:A:43:LYS:NZ	4:A:411:MES:H31	1.34	1.37
1:A:278:GLN:OE1	1:A:278:GLN:CD	1.71	1.28
1:A:264:ARG:HG2	11:A:1047:HOH:O	1.19	1.26

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/314 (115%)	343 (95%)	13 (4%)	4 (1%)	12	1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3[A]	ILE
1	A	3[B]	ILE
1	A	313[A]	ILE
1	A	313[B]	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/263 (117%)	302 (98%)	5 (2%)	58	22

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	GLU
1	A	206[A]	LYS
1	A	206[B]	LYS
1	A	313[A]	ILE
1	A	313[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	258	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 2 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	EDO	A	428	-	3,3,3	0.87	0	2,2,2	0.42	0
9	GOL	A	403	-	5,5,5	3.86	4 (80%)	5,5,5	1.15	1 (20%)
5	EDO	A	424	-	3,3,3	0.91	0	2,2,2	0.03	0
7	DXE	A	443	-	5,5,5	0.70	0	4,4,4	1.90	1 (25%)
9	GOL	A	401	-	5,5,5	4.26	3 (60%)	5,5,5	2.58	2 (40%)
5	EDO	A	426	-	3,3,3	1.17	0	2,2,2	0.55	0
5	EDO	A	423	-	3,3,3	0.89	0	2,2,2	1.64	1 (50%)
8	MXE	A	451	-	4,4,4	0.95	0	3,3,3	4.76	2 (66%)
10	PEG	A	461	-	6,6,6	0.69	0	5,5,5	3.02	3 (60%)
3	ME2	A	471	-	9,9,9	0.96	0	8,8,8	1.13	1 (12%)
5	EDO	A	427	-	3,3,3	0.50	0	2,2,2	1.78	1 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	DXE	A	441	8	5,5,5	0.96	0	4,4,4	1.17	0
6	PGE	A	433	-	9,9,9	1.01	0	8,8,8	1.75	2 (25%)
4	MES	A	411	-	12,12,12	2.33	5 (41%)	15,16,16	3.73	9 (60%)
4	MES	A	412	-	12,12,12	1.80	4 (33%)	15,16,16	2.49	5 (33%)
5	EDO	A	422	-	3,3,3	0.65	0	2,2,2	2.15	2 (100%)
10	PEG	A	462	-	6,6,6	1.19	0	5,5,5	2.16	2 (40%)
7	DXE	A	442	-	5,5,5	0.58	0	4,4,4	3.05	3 (75%)
5	EDO	A	421	-	3,3,3	0.50	0	2,2,2	0.51	0
8	MXE	A	452	7	4,4,4	0.50	0	3,3,3	3.53	2 (66%)
5	EDO	A	425	-	3,3,3	0.70	0	2,2,2	0.68	0
6	PGE	A	434	-	9,9,9	1.07	0	8,8,8	2.18	5 (62%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	428	-	-	0/1/1/1	-
9	GOL	A	403	-	-	0/4/4/4	-
5	EDO	A	424	-	-	0/1/1/1	-
7	DXE	A	443	-	-	2/3/3/3	-
9	GOL	A	401	-	-	2/4/4/4	-
5	EDO	A	426	-	-	0/1/1/1	-
5	EDO	A	423	-	-	1/1/1/1	-
8	MXE	A	451	-	-	2/2/2/2	-
10	PEG	A	461	-	-	3/4/4/4	-
3	ME2	A	471	-	-	6/7/7/7	-
5	EDO	A	427	-	-	1/1/1/1	-
7	DXE	A	441	8	-	2/3/3/3	-
6	PGE	A	433	-	-	5/7/7/7	-
4	MES	A	411	-	-	0/6/14/14	0/1/1/1
4	MES	A	412	-	-	0/6/14/14	0/1/1/1
5	EDO	A	422	-	-	0/1/1/1	-
10	PEG	A	462	-	-	3/4/4/4	-
7	DXE	A	442	-	-	1/3/3/3	-
5	EDO	A	421	-	-	0/1/1/1	-
8	MXE	A	452	7	-	1/2/2/2	-
5	EDO	A	425	-	-	1/1/1/1	-

Continued on next page...



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	A	434	-	-	7/7/7/7	-

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	401	GOL	C3-C2	-8.13	1.20	1.51
9	A	403	GOL	C3-C2	-6.94	1.25	1.51
4	A	411	MES	O2S-S	4.49	1.57	1.45
4	A	411	MES	C8-S	3.96	1.83	1.77
4	A	412	MES	C5-N4	3.70	1.56	1.46

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	411	MES	O3S-S-O1S	8.00	131.41	111.40
4	A	411	MES	O1S-S-C8	-6.94	96.24	106.73
8	A	451	MXE	C3-O2-C2	6.81	153.96	112.90
4	A	412	MES	C2-C3-N4	-5.78	101.34	110.12
4	A	411	MES	O3S-S-C8	-5.70	94.85	106.00

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	A	401	GOL	O1-C1-C2-C3
10	A	462	PEG	C1-C2-O2-C3
3	A	471	ME2	C2-C3-O2-C4
6	A	434	PGE	C6-C5-O3-C4
7	A	442	DXE	O1-C2-C3-O2

There are no ring outliers.

20 monomers are involved in 135 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	428	EDO	4	0
5	A	424	EDO	2	0
7	A	443	DXE	10	0
5	A	426	EDO	2	0
5	A	423	EDO	4	0
8	A	451	MXE	8	0
10	A	461	PEG	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	471	ME2	11	0
5	A	427	EDO	11	0
7	A	441	DXE	17	0
6	A	433	PGE	6	0
4	A	411	MES	10	0
4	A	412	MES	7	0
5	A	422	EDO	3	0
10	A	462	PEG	2	0
7	A	442	DXE	1	0
5	A	421	EDO	2	0
8	A	452	MXE	29	0
5	A	425	EDO	5	0
6	A	434	PGE	10	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	314/314 (100%)	0.22	16 (5%) 34 31	3, 8, 17, 27	59 (18%)

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1[A]	MET	6.4
1	A	2[A]	GLU	6.1
1	A	3[A]	ILE	5.5
1	A	4	PRO	4.2
1	A	40	VAL	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
10	PEG	A	462	7/7	0.55	0.24	39,42,50,53	7
5	EDO	A	428	4/4	0.60	0.23	56,67,75,78	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
10	PEG	A	461	7/7	0.67	0.15	24,37,50,53	7
5	EDO	A	423	4/4	0.73	0.18	18,30,30,40	4
7	DXE	A	443	6/6	0.73	0.19	22,42,50,51	5
3	ME2	A	471	10/10	0.75	0.14	23,34,40,41	10
5	EDO	A	426	4/4	0.75	0.18	23,23,29,56	4
7	DXE	A	442	6/6	0.76	0.16	15,30,31,34	6
6	PGE	A	433	10/10	0.77	0.15	16,25,29,30	10
4	MES	A	412	12/12	0.79	0.19	16,34,44,45	12
4	MES	A	411	12/12	0.80	0.16	10,24,31,32	12
5	EDO	A	421	4/4	0.81	0.13	24,26,33,35	4
7	DXE	A	441	6/6	0.82	0.13	14,24,26,29	6
5	EDO	A	425	4/4	0.85	0.12	22,24,29,41	4
8	MXE	A	451	5/5	0.87	0.14	13,16,24,25	5
9	GOL	A	401	6/6	0.87	0.13	12,18,22,24	0
6	PGE	A	434	10/10	0.87	0.14	12,22,34,36	10
5	EDO	A	427	4/4	0.87	0.13	22,27,32,37	4
9	GOL	A	403	6/6	0.89	0.12	10,13,16,19	6
5	EDO	A	424	4/4	0.92	0.11	14,16,22,25	0
8	MXE	A	452	5/5	0.92	0.09	13,15,22,25	5
5	EDO	A	422	4/4	0.93	0.12	8,11,16,24	4
2	CA	A	491	1/1	1.00	0.01	4,4,4,4	0
2	CA	A	492	1/1	1.00	0.01	4,4,4,4	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.