



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 28, 2026 – 02:07 PM EDT

PDB ID : 9PSL / pdb_00009psl
Title : Co-crystal structure of BAF60A (SMARCD1) in complex with PU.1 peptide
Authors : Clasman, J.C.; Sadalge, D.S.; Von der Porten, L.; Wu, S.; Nelen, M.; Davenport, Y.W.; Bellon, S.F.; Taherbhoy, A.M.
Deposited on : 2025-07-25
Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	12% 86% 10%
1	C	223	21% 84% 13%
1	E	223	19% 87% 10%
1	G	223	19% 86% 11%
2	B	48	25% 94% ••

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	48	17% 63% 6% 31%
2	F	48	15% 75% • 21%
2	H	48	19% 77% 6% 17%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8274 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily D member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	2	0
			1667	1067	293	301	6			
1	C	195	Total	C	N	O	S	0	1	0
			1587	1022	274	284	7			
1	E	200	Total	C	N	O	S	0	1	0
			1649	1059	290	293	7			
1	G	199	Total	C	N	O	S	0	1	0
			1645	1056	290	292	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	175	GLY	-	expression tag	UNP Q96GM5
A	176	SER	-	expression tag	UNP Q96GM5
C	175	GLY	-	expression tag	UNP Q96GM5
C	176	SER	-	expression tag	UNP Q96GM5
E	175	GLY	-	expression tag	UNP Q96GM5
E	176	SER	-	expression tag	UNP Q96GM5
G	175	GLY	-	expression tag	UNP Q96GM5
G	176	SER	-	expression tag	UNP Q96GM5


- Molecule 2 is a protein called Transcription factor PU.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	46	Total	C	N	O	S	0	0	0
			386	243	63	77	3			
2	D	33	Total	C	N	O	S	0	0	0
			278	174	48	53	3			
2	F	38	Total	C	N	O	S	0	0	0
			328	208	54	63	3			
2	H	40	Total	C	N	O	S	0	0	0
			324	203	55	63	3			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	59	MET	-	initiating methionine	UNP P17947
B	101	GLU	-	expression tag	UNP P17947
B	102	ASN	-	expression tag	UNP P17947
B	103	LEU	-	expression tag	UNP P17947
B	104	TYR	-	expression tag	UNP P17947
B	105	PHE	-	expression tag	UNP P17947
B	106	GLN	-	expression tag	UNP P17947
D	59	MET	-	initiating methionine	UNP P17947
D	101	GLU	-	expression tag	UNP P17947
D	102	ASN	-	expression tag	UNP P17947
D	103	LEU	-	expression tag	UNP P17947
D	104	TYR	-	expression tag	UNP P17947
D	105	PHE	-	expression tag	UNP P17947
D	106	GLN	-	expression tag	UNP P17947
F	59	MET	-	initiating methionine	UNP P17947
F	101	GLU	-	expression tag	UNP P17947
F	102	ASN	-	expression tag	UNP P17947
F	103	LEU	-	expression tag	UNP P17947
F	104	TYR	-	expression tag	UNP P17947
F	105	PHE	-	expression tag	UNP P17947
F	106	GLN	-	expression tag	UNP P17947
H	59	MET	-	initiating methionine	UNP P17947
H	101	GLU	-	expression tag	UNP P17947
H	102	ASN	-	expression tag	UNP P17947
H	103	LEU	-	expression tag	UNP P17947
H	104	TYR	-	expression tag	UNP P17947
H	105	PHE	-	expression tag	UNP P17947
H	106	GLN	-	expression tag	UNP P17947

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	G	1	Total	C	O	0	0
			4	2	2		
3	G	1	Total	C	O	0	0
			4	2	2		
3	G	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	82	Total	O	0	4
			86	86		
5	B	26	Total	O	0	1
			26	26		
5	C	57	Total	O	0	0
			57	57		
5	D	11	Total	O	0	0
			11	11		
5	E	70	Total	O	0	5
			75	75		
5	F	20	Total	O	0	0
			20	20		
5	G	70	Total	O	0	1
			71	71		
5	H	16	Total	O	0	0
			16	16		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family D member 1

Chain A: 




- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family D member 1

Chain C: 





- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family D member 1


Chain E: 




- Molecule 1: SWI/SNF-related matrix-associated actin-dependent regulator of chromatin sub-family D member 1





- Molecule 2: Transcription factor PU.1





- Molecule 2: Transcription factor PU.1



- Molecule 2: Transcription factor PU.1



- Molecule 2: Transcription factor PU.1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.75Å 131.85Å 75.36Å 90.00° 93.44° 90.00°	Depositor
Resolution (Å)	49.63 – 2.39 49.63 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.63-2.39) 99.6 (49.63-2.39)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.224 , 0.284 0.231 , 0.285	Depositor DCC
R_{free} test set	2341 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8274	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/1705	0.79	0/2310
1	C	0.46	0/1624	0.79	0/2207
1	E	0.46	0/1687	0.79	0/2286
1	G	0.46	0/1683	0.80	0/2280
2	B	0.45	0/395	0.85	0/534
2	D	0.45	0/284	0.83	0/385
2	F	0.43	0/336	0.85	0/454
2	H	0.46	0/331	0.87	0/449
All	All	0.46	0/8045	0.80	0/10905

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1667	0	1656	9	0
1	C	1587	0	1569	5	0
1	E	1649	0	1647	4	0
1	G	1645	0	1650	3	0
2	B	386	0	350	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	278	0	266	3	0
2	F	328	0	309	1	0
2	H	324	0	297	1	0
3	A	8	0	12	0	0
3	C	8	0	12	0	0
3	G	12	0	18	0	0
4	B	5	0	0	0	0
4	D	5	0	0	0	0
4	F	5	0	0	0	0
4	H	5	0	0	0	0
5	A	86	0	0	1	0
5	B	26	0	0	0	0
5	C	57	0	0	0	0
5	D	11	0	0	0	0
5	E	75	0	0	0	0
5	F	20	0	0	0	0
5	G	71	0	0	0	0
5	H	16	0	0	0	0
All	All	8274	0	7786	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 23 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:GLU:OE1	1:A:210[A]:ARG:NH2	2.37	0.57
1:A:210[A]:ARG:HH11	1:A:210[A]:ARG:HG2	1.71	0.56
1:E:234:SER:HB3	1:E:267:THR:HG22	1.89	0.54
1:G:190:PHE:C	1:G:191:ASN:HD22	2.20	0.50
1:A:240:LEU:C	1:A:240:LEU:HD23	2.37	0.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/223 (88%)	196 (100%)	1 (0%)	0	100	100
1	C	190/223 (85%)	186 (98%)	4 (2%)	0	100	100
1	E	195/223 (87%)	192 (98%)	3 (2%)	0	100	100
1	G	194/223 (87%)	193 (100%)	1 (0%)	0	100	100
2	B	44/48 (92%)	42 (96%)	2 (4%)	0	100	100
2	D	31/48 (65%)	31 (100%)	0	0	100	100
2	F	36/48 (75%)	36 (100%)	0	0	100	100
2	H	38/48 (79%)	38 (100%)	0	0	100	100
All	All	925/1084 (85%)	914 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/205 (91%)	186 (100%)	0	100	100
1	C	175/205 (85%)	175 (100%)	0	100	100
1	E	183/205 (89%)	182 (100%)	1 (0%)	81	91
1	G	184/205 (90%)	181 (98%)	3 (2%)	55	76
2	B	43/47 (92%)	43 (100%)	0	100	100
2	D	33/47 (70%)	33 (100%)	0	100	100
2	F	38/47 (81%)	37 (97%)	1 (3%)	40	63
2	H	36/47 (77%)	35 (97%)	1 (3%)	38	60
All	All	878/1008 (87%)	872 (99%)	6 (1%)	76	88

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	271	GLN
1	G	337	VAL
2	H	95	VAL
2	F	80	GLN
1	E	188	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	344	GLN
2	H	69	ASN
2	H	68	ASN
1	E	271	GLN
1	G	329	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	B	201	-	4,4,4	0.35	0	6,6,6	0.08	0
4	SO4	F	201	-	4,4,4	0.35	0	6,6,6	0.06	0
3	EDO	G	401	-	3,3,3	0.10	0	2,2,2	0.15	0
3	EDO	A	402	-	3,3,3	0.06	0	2,2,2	0.06	0
3	EDO	G	402	-	3,3,3	0.07	0	2,2,2	0.17	0
3	EDO	A	401	-	3,3,3	0.12	0	2,2,2	0.23	0
3	EDO	C	402	-	3,3,3	0.05	0	2,2,2	0.12	0
3	EDO	G	403	-	3,3,3	0.08	0	2,2,2	0.13	0
4	SO4	H	201	-	4,4,4	0.35	0	6,6,6	0.10	0
4	SO4	D	201	-	4,4,4	0.35	0	6,6,6	0.07	0
3	EDO	C	401	-	3,3,3	0.06	0	2,2,2	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	G	401	-	-	1/1/1/1	-
3	EDO	A	402	-	-	1/1/1/1	-
3	EDO	G	402	-	-	1/1/1/1	-
3	EDO	A	401	-	-	1/1/1/1	-
3	EDO	C	402	-	-	1/1/1/1	-
3	EDO	G	403	-	-	1/1/1/1	-
3	EDO	C	401	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	EDO	O1-C1-C2-O2
3	A	402	EDO	O1-C1-C2-O2
3	G	401	EDO	O1-C1-C2-O2
3	G	403	EDO	O1-C1-C2-O2
3	G	402	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/223 (90%)	1.13	27 (13%) 7 5	26, 58, 88, 124	2 (0%)
1	C	195/223 (87%)	1.45	47 (24%) 2 1	30, 65, 98, 115	1 (0%)
1	E	200/223 (89%)	1.42	43 (21%) 2 2	29, 66, 104, 117	1 (0%)
1	G	199/223 (89%)	1.31	43 (21%) 2 2	28, 61, 91, 100	1 (0%)
2	B	46/48 (95%)	1.61	12 (26%) 1 1	46, 61, 106, 109	0
2	D	33/48 (68%)	1.49	8 (24%) 2 1	47, 62, 90, 104	0
2	F	38/48 (79%)	1.32	7 (18%) 3 3	48, 62, 81, 91	0
2	H	40/48 (83%)	1.55	9 (22%) 2 2	49, 61, 100, 109	0
All	All	952/1084 (87%)	1.36	196 (20%) 2 2	26, 62, 97, 124	5 (0%)

The worst 5 of 196 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	100	MET	5.1
1	E	193	ALA	5.0
2	B	105	PHE	4.9
2	B	62	PHE	4.7
1	A	219	SER	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	402	4/4	0.77	0.24	66,69,71,72	0
3	EDO	A	401	4/4	0.80	0.24	64,67,71,72	0
3	EDO	G	402	4/4	0.81	0.20	69,70,71,72	0
3	EDO	G	403	4/4	0.81	0.21	82,82,83,84	0
3	EDO	C	402	4/4	0.83	0.24	72,73,75,75	0
3	EDO	G	401	4/4	0.84	0.21	64,69,72,73	0
3	EDO	C	401	4/4	0.86	0.21	66,66,68,69	0
4	SO4	B	201	5/5	0.96	0.09	57,58,61,67	0
4	SO4	H	201	5/5	0.96	0.09	58,60,61,65	0
4	SO4	F	201	5/5	0.97	0.07	48,50,51,52	0
4	SO4	D	201	5/5	0.97	0.07	56,56,57,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.