



wwPDB EM Validation Summary Report ⓘ

Jul 27, 2026 – 03:17 PM EDT

PDB ID : 9PT8 / pdb_00009pt8
EMDB ID : EMD-71835
Title : Alpha-synuclein acetylated lysine 80
Authors : Li, M.H.
Deposited on : 2025-07-28
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	
1	B	140	
1	C	140	
1	D	140	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	140	
1	F	140	
1	G	140	
1	H	140	
1	I	140	
1	J	140	
1	K	140	
1	L	140	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-synuclein.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	60	Total 411	C 254	N 73	O 84	0	0
1	B	60	Total 411	C 254	N 73	O 84	0	0
1	C	60	Total 411	C 254	N 73	O 84	0	0
1	D	60	Total 411	C 254	N 73	O 84	0	0
1	E	60	Total 411	C 254	N 73	O 84	0	0
1	F	60	Total 411	C 254	N 73	O 84	0	0
1	G	60	Total 411	C 254	N 73	O 84	0	0
1	H	60	Total 411	C 254	N 73	O 84	0	0
1	I	60	Total 411	C 254	N 73	O 84	0	0
1	J	60	Total 411	C 254	N 73	O 84	0	0
1	K	60	Total 411	C 254	N 73	O 84	0	0
1	L	60	Total 411	C 254	N 73	O 84	0	0

LEU GLY ASP LYS ASN ASN GLU GLY GLY ALA PRO PRO GLN GLY ILE LEU GLU ASP MET MET PRO VAL ASP PRO ASP ASN GLU ALA TYR GLU MET MET PRO PRO SER SER GLU GLY GLY TYR GLN ASP TYR GLU PRO GLU ALA

• Molecule 1: Alpha-synuclein

Chain E:  35% 8% 57%

MET ASP VAL PHE MET LYS GLY LEU SER LYS LYS ALA GLY ILE LEU GLU VAL ASP VAL ALA ALA ALA GLU LYS THR ASP LYS GLN GLY VAL ALA GLU TYR V40 V49 H50 G51 V55 T75 K30 T81 V82 E83 I88 K96 K97 D98 Q99

LEU GLY LYS ASN ASN GLU GLY GLY ALA PRO PRO GLN GLY ILE LEU GLU ASP MET MET PRO VAL ASP PRO ASP ASN GLU ALA TYR GLU MET MET PRO PRO SER SER GLU GLY TYR VAL ALA GLU TYR GLN GLY LYS THR GLY TYR GLN GLY VAL TYR GLU PRO GLU ALA

• Molecule 1: Alpha-synuclein

Chain F:  38% 5% 57%

MET ASP VAL PHE MET LYS GLY LEU SER LYS LYS ALA GLY ILE LEU GLU VAL ASP VAL ALA ALA ALA GLU LYS THR ASP LYS GLN GLY VAL ALA GLU TYR V40 H50 G51 T75 V82 E83 K96 K97 D98 Q99 LEU GLY LYS ASN GLU GLY

ALA PRO GLN GLU ILE LEU GLU ASP MET PRO VAL ASP PRO ASP ASN GLU ALA TYR GLU MET MET LYS THR ASP LYS GLN GLY TYR VAL ALA GLU TYR ASP TYR GLU PRO GLU ALA

• Molecule 1: Alpha-synuclein

Chain G:  36% 6% 57%

MET ASP VAL PHE MET LYS GLY LEU SER LYS LYS ALA GLY ILE LEU GLU VAL ASP VAL ALA ALA ALA GLU LYS THR ASP LYS GLN GLY VAL ALA GLU TYR V40 H50 G51 V55 T59 K80 T81 V82 E83 K96 K97 D98 Q99 LEU GLY LYS

ASN GLU GLY ALA MET LYS LYS ILE LEU GLU ASP MET PRO VAL ASP ASP ASP ASN GLU LYS THR ALA TYR LYS MET PRO SER GLU GLU ALA GLY TYR GLN ASP TYR LYS PRO GLU ALA

• Molecule 1: Alpha-synuclein

Chain H:  34% 9% 57%

MET ASP VAL PHE MET LYS GLY LEU SER LYS LYS ALA GLY ILE LEU VAL VAL ALA ALA ALA ALA GLU LYS THR LYS LYS GLN GLY VAL VAL ALA ALA GLU TYR V40 E46 V49 H50 G51 V55 T59 T75 K80 T81 V82 E83 K96 K97

D98 Q99 LEU GLY LYS ASN ASN GLU GLY ALA PRO PRO GLN GLY ILE LEU GLU ASP MET MET PRO VAL ASP PRO ASP ASN GLU ALA TYR GLU ALA ALA MET PRO LYS THR LYS LYS GLN GLY TYR ASP TYR GLU PRO GLU ALA



• Molecule 1: Alpha-synuclein

Chain I:  34% 9% 57%



MET ASP VAL PHE MET LYS GLY LEU SER LYS LYS ALA GLY ILE LEU VAL VAL ALA ALA ALA ALA GLU LYS THR LYS LYS GLN GLY VAL VAL ALA ALA GLU TYR V40 E46 V49 H50 G51 V55 T59 T75 K80 T81 V82 E83 I88





• Molecule 1: Alpha-synuclein



• Molecule 1: Alpha-synuclein



• Molecule 1: Alpha-synuclein



4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-1.11°, rise=4.76 Å, axial sym=C1	Depositor
Number of segments used	296473	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	14.8	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.103	Depositor
Minimum map value	-0.057	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.017	Depositor
Map size (Å)	296.96, 296.96, 296.96	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.16, 1.16, 1.16	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/412	0.31	0/556
1	B	0.11	0/412	0.31	0/556
1	C	0.11	0/412	0.31	0/556
1	D	0.11	0/412	0.31	0/556
1	E	0.11	0/412	0.31	0/556
1	F	0.11	0/412	0.31	0/556
1	G	0.11	0/412	0.31	0/556
1	H	0.11	0/412	0.31	0/556
1	I	0.11	0/412	0.31	0/556
1	J	0.11	0/412	0.31	0/556
1	K	0.11	0/412	0.31	0/556
1	L	0.11	0/412	0.31	0/556
All	All	0.11	0/4944	0.31	0/6672

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	411	0	430	9	0
1	B	411	0	430	13	0
1	C	411	0	430	12	0
1	D	411	0	430	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	411	0	430	14	0
1	F	411	0	430	6	0
1	G	411	0	430	8	0
1	H	411	0	430	13	0
1	I	411	0	430	16	0
1	J	411	0	430	15	0
1	K	411	0	430	11	0
1	L	411	0	430	5	0
All	All	4932	0	5160	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 83 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:VAL:HG11	1:K:51:GLY:HA2	1.78	0.66
1:B:51:GLY:HA2	1:K:55:VAL:HG11	1.85	0.59
1:A:51:GLY:HA2	1:L:55:VAL:HG11	1.85	0.57
1:B:55:VAL:HG11	1:J:51:GLY:HA2	1.87	0.56
1:I:98:ASP:HB2	1:J:98:ASP:OD1	2.06	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	B	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	C	58/140 (41%)	57 (98%)	1 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	E	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	F	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	G	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	H	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	I	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	J	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	K	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
1	L	58/140 (41%)	57 (98%)	1 (2%)	0	100	100
All	All	696/1680 (41%)	684 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	42/103 (41%)	42 (100%)	0	100	100
1	B	42/103 (41%)	42 (100%)	0	100	100
1	C	42/103 (41%)	42 (100%)	0	100	100
1	D	42/103 (41%)	42 (100%)	0	100	100
1	E	42/103 (41%)	42 (100%)	0	100	100
1	F	42/103 (41%)	42 (100%)	0	100	100
1	G	42/103 (41%)	42 (100%)	0	100	100
1	H	42/103 (41%)	42 (100%)	0	100	100
1	I	42/103 (41%)	42 (100%)	0	100	100
1	J	42/103 (41%)	42 (100%)	0	100	100
1	K	42/103 (41%)	42 (100%)	0	100	100
1	L	42/103 (41%)	42 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	504/1236 (41%)	504 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	79	GLN
1	I	79	GLN
1	L	79	GLN
1	J	79	GLN
1	D	79	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-71835. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 123




Y Index: 110




Z Index: 119


6.3.2 Raw map



X Index: 128



Y Index: 142




Z Index: 119


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.017. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_71835_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 65 nm³; this corresponds to an approximate mass of 58 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.333 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	2.95	3.31	2.99
Unmasked-calculated*	3.24	3.52	3.28

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-71835 and PDB model 9PT8. Per-residue inclusion information can be found in section [3](#) on page [5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.017 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.017).

9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.017) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7620	0.5720
A	0.7590	0.5810
B	0.7790	0.5810
C	0.7760	0.5800
D	0.7660	0.5730
E	0.7710	0.5750
F	0.7710	0.5670
G	0.7400	0.5660
H	0.7540	0.5720
I	0.7450	0.5660
J	0.7620	0.5630
K	0.7590	0.5700
L	0.7590	0.5680

