



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 03:20 pm BST

PDB ID : 6Q3E / pdb\_00006q3e  
Title : Dye type peroxidase Aa from Streptomyces lividans: 274.4 kGy structure  
Authors : Ebrahim, A.; Moreno-Chicano, T.; Worrall, J.A.R.; Strange, R.W.; Axford, D.; Sherrell, D.A.; Appleby, M.; Owen, R.L.  
Deposited on : 2018-12-04  
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	12358 (2.04-2.00)
Clashscore	180529	13897 (2.04-2.00)
Ramachandran outliers	177936	13770 (2.04-2.00)
Sidechain outliers	177891	13769 (2.04-2.00)
RSRZ outliers	164620	12358 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span>%</span> <div style="width: 81%; height: 10px; background-color: green;"></div> <div style="width: 19%; height: 10px; background-color: yellow;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span></span> <span>81%</span> <span>19%</span> </div> </div>
1	B	362	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span>%</span> <div style="width: 82%; height: 10px; background-color: green;"></div> <div style="width: 18%; height: 10px; background-color: yellow;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> <span></span> <span>82%</span> <span>18%</span> </div> </div>



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Deferrochelataase/peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total 2773	C 1745	N 509	O 514	S 5	0	0	0
1	B	361	Total 2783	C 1752	N 510	O 516	S 5	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	228	LYS	GLU	conflict	UNP Q9RKQ2
B	228	LYS	GLU	conflict	UNP Q9RKQ2

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	104	Total	O	0	0
			104	104		
3	B	107	Total	O	0	0
			107	107		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deferrochelataase/peroxidase



- Molecule 1: Deferrochelataase/peroxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.07Å 68.38Å 75.10Å 90.00° 105.85° 90.00°	Depositor
Resolution (Å)	37.39 – 2.03 37.39 – 2.03	Depositor EDS
% Data completeness (in resolution range)	100.0 (37.39-2.03) 93.1 (37.39-2.03)	Depositor EDS
$R_{merge}$	0.77	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.45 (at 2.03Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.182 , 0.249 0.182 , 0.250	Depositor DCC
$R_{free}$ test set	44150 reflections (4.33%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.4	Xtriage
Anisotropy	0.165	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 38.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.048 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5913	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2841	0.55	0/3855
1	B	0.36	0/2851	0.53	0/3869
All	All	0.36	0/5692	0.54	0/7724

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2773	0	2704	49	0
1	B	2783	0	2715	49	0
2	A	43	30	30	4	0
2	B	43	30	30	4	0
3	A	104	0	0	8	0
3	B	107	0	0	7	0
All	All	5853	60	5479	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:ASP:HB2	3:A:695:HOH:O	1.82	0.80
1:B:108:GLY:HA3	3:B:618:HOH:O	1.82	0.79
1:B:308:GLY:HA3	3:B:643:HOH:O	1.86	0.74
1:A:254:ARG:HD3	1:A:255:PHE:CE2	2.24	0.73
1:B:358:ASP:HB3	3:B:629:HOH:O	1.91	0.70
1:B:404:VAL:HG13	1:B:409:ASP:HB3	1.71	0.69
2:A:501:HEM:HBB2	2:A:501:HEM:HMB2	1.74	0.69
1:A:268:THR:O	1:A:402:PRO:HG3	1.94	0.67
1:A:367:GLN:HE21	1:A:374:PHE:H	1.43	0.66
1:B:260:GLU:HG3	3:B:676:HOH:O	1.94	0.66
1:A:212:ARG:HG2	1:A:213:TRP:HD1	1.61	0.64
1:A:367:GLN:NE2	1:A:374:PHE:H	1.97	0.63
2:B:501:HEM:HMB2	2:B:501:HEM:HBB2	1.82	0.60
1:B:404:VAL:CG1	1:B:409:ASP:HB3	2.32	0.60
1:B:244:ILE:HD11	1:B:339:THR:HG22	1.84	0.59
1:B:259:ALA:O	1:B:262:ASP:HB2	2.04	0.58
1:B:81:PHE:HB2	1:B:182:LEU:CD1	2.34	0.57
1:A:409:ASP:OD1	1:A:410:TRP:N	2.36	0.57
1:A:82:ASP:HA	1:A:181:ASP:HB2	1.87	0.57
1:B:244:ILE:HD12	1:B:340:LEU:O	2.06	0.56
1:B:111:VAL:HG11	1:B:204:ILE:HD11	1.88	0.55
1:B:135:PRO:HG3	3:B:618:HOH:O	2.06	0.55
1:A:168:PHE:O	1:A:171:ASP:HB2	2.07	0.55
2:A:501:HEM:HBB2	2:A:501:HEM:CMB	2.37	0.55
1:A:285:TRP:CH2	1:A:293:GLN:HG2	2.42	0.55
1:A:403:GLY:O	1:A:405:ARG:HD2	2.07	0.55
1:A:162:LEU:HD11	1:A:397:VAL:HG23	1.88	0.54
1:B:81:PHE:HB2	1:B:182:LEU:HD11	1.89	0.54
1:B:127:THR:HG21	1:B:199:ARG:NE	2.23	0.54
1:A:104:ARG:NH2	1:A:109:HIS:HB3	2.24	0.53
1:A:266:TRP:CZ2	1:A:415:LEU:HD13	2.45	0.52
1:A:415:LEU:HD23	1:A:416:PHE:CE1	2.45	0.52
1:B:74:ASP:HB3	1:B:75:ARG:NH1	2.25	0.52
1:B:290:LEU:O	1:B:294:GLU:HG3	2.09	0.52
1:B:274:VAL:HG22	1:B:397:VAL:HG12	1.92	0.51
2:B:501:HEM:HBB2	2:B:501:HEM:CMB	2.40	0.51
1:A:285:TRP:CZ2	1:A:293:GLN:HG2	2.46	0.51
1:A:87:ASP:HB3	1:A:90:ALA:CB	2.41	0.51
1:B:297:PHE:O	1:B:327:VAL:HG23	2.10	0.51
1:B:351:THR:HG22	1:B:357:LEU:HA	1.92	0.49
1:A:191:PRO:O	1:A:195:VAL:HG23	2.12	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:68:ILE:HD11	1:B:400:VAL:HG11	1.93	0.49
1:A:283:GLU:OE1	1:A:356:ARG:HD3	2.12	0.49
1:A:280:MET:HG3	1:A:360:GLY:HA2	1.95	0.49
1:A:367:GLN:HE22	1:A:374:PHE:HB2	1.78	0.48
1:B:248:GLU:CD	1:B:251:ARG:HD3	2.38	0.48
1:A:138:LEU:HD12	1:A:187:CYS:O	2.12	0.48
1:B:244:ILE:HD11	1:B:339:THR:CG2	2.43	0.48
1:B:330:ALA:HB1	2:B:501:HEM:C2D	2.49	0.48
1:A:405:ARG:HG2	1:A:409:ASP:OD2	2.13	0.48
1:A:175:ARG:HG3	3:A:612:HOH:O	2.12	0.48
1:A:243:ASN:ND2	3:A:607:HOH:O	2.46	0.48
1:B:276:ARG:HD2	1:B:395:SER:HB2	1.96	0.48
1:A:315:GLU:OE1	1:A:316:PRO:HD2	2.13	0.47
1:B:362:PHE:CE2	1:B:364:LEU:HD21	2.49	0.47
1:A:110:ALA:HB3	1:A:114:GLY:HA2	1.97	0.46
1:A:119:LEU:HB2	3:A:690:HOH:O	2.14	0.46
1:A:299:ARG:NH1	1:A:304:GLY:O	2.45	0.46
1:A:388:GLU:O	1:A:388:GLU:HG3	2.15	0.46
1:B:226:PRO:O	1:B:229:GLN:HB3	2.16	0.45
1:A:261:LYS:HB3	1:A:261:LYS:HE2	1.78	0.45
1:A:75:ARG:HE	1:A:75:ARG:HB3	1.62	0.45
1:A:212:ARG:HG2	1:A:213:TRP:CD1	2.47	0.45
1:A:234:LEU:HD23	1:A:234:LEU:HA	1.56	0.45
2:A:501:HEM:HBC2	2:A:501:HEM:HMC1	1.99	0.45
1:A:87:ASP:HB3	1:A:90:ALA:HB3	1.98	0.44
1:A:56:SER:HB3	3:A:680:HOH:O	2.17	0.44
1:B:166:PRO:HG2	1:B:168:PHE:CZ	2.52	0.44
1:B:306:PRO:O	1:B:307:VAL:C	2.61	0.44
1:A:220:LYS:HG2	3:A:655:HOH:O	2.18	0.44
2:A:501:HEM:HBC2	2:A:501:HEM:CMC	2.48	0.44
1:B:349:ASP:C	3:B:629:HOH:O	2.60	0.43
1:A:348:THR:HG23	1:A:348:THR:O	2.18	0.43
1:B:74:ASP:HB2	1:B:189:ASP:HA	2.01	0.43
1:B:172:ASN:O	1:B:393:VAL:HA	2.19	0.43
1:B:289:SER:N	1:B:292:GLU:OE1	2.43	0.43
1:B:199:ARG:HG3	1:B:199:ARG:HH11	1.83	0.43
1:A:83:VAL:N	1:A:181:ASP:OD2	2.39	0.43
1:B:242:ARG:HH21	1:B:242:ARG:HG2	1.82	0.43
1:B:101:ALA:O	1:B:105:LEU:HG	2.20	0.42
1:A:144:PHE:HB3	1:A:148:LEU:HD23	2.01	0.42
1:A:401:PRO:HB2	1:A:402:PRO:HD2	2.02	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:HIS:ND1	1:B:195:VAL:HA	2.35	0.42
1:B:77:HIS:CD2	1:B:216:LEU:HD12	2.55	0.42
1:B:260:GLU:HA	1:B:268:THR:OG1	2.20	0.42
1:A:68:ILE:HB	3:A:620:HOH:O	2.20	0.42
1:B:251:ARG:O	1:B:255:PHE:HB2	2.20	0.42
1:A:367:GLN:HE22	1:A:374:PHE:CB	2.33	0.41
1:B:244:ILE:CD1	1:B:339:THR:HG22	2.48	0.41
1:B:341[B]:LEU:C	1:B:341[B]:LEU:HD23	2.45	0.41
1:B:134:LYS:HB3	1:B:135:PRO:CD	2.50	0.41
1:B:134:LYS:HB3	1:B:135:PRO:HD2	2.03	0.41
1:B:199:ARG:HG3	1:B:199:ARG:NH1	2.35	0.41
1:B:242:ARG:NE	2:B:501:HEM:O1D	2.50	0.41
1:A:361:LEU:HD23	1:A:363:PHE:HB2	2.01	0.41
1:A:279:ARG:NH2	1:A:358:ASP:OD2	2.54	0.41
1:B:131:LEU:O	3:B:602:HOH:O	2.22	0.41
1:A:94:LEU:HD22	1:A:208:LYS:HB2	2.03	0.41
1:B:235:LEU:HD22	1:B:285:TRP:CD2	2.56	0.41
1:A:173:LEU:HD13	1:A:395:SER:HB3	2.03	0.40
1:A:416:PHE:C	3:A:649:HOH:O	2.65	0.40
1:A:161:ALA:HB3	1:A:272:TYR:CZ	2.55	0.40
1:B:70:THR:O	1:B:71:PRO:C	2.64	0.40
1:B:81:PHE:HB2	1:B:182:LEU:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/362 (99%)	348 (97%)	11 (3%)	1 (0%)	37	34
1	B	361/362 (100%)	348 (96%)	13 (4%)	0	100	100
All	All	721/724 (100%)	696 (96%)	24 (3%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	56	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/277 (100%)	276 (100%)	1 (0%)	89	92
1	B	279/277 (101%)	279 (100%)	0	100	100
All	All	556/554 (100%)	555 (100%)	1 (0%)	92	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	249	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	367	GLN
1	A	378	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	B	501	1,3	41,50,50	1.48	6 (14%)	45,82,82	1.37	4 (8%)
2	HEM	A	501	1	41,50,50	1.58	6 (14%)	45,82,82	1.38	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	501	1,3	-	4/12/54/54	-
2	HEM	A	501	1	-	5/12/54/54	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	C3C-C2C	-4.70	1.33	1.40
2	B	501	HEM	C3C-C2C	-3.58	1.35	1.40
2	B	501	HEM	C3C-CAC	3.48	1.54	1.47
2	A	501	HEM	C3C-CAC	3.23	1.54	1.47
2	A	501	HEM	CAA-C2A	2.93	1.56	1.52
2	B	501	HEM	CAB-C3B	2.83	1.55	1.47
2	A	501	HEM	FE-ND	2.76	2.10	1.96
2	B	501	HEM	CAA-C2A	2.67	1.56	1.52
2	A	501	HEM	CAB-C3B	2.62	1.54	1.47
2	A	501	HEM	C1B-NB	-2.51	1.36	1.40
2	B	501	HEM	CMD-C2D	2.17	1.55	1.50
2	B	501	HEM	CMB-C2B	2.06	1.55	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	CBA-CAA-C2A	4.39	120.11	112.62
2	B	501	HEM	C4C-CHD-C1D	3.25	126.84	122.56
2	A	501	HEM	C1B-NB-C4B	2.60	107.76	105.07
2	B	501	HEM	CMC-C2C-C3C	2.56	129.48	124.68
2	B	501	HEM	C4B-CHC-C1C	2.44	125.77	122.56
2	A	501	HEM	CMA-C3A-C4A	-2.43	124.72	128.46
2	B	501	HEM	C1B-NB-C4B	2.43	107.58	105.07
2	A	501	HEM	C4A-C3A-C2A	2.41	108.67	107.00
2	A	501	HEM	C4C-CHD-C1D	2.15	125.39	122.56
2	A	501	HEM	O1A-CGA-CBA	-2.04	116.54	123.08

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	HEM	CAD-CBD-CGD-O1D
2	A	501	HEM	CAD-CBD-CGD-O2D
2	B	501	HEM	CAD-CBD-CGD-O1D
2	B	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAA-CBA-CGA-O2A
2	B	501	HEM	CAA-CBA-CGA-O1A
2	B	501	HEM	CAA-CBA-CGA-O2A
2	A	501	HEM	CAA-CBA-CGA-O1A
2	A	501	HEM	C2A-CAA-CBA-CGA


There are no ring outliers.


2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	HEM	4	0
2	A	501	HEM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	362/362 (100%)	0.15	2 (0%) 85 85	21, 31, 49, 62	0
1	B	361/362 (99%)	0.10	5 (1%) 73 73	10, 29, 50, 86	2 (0%)
All	All	723/724 (99%)	0.12	7 (0%) 79 79	10, 30, 50, 86	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	225	THR	3.2
1	B	226	PRO	2.7
1	B	224	THR	2.5
1	A	357	LEU	2.3
1	B	228	LYS	2.3
1	B	341[A]	LEU	2.2
1	A	55	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.






Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HEM	A	501	43/43	0.96	0.07	14,26,34,34	0
2	HEM	B	501	43/43	0.97	0.06	16,27,35,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around HEM A 501:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.