



wwPDB EM Validation Summary Report ⓘ

Jul 23, 2026 – 01:37 pm BST

PDB ID : 9QGK / pdb_00009qgk
EMDB ID : EMD-53133
Title : F-actin decorated by ITPKA
Authors : Yuan, B.; Paraschiakos, T.; Windhorst, S.; Marlovits, T.C.
Deposited on : 2025-03-13
Resolution : 2.97 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY




The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13205 (2.47 - 3.47)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	 5% 95%
1	B	461	 5% 95%
1	C	461	 5% 95%
1	D	461	 5% 95%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	461	 5% 95%
2	F	377	 89% 9%
2	G	377	 89% 9%
2	H	377	 88% 11%
2	I	377	 89% 9%
2	J	377	 88% 10%
3	X	7	 29% 43% 29%
3	Y	7	 29% 43% 29%
3	Z	7	 29% 57% 14%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 30895 atoms, of which 15255 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol-trisphosphate 3-kinase A.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	23	Total	C	H	N	O	S	0	0
			344	103	177	35	28	1		
1	B	23	Total	C	H	N	O	S	0	0
			344	103	177	35	28	1		
1	C	23	Total	C	H	N	O	S	0	0
			344	103	177	35	28	1		
1	D	23	Total	C	H	N	O	S	0	0
			344	103	177	35	28	1		
1	E	23	Total	C	H	N	O	S	0	0
			344	103	177	35	28	1		

- Molecule 2 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	F	371	Total	C	H	N	O	S	0	0
			5774	1837	2874	489	553	21		
2	G	371	Total	C	H	N	O	S	0	0
			5774	1837	2874	489	553	21		
2	H	371	Total	C	H	N	O	S	0	0
			5774	1837	2874	489	553	21		
2	I	371	Total	C	H	N	O	S	0	0
			5774	1837	2874	489	553	21		
2	J	371	Total	C	H	N	O	S	0	0
			5774	1837	2874	489	553	21		


- Molecule 3 is a protein called Phalloidin.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	X	7	Total	C	N	O	S	0	0
			55	35	8	11	1		
3	Y	7	Total	C	N	O	S	0	0
			55	35	8	11	1		
3	Z	7	Total	C	N	O	S	0	0
			55	35	8	11	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
4	F	1	Total	Mg	0
			1	1	
4	G	1	Total	Mg	0
			1	1	
4	H	1	Total	Mg	0
			1	1	
4	I	1	Total	Mg	0
			1	1	
4	J	1	Total	Mg	0
			1	1	

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
5	F	1	Total	C	N	O	P	0
			27	10	5	10	2	
5	G	1	Total	C	N	O	P	0
			27	10	5	10	2	
5	H	1	Total	C	N	O	P	0
			27	10	5	10	2	
5	I	1	Total	C	N	O	P	0
			27	10	5	10	2	
5	J	1	Total	C	N	O	P	0
			27	10	5	10	2	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Inositol-trisphosphate 3-kinase A



• Molecule 1: Inositol-trisphosphate 3-kinase A



TRP	GLU	ARG	ASN	TYR	PHE
	GLU	ARG	LEU		GLY
	GLU	LEU	GLN		TRP
	GLY	GLN	ARG		ASP
	ARG	ILE	ILE		GLU
	GLU	ARG	GLY		GLY
	ASP	ASP	ILE		ASP
	GLY	THR	THR		SER
	TYR	LEU	LEU		SER
	LEU	GLU	GLU		MET
LEU	VAL	VAL	THR	GLY	
GLY	SER	SER	THR	VAL	
LEU	GLU	GLU	GLY	ARG	
ASP	PHE	PHE	LEU	THR	
ASP	PHE	PHE	ARG	THR	
GLU	HIS	HIS	ILE	THR	
LEU	GLY	GLY	GLY	LYS	
LEU	VAL	VAL	LYS	ALA	
LEU	ILE	ILE	ALA	LYS	
LEU	SER	SER	SER	ALA	
LEU	SER	SER	GLY	ALA	
ALA	SER	SER	LEU	ARG	
GLU	LEU	LEU	LEU	GLU	
ARG	PHE	PHE	VAL	GLY	
	VAL	HIS	VAL	GLY	
	HIS	HIS	ASP	GLY	
	GLY	GLY	ASP	GLY	
	VAL	VAL	ASP	GLY	
	TRP	TRP	THR	GLU	
	LEU	LEU	THR	GLU	
	LEU	ILE	THR	THR	
	ASP	ASP	PRO	PHE	
	GLY	GLY	LEU	VAL	
	GLN	GLN	GLU	GLN	
	ILE	ILE	GLU	ARG	
	LEU	LEU	VAL	ALA	
	ASP	ASP	LEU	VAL	
	HIS	HIS	ARG	THR	
	ARG	ARG	LYS	ARG	
	PRO	PRO	THR	PRO	
	LEU	LEU	THR	LEU	

- Molecule 1: Inositol-trisphosphate 3-kinase A

Chain C: 5% 95%

[illegible]

- Molecule 1: Inositol-trisphosphate 3-kinase A


Chain D: 5% 95%

GLY	SER	SER	SER	PRO	VAL	THR	MET
CYS	PHE	LYS	GLU	ASP	ILE	LEU	THR
VAL	LYS	ASP	ASP	PRO	PRO	GLN	PRO
LEU	ALA	LEU	LEU	LEU	LEU	LEU	GLY
ASP	GLY	THR	THR	THR	THR	THR	GLY
CYS	THR	SER	SER	ASP	VAL	THR	THR
LYS	SER	GLY	SER	SER	THR	THR	GLY
MET	GLY	LEU	GLU	GLU	ALA	ALA	MET
GLY	ILE	SER	SER	GLU	GLU	ALA	ALA
VAL	LEU	LEU	ARG	ARG	GLU	GLU	ARG
ARG	LEU	LYS	ARG	ARG	PRO	PRO	ARG
THR	THR	ARG	CYS	GLY	VAL	VAL	GLY
THR	ARG	CYS	ASN	PRO	PRO	PRO	ALA
LEU	GLU	GLU	GLN	THR	THR	THR	ALA
GLU	PRO	PRO	LEU	SER	SER	SER	ARG
GLU	GLU	GLU	GLU	GLU	GLU	GLU	CYS
THR	ARG	TYR	ALA	GLY	VAL	VAL	THR
LYS	TYR	LYS	ALA	GLY	GLY	GLY	PRO
ALA	CYS	GLY	GLY	PRO	PRO	PRO	GLY
ARG	LEU	GLU	GLU	PRO	PRO	PRO	LEU
GLU	ALA	ASP	ASP	GLU	GLU	GLU	GLU
ARG	ARG	ARG	VAL	ARG	ARG	ARG	ARG
PRO	LEU	LEU	GLN	GLU	GLU	GLU	ALA
LYS	MET	GLY	GLN	GLY	ARG	ARG	PRO
LEU	ALA	ALA	LYS	ASP	ASP	ASP	PRO
ARG	ASP	ASP	ASN	CYS	CYS	CYS	R28
LYS	ALA	ALA	HIS	LEU	LEU	LEU	L37
ASP	LEU	ARG	TRP	PRO	ALA	ALA	A48
MET	ARG	GLY	GLN	ALA	ALA	ALA	A49
TYR	GLY	LYS	LYS	ILE	GLY	GLY	A50
LYS	VAL	VAL	ARG	ARG	SER	SER	ALA
MET	PRO	PRO	THR	THR	SER	SER	GLY
LEU	ALA	ALA	MET	HIS	HIS	GLY	GLY
ALA	PHE	HIS	ASN	GLN	LEU	LEU	PRO
VAL	HIS	ASN	ASN	GLN	GLN	GLN	ALA
ASP	GLY	GLY	LEU	LEU	GLN	GLN	ALA
PRO	VAL	PRO	PRO	PRO	PRO	ARG	ARG
GLU	VAL	VAL	VAL	ARG	ARG	GLY	GLY
ALA	GLU	GLU	ILE	ARG	ARG	ALA	LYS
PRO	PRO	ARG	SER	SER	LEU	LYS	VAL
THR	ASP	ASP	PHE	SER	SER	THR	ARG
GLU	GLY	GLY	LYS	SER	SER	SER	ARG
GLU	GLU	SER	LYS	LYS	VAL	GLY	GLY
HIS	TYR	ARG	ARG	VAL	VAL	GLN	GLN
ALA	LEU	GLN	TYR	SER	SER	VAL	VAL
GLN	LEU	LEU	TRP	THR	THR	ASN	PRO
ARG	LEU	GLN	VAL	GLY	SER	ASN	GLY
ALA	ASP	GLN	GLN	SER	SER	LEU	GLY
VAL	ASP	ASP	VAL	LEU	SER	LEU	LEU
THR	LEU	LEU	LEU	LEU	SER	SER	PRO
LYS	LEU	ALA	ALA	GLY	GLY	GLY	ALA
PRO	ASP	GLY	HIS	LEU	LEU	LEU	ALA
ARG	GLY	THR	THR	HIS	LEU	LEU	PRO
ARG	GLY	THR	GLY	THR	GLU	GLU	PRO
TYR	PHE	ASP	CYS	GLY	ASP	ASP	PRO
MET	ASP	THR	THR	THR	THR	THR	THR




- Molecule 2: Actin, alpha skeletal muscle

Chain H: 88% 11% .




- Molecule 2: Actin, alpha skeletal muscle

Chain I: 89% 9% .




- Molecule 2: Actin, alpha skeletal muscle

Chain J: 88% 10% .




- Molecule 3: Phalloidin

Chain X: 29% 43% 29%



- Molecule 3: Phalloidin

Chain Y: 29% 43% 29%



- Molecule 3: Phalloidin

Chain Z: 29% 57% 14%

P1	A2		A5	T6	C7
----	----	--	----	----	----

4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-166.86°, rise=27.58 Å, axial sym=C1	Depositor
Number of segments used	65917	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{Å}^2$)	60	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.036	Depositor
Minimum map value	-0.011	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.00166	Depositor
Map size (Å)	297.36, 297.36, 297.36	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.826, 0.826, 0.826	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, HIC, HYP, ADP, G5G, ALO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.08	0/167	0.20	0/223
1	B	0.08	0/167	0.26	0/223
1	C	0.08	0/167	0.18	0/223
1	D	0.07	0/167	0.15	0/223
1	E	0.07	0/167	0.16	0/223
2	F	0.11	0/2950	0.23	0/3994
2	G	0.11	0/2950	0.24	0/3994
2	H	0.11	0/2950	0.25	0/3994
2	I	0.12	0/2950	0.24	0/3994
2	J	0.11	0/2950	0.24	0/3994
3	X	0.06	0/29	0.14	0/36
3	Y	0.07	0/29	0.06	0/36
3	Z	0.06	0/29	0.09	0/36
All	All	0.11	0/15672	0.24	0/21193

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	167	177	176	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	167	177	176	2	0
1	C	167	177	176	1	0
1	D	167	177	176	1	0
1	E	167	177	176	1	0
2	F	2900	2874	2872	25	0
2	G	2900	2874	2872	23	0
2	H	2900	2874	2872	24	0
2	I	2900	2874	2872	22	0
2	J	2900	2874	2872	28	0
3	X	55	0	39	7	0
3	Y	55	0	37	7	0
3	Z	55	0	39	6	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
5	F	27	0	12	3	0
5	G	27	0	12	2	0
5	H	27	0	12	3	0
5	I	27	0	12	4	0
5	J	27	0	12	1	0
All	All	15640	15255	15415	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 128 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:157:ASP:OD1	5:G:402:ADP:O3'	1.98	0.82
2:I:157:ASP:OD1	5:I:402:ADP:O3'	1.96	0.81
2:H:323:SER:O	2:H:324:THR:OG1	1.98	0.81
2:G:336:LYS:NZ	5:G:402:ADP:N7	2.29	0.80
2:H:157:ASP:OD1	5:H:402:ADP:O3'	1.98	0.79

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	21/461 (5%)	21 (100%)	0	0	100	100
1	B	21/461 (5%)	20 (95%)	1 (5%)	0	100	100
1	C	21/461 (5%)	21 (100%)	0	0	100	100
1	D	21/461 (5%)	21 (100%)	0	0	100	100
1	E	21/461 (5%)	21 (100%)	0	0	100	100
2	F	368/377 (98%)	363 (99%)	5 (1%)	0	100	100
2	G	368/377 (98%)	362 (98%)	6 (2%)	0	100	100
2	H	368/377 (98%)	358 (97%)	10 (3%)	0	100	100
2	I	368/377 (98%)	362 (98%)	6 (2%)	0	100	100
2	J	368/377 (98%)	360 (98%)	8 (2%)	0	100	100
3	X	3/7 (43%)	1 (33%)	2 (67%)	0	100	100
3	Y	3/7 (43%)	1 (33%)	2 (67%)	0	100	100
3	Z	3/7 (43%)	1 (33%)	2 (67%)	0	100	100
All	All	1954/4211 (46%)	1912 (98%)	42 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	14/383 (4%)	14 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	14/383 (4%)	14 (100%)	0	100	100
1	C	14/383 (4%)	14 (100%)	0	100	100
1	D	14/383 (4%)	14 (100%)	0	100	100
1	E	14/383 (4%)	14 (100%)	0	100	100
2	F	313/319 (98%)	312 (100%)	1 (0%)	86	91
2	G	313/319 (98%)	313 (100%)	0	100	100
2	H	313/319 (98%)	313 (100%)	0	100	100
2	I	313/319 (98%)	313 (100%)	0	100	100
2	J	313/319 (98%)	313 (100%)	0	100	100
3	X	2/2 (100%)	2 (100%)	0	100	100
3	Y	2/2 (100%)	2 (100%)	0	100	100
3	Z	2/2 (100%)	2 (100%)	0	100	100
All	All	1641/3516 (47%)	1640 (100%)	1 (0%)	87	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	62	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	225	ASN
2	H	353	GLN
2	J	246	GLN
2	I	173	HIS
2	H	121	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

14 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HIC	G	73	2	10,11,12	1.54	2 (20%)	8,14,16	0.99	1 (12%)
2	HIC	F	73	2	10,11,12	1.54	2 (20%)	8,14,16	1.01	1 (12%)
2	HIC	H	73	2	10,11,12	1.55	2 (20%)	8,14,16	1.00	1 (12%)
3	HYP	X	1	3	6,8,9	0.97	0	5,10,12	1.56	1 (20%)
3	G5G	Z	4	3	8,9,10	0.94	0	5,12,14	1.15	0
3	ALO	Y	6	3	5,6,7	0.72	0	6,7,9	1.01	1 (16%)
3	HYP	Y	1	3	6,8,9	0.89	0	5,10,12	1.73	2 (40%)
3	G5G	X	4	3	8,9,10	0.95	0	5,12,14	1.16	0
3	G5G	Y	4	3	8,9,10	0.94	0	5,12,14	1.15	0
3	ALO	Z	6	3	5,6,7	0.71	0	6,7,9	1.02	1 (16%)
3	HYP	Z	1	3	6,8,9	1.02	0	5,10,12	1.70	1 (20%)
2	HIC	J	73	2	10,11,12	1.53	2 (20%)	8,14,16	1.08	1 (12%)
3	ALO	X	6	3	5,6,7	0.72	0	6,7,9	0.99	1 (16%)
2	HIC	I	73	2	10,11,12	1.54	2 (20%)	8,14,16	1.04	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HIC	G	73	2	-	0/5/6/8	0/1/1/1
2	HIC	F	73	2	-	0/5/6/8	0/1/1/1
2	HIC	H	73	2	-	0/5/6/8	0/1/1/1
3	HYP	X	1	3	-	0/0/11/13	0/1/1/1
3	G5G	Z	4	3	-	8/9/10/12	-
3	ALO	Y	6	3	-	3/5/6/8	-
3	HYP	Y	1	3	-	0/0/11/13	0/1/1/1
3	G5G	X	4	3	-	5/9/10/12	-
3	G5G	Y	4	3	-	5/9/10/12	-
3	ALO	Z	6	3	-	5/5/6/8	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HYP	Z	1	3	-	0/0/11/13	0/1/1/1
2	HIC	J	73	2	-	0/5/6/8	0/1/1/1
3	ALO	X	6	3	-	3/5/6/8	-
2	HIC	I	73	2	-	0/5/6/8	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	73	HIC	CD2-CG	3.03	1.41	1.36
2	G	73	HIC	CD2-CG	3.02	1.41	1.36
2	I	73	HIC	CD2-CG	3.02	1.41	1.36
2	F	73	HIC	CD2-CG	3.02	1.41	1.36
2	J	73	HIC	CD2-CG	2.98	1.41	1.36

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	73	HIC	CB-CA-C	-2.33	107.10	111.47
3	X	1	HYP	OD1-CG-CB	2.26	115.62	110.03
3	Z	1	HYP	OD1-CG-CB	2.22	115.52	110.03
3	Y	1	HYP	CG-CB-CA	2.21	106.76	103.96
2	I	73	HIC	CB-CA-C	-2.20	107.34	111.47

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	X	4	G5G	CA-CB-CG-O1
3	X	4	G5G	CA-CB-CG-CD1
3	X	4	G5G	N-CA-CB-CG
3	Y	4	G5G	CA-CB-CG-CD1
3	Y	4	G5G	N-CA-CB-CG

There are no ring outliers.

7 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	73	HIC	1	0
3	X	1	HYP	1	0
3	Y	6	ALO	2	0
3	Y	1	HYP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Z	6	ALO	3	0
2	J	73	HIC	1	0
3	X	6	ALO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	ADP	H	402	4	27,29,29	1.34	4 (14%)	42,45,45	1.92	11 (26%)
5	ADP	I	402	4	27,29,29	1.34	4 (14%)	42,45,45	1.93	11 (26%)
5	ADP	J	402	4	27,29,29	1.34	4 (14%)	42,45,45	1.94	10 (23%)
5	ADP	F	402	4	27,29,29	1.34	4 (14%)	42,45,45	1.94	11 (26%)
5	ADP	G	402	4	27,29,29	1.35	4 (14%)	42,45,45	1.92	11 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	H	402	4	-	6/16/32/32	0/3/3/3
5	ADP	I	402	4	-	5/16/32/32	0/3/3/3
5	ADP	J	402	4	-	5/16/32/32	0/3/3/3
5	ADP	F	402	4	-	5/16/32/32	0/3/3/3
5	ADP	G	402	4	-	7/16/32/32	0/3/3/3

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	402	ADP	C5-C4	4.43	1.47	1.39
5	H	402	ADP	C5-C4	4.41	1.47	1.39
5	I	402	ADP	C5-C4	4.39	1.47	1.39
5	J	402	ADP	C5-C4	4.38	1.47	1.39
5	F	402	ADP	C5-C4	4.38	1.47	1.39

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	402	ADP	C5-C4-N3	-6.20	118.66	126.75
5	H	402	ADP	C5-C4-N3	-6.15	118.72	126.75
5	J	402	ADP	C5-C4-N3	-6.10	118.79	126.75
5	F	402	ADP	C5-C4-N3	-6.07	118.83	126.75
5	I	402	ADP	C5-C4-N3	-6.07	118.83	126.75

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	402	ADP	C5'-O5'-PA-O2A
5	F	402	ADP	C5'-O5'-PA-O3A
5	F	402	ADP	O4'-C4'-C5'-O5'
5	F	402	ADP	C3'-C4'-C5'-O5'
5	G	402	ADP	C5'-O5'-PA-O3A



There are no ring outliers.



5 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	402	ADP	3	0
5	I	402	ADP	4	0
5	J	402	ADP	1	0
5	F	402	ADP	3	0
5	G	402	ADP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-53133. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 191




Y Index: 172




Z Index: 143


6.3.2 Raw map



X Index: 191



Y Index: 172




Z Index: 185


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00166. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 505 nm^3 ; this corresponds to an approximate mass of 456 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.337 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.337 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.97	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.39	4.13	3.48

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.39 differs from the reported value 2.97 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-53133 and PDB model 9Q GK. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.00166 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00166).

9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 99% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.00166) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9920	 0.5970
A	 0.8870	 0.5050
B	 0.9180	 0.4920
C	 0.9180	 0.5030
D	 0.9060	 0.4900
E	 0.9120	 0.4640
F	 0.9970	 0.6070
G	 0.9970	 0.6020
H	 0.9980	 0.6060
I	 0.9970	 0.6000
J	 0.9960	 0.6060
X	 0.9070	 0.4760
Y	 0.8890	 0.4990
Z	 0.8890	 0.4770

